

A proof of the Willmore conjecture

Martin U. Schmidt*

Institut für Theoretische Physik

Freie Universität Berlin

Arnimallee 14

D-14195 Berlin

email: martin.schmidt@physik.fu-berlin.de

*Supported by DFG, SFB 288 “Differentialgeometrie und Quantenphysik”

Contents

1	Introduction	3
1.1	The main result	3
1.2	A transformation of the variational problem	3
1.3	The solution of the transformed problem	9
2	The Bloch theory	14
2.1	The resolvent of the Dirac operator	15
2.2	The Bloch variety and the Fermi curve	25
2.3	Several reductions	29
2.4	Spectral projections	32
2.5	<i>Complex Fermi curves</i> of finite genus	42
2.5.1	The inverse problem	44
2.5.2	The Bäcklund transformations	47
2.5.3	A compactification of the <i>isospectral sets</i>	51
3	The moduli space	57
3.1	Variations of <i>complex Fermi curves</i>	57
3.2	Deformations of <i>complex Fermi curves</i>	68
3.3	The compactified moduli spaces with bounded genus	78
3.4	Limits of <i>complex Fermi curves</i>	91
3.4.1	Finite rank perturbations	92
3.4.2	Spectral theory of Dirac operators on the sphere	106
3.4.3	An illustrative family of potentials	112
3.4.4	The order of zeroes of spinors	120
3.4.5	The general case	129
3.5	The compactified moduli spaces	139
4	The <i>Generalized Willmore functional</i>	154
4.1	The <i>Weak Singularity condition</i>	154
4.2	Relative minimizers of the <i>Generalized Willmore functional</i>	157
4.3	Absolute minimizers of the <i>Generalized Willmore functional</i>	170
5	<i>Weierstraß potentials</i>	194
5.1	The <i>Singularity condition</i>	194
5.2	Variations of immersions	197
5.3	Constrained Willmore tori	201

1 Introduction

1.1 The main result

The Willmore functional associates to each immersion of the two-dimensional torus $\mathbb{T}^2$ into the three-dimensional Euclidean space $\mathbb{R}^3$ the integral $\int_{\mathbb{T}^2} H^2 d\mu$ of the square of the mean curvature of the induced metric with respect to the corresponding measure over the two-dimensional torus. This functional has an obvious extension to immersions of two-dimensional manifolds into higher-dimensional Riemannian manifolds. In this article we shall mainly consider immersions of $\mathbb{T}^2$ into $\mathbb{R}^3$ and sometimes into $\mathbb{R}^4$. This functional is invariant under conformal transformations of $\mathbb{R}^3$. The Willmore conjecture states that up to conformal transformations this functional has a unique absolute minimum. The corresponding immersion is the *Clifford torus*, which is the surface of a rotated circle of radius r around some axis in the plane spanned by the circle with distance $\sqrt{2}r$ from the center of the circle. The corresponding Willmore functional is equal to $2\pi^2$.

All smooth metrics on $\mathbb{T}^2$ are conformally equivalent to a flat metric and each flat metric corresponds to exactly one complex structure on $\mathbb{T}^2$. Hence, the conformal classes of these immersions are equal to the real two-dimensional manifold $\mathcal{M}_1$ of moduli of compact Riemann surfaces of genus one. Consequently, the Willmore functional may be restricted to all immersions of some given conformal class, and the infima of these restrictions fit together to some function on this moduli space $\mathcal{M}_1$. The main result of this article is the calculation of another function on $\mathcal{M}_1$, which yields a lower bound of the former function. Furthermore, this latter function has a unique minimum at the conformal class corresponding to the flat torus $\mathbb{R}^2/\mathbb{Z}^2$. Finally, for all rectangular conformal classes this lower bound is sharp and the corresponding immersions are unique up to conformal transformations of $\mathbb{R}^3$. The minimum is realized by the *Clifford torus*. We conjecture that the latter function describes the minimum of the restriction of the Willmore functional to conformal immersions of two-dimensional tori $\mathbb{T}^2$ into $\mathbb{R}^4$ in dependence of the conformal class of $\mathbb{T}^2$.

We now outline the strategy of the proof. It contains

1.2 A transformation of the variational problem

Usually the determination of the minimum splits into two problems:

Proof of the existence of a minimizer.

Determination of the lowest critical point.

As part of the *proof of the existence* we shall specify the class of objects (immersions) we consider. In view of the first problem this class should be chosen large enough in order to contain enough relative minimizers. But in view of the second problem this class should be chosen small enough in order to simplify the classification. For the Willmore functional both problems seem to be difficult to solve.

The first problem was solved by L. Simon [Si-1, Si-2] using geometric measure theory. With his methods, however, it seems hopeless to prove or disprove the existence of a minimizer for the restrictions to arbitrary conformal classes.

The critical points of the Willmore functional are called Willmore tori, and the critical points of the restrictions to some conformal classes are called *Constrained Willmore tori* (the precise definition is given in *Constrained Willmore tori* 5.8). The stereographic projections of minimal tori in S^3 are Willmore tori, and the area of these minimal tori is equal to the Willmore functional. In spite of a quite satisfactory description of the minimal tori in S^3 given in [Hi] it was not clear which minimal torus corresponds to the lowest value of the Willmore functional. U. Pinkall was the first to construct Willmore tori which are not stereographic projections of minimal tori in S^3 [Pi-1]. Afterwards many other Willmore tori were constructed [Ba, B-B], indicating that a general classification of the lowest critical points seems to be even more difficult.

Thirdly, there exist several attempts to pass around these two problems. In this direction first we mention the topological lower bound 4π (compare with [Wi-2, Section 7.6] and references therein), which is sharp for immersions from S^2 into $\mathbb{R}^3$. Furthermore, in [L-Y] P. Li and S. T. Yau proved several lower bounds of the Willmore functional, one of which implies that all immersions, whose conformal classes belong to a closed subset including the conformal class of the *Clifford torus*, have a Willmore functional larger or equal to $2\pi^2$.

We want to present a new approach to both problems. The starting point is an observation due to I. Taimanov and which initiated this project. It states that the local Weierstraß representation [Ei] of immersions of surfaces in $\mathbb{R}^3$ can be extended to a global Weierstraß representation. More precisely, for all conformal immersions of some flat torus $\mathbb{T}^2$ into $\mathbb{R}^3$, there exists a unique real potential U on $\mathbb{T}^2$ and some spinor $\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$ in the kernel of the Dirac operator $\begin{pmatrix} U & \partial \\ -\bar{\partial} & U \end{pmatrix}$ on $\mathbb{T}^2$ with potential U , such that the differential of the immersion is the $\mathbb{R}^3$ -valued one form $(\Re(\psi_1^2 dz - \psi_2^2 d\bar{z}), \Im(\psi_1^2 dz - \psi_2^2 d\bar{z}), \psi_1 \bar{\psi}_2 dz + \bar{\psi}_1 \psi_2 d\bar{z})$ on $\mathbb{T}^2$ [Ta-1, Fr-2]. With respect to the spinor this $\mathbb{R}^3$ -valued one-form is a quadratic form. Therefore, it is single-valued on $\mathbb{T}^2$. We should remark that these spinors $\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$ are in general not single-valued functions on $\mathbb{T}^2$. The non-uniqueness, however, is soft in the sense that the difference is only given by the factor -1 . More precisely, the universal covering $\mathbb{R}^2$ is a principal bundle over $\mathbb{T}^2$ whose structure group is the fundamental group of $\mathbb{T}^2$, which is isomorphic to $\mathbb{Z}^2$. Any homomorphism of this group into the group $\{1, -1\} \simeq \mathbb{Z}_2$, induces some line bundle on $\mathbb{T}^2$. The spinor bundle is then given by this line bundle tensored with the trivial vector bundle of rank two over $\mathbb{T}^2$. Obviously there exist four different homomorphisms from $\mathbb{Z}^2$ into $\mathbb{Z}_2$, and therefore also four different spin structures on $\mathbb{T}^2$. The spin structures of the global Weierstraß representations are obviously homotopy invariants of the corresponding immersions, and in [Pi-2] it was shown that they are the only homotopy invariants.

This global Weierstraß representation was generalized to immersions into $\mathbb{R}^4$ by F. Pedit and U. Pinkall [P-P]. For this purpose generalized Dirac operators with complex potentials were considered. They act on the trivial vector bundle of rank two over $\mathbb{T}^2$ tensored with the line bundles induced by a real character of the fundamental group of $\mathbb{T}^2$. These bundles may be considered as quaternionic line bundles

Conversely, for all real potentials, whose Dirac operators have a non-empty kernel, the corresponding quadratic form of any element of this kernel is a closed $\mathbb{R}^3$ -valued one-form on the torus $\mathbb{T}^2$. The integral of this one-form yields a mapping from the universal covering $\mathbb{R}^2$ of $\mathbb{T}^2$ into $\mathbb{R}^3$. If the spinor is everywhere non-vanishing, this mapping is a conformal immersion. Moreover, if this mapping is the composition of the universal covering map from $\mathbb{R}^2$ to $\mathbb{T}^2$ with some mapping from $\mathbb{T}^2$ into $\mathbb{R}^3$, or equivalently if the integrals of the $\mathbb{R}^3$ -valued one-form over all cycles of $\mathbb{T}^2$ vanish, then the corresponding Willmore functional is equal to the square of the L^2 -norm of the potential times four.

Periodicity condition 1.1. *The integrals of the two closed one forms $\psi_1^2 dz - \psi_2^2 d\bar{z}$ and $\psi_1 \bar{\psi}_2 dz + \bar{\psi}_1 \psi_2 d\bar{z}$ along all cycles of $\mathbb{T}^2$ vanish.*

An easy calculation (compare with Corollary 2.16 and Lemma 2.22) shows that this condition on any spinor $\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$ on $\mathbb{R}^2$ in the kernel of the Dirac operator $\begin{pmatrix} U & \partial \\ -\bar{\partial} & U \end{pmatrix}$ has another

Equivalent form of the Periodicity condition 1.2.

$$\int_{\mathbb{T}^2} \psi_1^2(x) d^2x = 0, \int_{\mathbb{T}^2} \psi_2^2(x) d^2x = 0 \text{ and } \int_{\mathbb{T}^2} \psi_1(x) \bar{\psi}_2(x) d^2x = 0.$$

Therefore, we may transform the variational problem of the restriction of the Willmore functional to some conformal class into the variational problem of the square of the L^2 -norm times four on the space of all real potentials on $\mathbb{T}^2$, whose Dirac operator has some non-trivial spinor in the kernel satisfying the *Periodicity condition* 1.1. The conformal classes of the corresponding immersions are all equal to the conformal class of the flat torus $\mathbb{T}^2$. For all L^2 -potentials the resulting Dirac operator defines an operator with compact resolvent on an appropriate Hilbert space of spinors on $\mathbb{T}^2$. For all spinors in the kernel of this Dirac operator both equivalent forms of the *Periodicity condition* 1.1 are well defined. In fact, we shall see in Remark 2.7 that the spinors in the kernel belong to the Sobolev spaces $W^{1,p}(\mathbb{T}^2) \times W^{1,p}(\mathbb{T}^2)$ with $1 < p < 2$. Due to the Sobolev embedding theorem [Ad, 5.4 Theorem] this implies that the restrictions of the spinors to a one-dimensional plane in $\mathbb{T}^2$ are L^2 -spinors. In this sense the *Periodicity condition* 1.1 is well defined. The *Equivalent form* 1.2 is well defined since these spinors in the kernel of the Dirac operator are L^2 -spinors. Therefore, it is natural to extend the Willmore functional (defined by the square of the L^2 -norm of the potential times four) to the space of all mappings from $\mathbb{T}^2$ into $\mathbb{R}^3$, defined as the integrals over the $\mathbb{R}^3$ -valued one-forms corresponding to some spinor in the kernel of the Dirac operator fulfilling the *Periodicity condition* 1.1. These mappings are always continuous, but in general not smooth. Furthermore, they contain also smooth mappings, which are not immersions, since the images have branch points. We consider these mappings to be the maximal domain of definition of the Willmore functional and therefore make the definitions

Weierstraß potentials 1.3. *Real potentials $U \in L^2(\mathbb{T}^2)$, whose Dirac operators contain in their kernel some non-trivial spinor fulfilling the *Periodicity condition* 1.1, are called Weierstraß potentials.*

The main achievement of the observation of I. Taimanov is the description of the space of conformal immersions from a flat torus $\mathbb{T}^2$ into $\mathbb{R}^3$ as a subspace of the phase space of the integrable system, whose Lax operator is the Dirac operator with a pair of potentials on $\mathbb{T}^2$. Consequently, we use the methods from the theory of integrable systems related to Lax operators and their spectral curves. The corresponding evolution is described by the Lax equation $\frac{\partial L}{\partial t} = [L, A]$ of a Lax pair (L, A) . This equation indicates that the spectrum of the Lax operator L is an integral of motion. Roughly speaking, the eigenvalues of the Lax operator fits together to all integrals of motion of a completely integrable system [D-K-N, Kon]. The application of these methods to differential geometry accompanied their development almost from the very beginning. But in contrast to other geometric applications the present approach uses these methods in order to describe the whole space of immersions, and not only some special immersions (Willmore tori, CMC-tori, minimal tori etc.). This makes the interplay between Hamiltonian systems and variational analysis available for this problem. Since Hamiltonian mechanics and variational analysis have as common origin Lagrangian mechanics, it seems natural to apply the new methods of the former theory to an unsolved problem of the latter.

These Dirac operators with potentials are the Lax operators of the Davey–Stewartson equation [Kon], and the corresponding spectral curves (i. e. a complete set of integrals of motion) are the *complex Fermi curves* of these Dirac operators corresponding to vanishing energy. These *complex Fermi curves* are the subsets of all complex characters of the fundamental group of $\mathbb{T}^2$ such that the Dirac operators $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ acting on the trivial vector bundle of rank two tensored with the line bundle induced by the character have a non-trivial kernel. The Bloch theory investigates the common spectrum of a periodic differential operator with the commuting space translations [Ku]. For three-dimensional periodic Schrödinger operators this common spectrum is called band structure. The physical *Fermi curves* are the intersection of the band structure with a fixed energy level (compare with [F-K-T, §18.-§20.]). Our *complex Fermi curves* are the complex continuations of the analogs for two-dimensional Dirac operators with periodic potentials instead of three-dimensional periodic Schrödinger operators.

Weierstraß curves 1.4. *The complex Fermi curves of the Weierstraß potentials are called Weierstraß curves. (This terminology was suggested by I. Taimanov.)*

The linear involution $(V, W) \mapsto (W, V)$ and the anti-linear involution $(V, W) \mapsto (\bar{V}, \bar{W})$ induce two involutions on the space of *complex Fermi curves*. Therefore, the fixed points of these involutions (i. e. the space of real potentials U) is some reduction of this integrable system, and called the modified Novikov–Veselov equation, since it is a modification of the Novikov–Veselov equation analogous to the modified Korteweg–de Vries equation [Kon]. The corresponding *complex Fermi curves* have a holomorphic and an anti-holomorphic involution. Unfortunately, the reduction corresponding to the linear involution destroys the Hamiltonian structure of this integrable system. This is because the natural symplectic form vanishes on the fixed point set of this involution (compare with Remark 3.3). For this reason we shall consider the unreduced system. Moreover, due to the generalization of the global Weierstraß representation to immersions into $\mathbb{R}^4$, the corresponding *complex Fermi curves* are invari-

ant only under the anti-holomorphic involution, but in general not under the holomorphic involution. In [G-S-2] it was proven that the *Weierstraß curves* 1.4 are invariant under conformal transformations of $\mathbb{R}^3$ acting on the corresponding immersions. Finally, the Willmore functional is the *first integral* of this integrable system.

In a second step we will investigate the *Periodicity condition* 1.1. I. Taimanov conjectured at the beginning of this project that these *Weierstraß curves* 1.4 are conformal invariants and he confirmed this numerically. Usually, the real parts of the *isospectral sets* (i. e. all potentials fulfilling some reality condition, whose spectral curves are equal to some given spectral curve) are compact tori. On the other hand, the set of all potentials corresponding to all conformal transformations of a given immersion is in general not compact. For this reason the subset of some *isospectral set* containing all potentials fulfilling the *Periodicity condition* 1.1 seems not to be closed. But he proved that the isospectral flows preserve this subset [Ta-1, Ta-2]. This apparent inconsistency was the first hint that all *Weierstraß curves* 1.4 have some non-trivial singularity. In fact, in [Sch] it was shown that for singular spectral curves the real parts of the *isospectral sets* are compact, but decompose into a union of non-closed subsets of different dimensions which are invariant under the isospectral flows. We should remark that all *complex Fermi curves* corresponding to Dirac operators with non-trivial kernel have some kind of singularity. In fact, each character corresponding to some spin structure (which takes values in $\{1, -1\}$) is a fixed point of the anti-holomorphic involution. Since this involution may not have fixed points on the normalization of the *complex Fermi curve*, the preimage of these elements of the *complex Fermi curves* under the normalization map has to contain at least two elements. But in general these singularities do not give rise to non compact components of the real part of the *isospectral sets*.

Hence the characterization of *Weierstraß potentials* 1.3 refers to the characterization of *Weierstraß curves* 1.4. But we emphasize that not all potentials, whose *complex Fermi curves* are *Weierstraß curves* 1.4, are *Weierstraß potentials* 1.3. The *Weierstraß curves* 1.4 may be characterized as those *complex Fermi curves*, which obey the *Singularity condition* 5.3. This *Singularity condition* 5.3 assumes, roughly speaking, that the *complex Fermi curve* has some fourth-order multiple point. The exact form of this condition involves the first derivatives of the regular functions. There exists a natural simplified and weaker condition called

Weak Singularity condition 1.5. (*See also 4.2.*) *The complex Fermi curve contains a singular point corresponding to characters with values in $\{1, -1\}$, whose preimage in the normalization contains either at least four points, or two points, which are zeroes of both quasi-momenta (i. e. the logarithmic derivatives of the values of the characters at two generators of the fundamental group of $\mathbb{T}^2$).*

For immersions into $\mathbb{R}^4$ ‘quaternionic function theory’ developed by F. Pedit and U. Pinkall [P-P, B-F-L-P-P, F-L-P-P] leads to the

Quaternionic Singularity condition 1.6. *The complex Fermi curve contains a singular point corresponding to real characters (i. e. quaternionic line bundles), whose preimage in the normalization contains either at least four different points, or at least two different points, at which both quasi-momenta vanish.*

This *Quaternionic Singularity condition* 1.6 is equivalent to the condition that the corresponding *isospectral set* contains some complex potential, whose Dirac operator acting on some quaternionic line bundle (i. e. the trivial vector bundle of rank two on $\mathbb{T}^2$ tensored with the line bundle induced by some real character of the fundamental group of $\mathbb{T}^2$) has a kernel of quaternionic dimension larger than one. In general the quaternionic line bundles may have non-vanishing degree. In this case the action of the Dirac operator has to be modified. But we shall see that the notion of *complex Fermi curves* generalizes to these cases, and the construction of the *Baker–Akhiezer function* shows that at least in the finite genus case the *Quaternionic Singularity condition* 1.6 is equivalent to the condition that the *isospectral set* contains some Dirac operator acting on some quaternionic line bundle with at least two-dimensional kernel.

The *Weak Singularity condition* 1.5 is equivalent to the condition that some *complex Fermi curve* endowed with a holomorphic and an anti-holomorphic involution obeys the *Quaternionic Singularity condition* 1.6, and that the singularity is invariant under the holomorphic involution. In particular, all *complex Fermi curves* obeying the *Weak Singularity condition* 1.5 correspond to some not necessarily injective conformal mappings from $\mathbb{T}^2$ into $\mathbb{R}^4$.

The characterization of the *Weierstraß potentials* 1.3 in terms of their *complex Fermi curves* suggests to consider the slightly larger set of all potentials, whose *complex Fermi curves* satisfy the *Singularity condition* 5.3. Instead of this set we now introduce the even larger set of all

Generalized Weierstraß potentials 1.7. *Real potentials $U \in L^2(\mathbb{T}^2)$, whose complex Fermi curves fulfill the Weak Singularity condition 1.5, are called Generalized Weierstraß potentials.*

The set of all *Generalized Weierstraß potentials* is a subvariety of the Hilbert space of all L^2 -potentials. More precisely, this set is locally the zero set of at most ten analytic functions. Unfortunately, these functions are not weakly continuous. But the restriction of these functions to small balls are weakly continuous. Indeed, we shall explicitly see that these functions are not weakly continuous on balls, whose radius correspond to Willmore functionals larger than 4π . More precisely, the moduli space of all *complex Fermi curves* is not complete. There exist limits of *complex Fermi curves*, whose *first integrals* are equal to 4π , but which correspond to no real (or complex) potential. Nevertheless we shall call these limits *complex Fermi curves*, since they are the *complex Fermi curves* of *finite rank perturbations* of the Dirac operators. Moreover, they may be considered as *complex Fermi curves* of Dirac operators acting on line bundles of non-vanishing degree, which occur in ‘quaternionic function theory’ [P-P, B-F-L-P-P, F-L-P-P].

For this reason we transform in a last step the variational problem of the L^2 -norm on the space of all *Generalized Weierstraß potentials* 1.7 to a variational problem of the natural extension of the *first integral* to all

Generalized Weierstraß curves 4.4. *Roughly speaking, these are those complex Fermi curves of finite rank perturbations of Dirac operators with real potentials $U \in L^2(\mathbb{T}^2)$, which obey the Weak Singularity condition 1.5.*

To sum up we have the following chain of modifications of variational problems starting with the Willmore functional:

1. **Immersions.** On the space of smooth immersions from $\mathbb{T}^2$ into $\mathbb{R}^3$ we are given the Willmore functional. The space is an open subset of the vector space of smooth mappings from $\mathbb{T}^2$ into $\mathbb{R}^3$. But the functional is quite complicated involving second-order partial derivatives.
2. **Weierstraß potentials.** Consider the subspace of all L^2 -potentials, whose Dirac operators contain in their kernel some spinor fulfilling the *Periodicity condition* 1.1, and on this subspace the functional, which is the square of the L^2 -norm times four. This space is quite complicated, but the functional is very simple.
3. **Generalized Weierstraß potentials.** Consider the real subspace of all L^2 -potentials, whose *complex Fermi curves* fulfill the *Weak Singularity condition* 1.5, and on this subspace again the functional, which is the square of the L^2 -norm times four. The space is a real subvariety of L^2 . The functional is again very simple. The corresponding infimum yields only a lower bound for the Willmore functional.
4. **Generalized Weierstraß curves.** Consider the subspace of the completion of the moduli space containing all *complex Fermi curves* fulfilling the *Weak Singularity condition* 1.5, and the functional, which is the restriction of the *first integral*. The corresponding infimum yields only a lower bound of the Willmore functional.

In a concluding remark we emphasize the motivation of the modification of the second problem into the third problem. From the point of view of ‘quaternionic function theory’ it seems to be more natural to consider the variational problem of the Willmore functional on immersions into $\mathbb{R}^4$ instead of $\mathbb{R}^3$. Comparing these two problems the latter has the advantage of providing a smaller space, and the former has the advantage of a stronger condition on relative minimizers. In some sense the *Weak Singularity condition* 1.5 uses this advantage of immersions into $\mathbb{R}^4$ without giving up the advantage of immersions into $\mathbb{R}^3$. In fact, we shall provide some conditions on relative minimizers of the fourth variational problem, which singles out many Willmore tori (in particular all Willmore tori described in [B-B]).

1.3 The solution of the transformed problem

Finally, we arrive at a variational problem of the *first integral* of an integrable system related to the two-dimensional Dirac operator with constraints on the corresponding spectral curves. Therefore, our main effort aims at investigating the moduli space, i. e. the space of all generalized *complex Fermi curves*. Roughly speaking, we are interested in two properties:

Compactness of the space of all *complex Fermi curves*, whose *first integral* is bounded from above, and a

Differentiable structure of the moduli space.

The first property contains the essential part of the *Proof of the existence of a minimizer*. The second property provides the main tools for the *Classification of the lowest critical points*. Both properties are proven in full generality.

For the *Compactness* we have to endow this moduli space with some topology. We shall use the topology of the Hausdorff metric on the space of all closed subsets of the one-point-compactification of $\mathbb{C}^2$. With this topology the corresponding space of closed subsets of the one-point-compactification of $\mathbb{C}^2$ becomes a compact Hausdorff space. So in order to obtain compactness of some subset of the moduli space, we only have to prove that this subset is closed. First we shall prove this for the subsets of the moduli space containing all generalized *complex Fermi curves* of bounded geometric genus and with bounded *first integral*. The main technical tools are a representation of the *complex Fermi curves* as two complex planes glued along several cuts and the evaluation of the *first integral* by some Dirichlet integral of some linear combination of the quasi-momenta over one of these planes. These tools should be considered as generalizations of analogous tools for the KdV equations, which were first used by V. A. Marchenko and I. V. Ostrowski [M-O] in their investigation of the moduli space corresponding to real periodic solutions of the KdV equation with the help of conformal mappings. In the present situation we encounter two additional technical complications.

1. Two-dimensionality of the system: Integrable systems having Lax operators may be divided into three groups.

Two trivial flows. These systems are finite-dimensional. The Lax operator is a spectral-parameter dependent matrix, and the spectral curves have two meromorphic functions with prescribed poles.

One trivial and one periodic flow. The phase space of these systems is a space of functions depending on one variable. The Lax operator is some ordinary differential operator. The associated spectral curves have one meromorphic function and one transcendental function with prescribed singularities.

Two periodic flows. The phase space of these systems is a space of functions depending on two variables. The Lax operator is some partial differential operator and the associated spectral curves have two transcendental functions with prescribed singularities.

The KdV and the NLS equation belong to the second group. In particular the corresponding spectral curves are hyperelliptic. In this case there exist natural parameters of the moduli space, which were introduced in [M-O] and studied further in [G-S-1, Tk]. The integrable system corresponding to the two-dimensional Dirac operator belongs to the third group. In this case natural parameters of the moduli space do not exist. We overcome this problem by considering the moduli space as a closed subspace of some higher-dimensional space, which has natural parameters.

2. A complicated reality condition: Some reality conditions imply that the corresponding spectral curves are *M-curves* (i. e. the real part has $g + 1$ connected components, where g is the geometric genus of the spectral curve). In this case the real cycles cut

the spectral curves into pieces, which may be described by some conformal mappings. The KdV equation has this property, but the integrable system corresponding to the two-dimensional Dirac operator does not. It might also be that the *complex Fermi curves* can be divided into pieces, which can be described by conformal mappings, but there does not exist a natural choice. This difficulty is overcome by dividing the *complex Fermi curve* again into two pieces, where we make use of holomorphic functions instead of univalent (or schlicht) functions.

In a second and independent step we determine the limits of the resolvents of a weakly convergent sequence of potentials. It is contained in the technically most involved Section 3.4 and is motivated by two results on the spectral theory of Dirac operators in Section 2.1 and on the compactifications of isospectral sets of *complex Fermi curves* of finite *geometric genus* in Section 2.5.3. The first result states that the resolvents are almost weakly continuous with respect to the potentials. More precisely, if for any $\varepsilon > 0$ the L^2 -norms of the restrictions to all ε -balls of a weakly convergent sequence of potentials is bounded by the inverse Sobolev constant, then the limit of the corresponding sequence of resolvents is the resolvent of the limit. Therefore, in general the limits of the resolvents of a weakly convergent sequence of potentials U_n should differ from the resolvent of the limit only in those points where the limit of the corresponding measures $U_n(x)\bar{U}_n(x)d^2x$ has a point measure whose mass is not smaller than the inverse of the square of the Sobolev constant. The second result states that the compactified isospectral sets of *complex Fermi curves* of finite *geometric genus* contain potentials on the complement of a finite set of the torus (i. e. the corresponding *Baker Akhiezer functions* are not eigenfunctions in small neighbourhoods of this finite set). The corresponding *first integrals* are equal to four times the square of the L^2 -norms of these potentials plus a multiple of 4π . In particular, the isospectral sets are compact, if the *first integral* is smaller than 4π . This follows also from the first result with the optimal Sobolev constant (compare with Remark 2.5).

Therefore, it seems natural to expect that in general the limits of resolvents of weakly convergent sequences of potentials are resolvents of perturbations of the Dirac operator corresponding to the weak limit. More precisely, the support of the perturbation is contained in a finite subset of $\mathbb{T}^2$ and therefore should be described by a finite linear combination of such distributions. This is proven by some kind of blowing up of the sequences of potentials nearby the singular points to potentials on $\mathbb{P}^1$. These finite rank perturbations of Dirac operators can be considered as holomorphic structures of quaternionic line bundles of non-vanishing degree, which occur in ‘quaternionic function theory’ [P-P, B-F-L-P-P, F-L-P-P].

As a consequence, we prove in Corollary 5.11 the existence of minimizers of the restrictions of the Willmore functional to all conformal classes. We remark that the application of the Heine–Borel property [Str, Chapter I. 1.1 Theorem] in this variational problem is possible, since the transformation to a variational problem on the *complex Fermi curves* throws out the non-compact conformal symmetry. Furthermore, the Willmore functional was extended to the maximal domain of definition.

The main techniques for establishing the *Differentiable structure* is an application of deformation theory of compact complex spaces combined with some general features of integrable systems having Lax operators. The spectral curves of these integrable systems should be

considered as a combination of all integrals. Moreover, the differentials of these integrals fit together to some linear map from the tangent space of the phase space into the space of holomorphic one-forms of the corresponding spectral curve. On the other hand, the real part of the connected component of the identity of the Picard group acts freely and transitively on the *isospectral sets* (i. e. the set of those elements of the phase space, whose spectral curves are equal to some given spectral curve). The differential of this action yields some mapping from the Lie algebra of the Picard group into the tangent space. On compact Riemann surfaces Serre duality establishes a canonical pairing between the Lie algebra of the Picard group and the space of holomorphic one-forms. On the other hand, the symplectic form gives a pairing on the tangent space of the phase space. Due to a general rule for such integrable systems these two mappings are dual to each other with respect to the pairings given by the symplectic form and Serre duality. As a starting point we establish this structure for the integrable system, whose Lax operator is the two-dimensional Dirac operator. Since both pairings (symplectic form and Serre duality) are non-degenerate, we conclude that the tangent space of the moduli space may be identified with the space of regular forms of the spectral curves. Afterwards we apply a theorem of Grauert, which is the base of the generalization of the deformation theory of compact complex manifolds initiated by K. Kodaira and D. C. Spencer [Ko] to compact complex spaces (see [G-P-R, Chapter III. §4.]). This application shows that some special subsets of the moduli space containing only *complex Fermi curves* of bounded geometric genus are complex manifolds. All *complex Fermi curves* of finite geometric genus are contained in some of these subsets.

In a second step we extend this deformation theory to all *complex Fermi curves*. For this purpose we show that all *complex Fermi curves* are arbitrary small perturbations of *complex Fermi curves* of finite *geometric genus*, and as a consequence we may extend the deformation theory to these perturbations. The essential tool of this step is the *asymptotic analysis* of *complex Fermi curves*. Usually this is done only for a restricted class of potentials. More precisely, since the mapping from the potentials to the spectral curves (i. e. the combined integrals of motion) is a non-linear perturbation of the mapping to the absolute values of the Fourier coefficients, differentiability assumptions on the potentials lead to decreasing Fourier coefficients and a simplification of the asymptotic analysis (compare with [F-K-T]). We shall establish the asymptotic analysis in Theorem 2.12 and Proposition 3.52 for all L^2 -potentials. As a consequence in Theorem 3.57 the moduli space is shown to be a Banach manifold modelled on Banach space ℓ_1 . For short we use the notation ℓ_1 -manifold. The corresponding topology differs from the topology used for the *Compactness*.

The *Differentiable structure* allows us to prove that all relative minimizers of the fourth variational problem have dividing real parts (i. e. the real part divides the normalization of the *complex Fermi curve* into two connected components). Here we make use of local one-dimensional deformation families, which for negative deformation parameters violate the reality condition. More precisely, the corresponding anti-holomorphic involutions have fixed points on the normalization of the *complex Fermi curves*. Therefore, the corresponding first-order derivative should correspond to a higher-order derivative of deformations of immersions. For relative minimizers the derivative of the *first integral* with respect to the deformation parameter has to be non-negative. This condition can be used with the help of a

calculation of intersection numbers of the real part to show that either this real part divides the normalization or it has self intersecting points. In the second case the corresponding double points can again be used to construct local deformations with decreasing *first integral*. To sum up, the dividing property of the real part is a consequence of the positivity of some derivative of even-order, and is definitely not fulfilled at all critical points. We remark that this dividing property of relative minimizers is not true for the original Willmore functional. Roughly speaking, the simplification of the *Weak Singularity condition* 1.5 results in a smaller set of relative minimizers comparable to the smaller set of stereographic projections of minimal tori in S^3 .

Finally, we investigate the space of all *Generalized Weierstraß curves* 4.4 with dividing real parts and with the *first integral* being smaller than 8π . Fortunately, this space is small enough to allow a classification of all relative minimizers in this space. Since the *geometric genera* of all *complex Fermi curves* with dividing real parts are bounded in terms of the *first integral*, this space is finite-dimensional. We shall first show that any *complex Fermi curve* of this space may be deformed within this space into a *Generalized Weierstraß curve* 4.4, whose normalization has two connected components and whose *first integral* is equal to 8π . One connected component of these *Generalized Weierstraß curves* 4.4 has to be the spectral curve of some elliptic solution of the Kadomtsev–Petviashvili Equation. Moreover, due to an estimate on the genus in terms of the *first integral*, which follows from a formula of Krichever [Kr-2], we will classify these *Generalized Weierstraß curves* 4.4. Finally, this classification is extended to the whole deformation family. As a result for each conformal class we have for all of the three non-trivial spin structures exactly one relative minimizer of the fourth variational problem, whose *first integrals* are not larger than 8π . This agrees with the identification of the homotopy invariants with the spin structures and a result of P. Li and S. T. Yau [L-Y], which implies that all immersions corresponding to the trivial spin structure have a self intersection, and therefore a Willmore functional larger than 8π (compare with Remark 4.23). The *Generalized Weierstraß curves* 4.4 of these relative minimizers fit together into one family of compact Riemann surfaces whose geometric genera vary between zero and two. Due to a special feature of this family the absolute minimum is attained for a *Generalized Weierstraß curve* 4.4 of geometric genus zero. The *Generalized Weierstraß curves* 4.4 of geometric genus zero and one are calculated explicitly, and the absolute minimum is attained for the *complex Fermi curve* of the *Clifford torus*.

Finally, we give a short summary of the content of the main sections. Section 2 contains the spectral theory of the Dirac operator with periodic L^2 -potentials on a flat torus. First we reproduce in the present situation several classical result of Bloch theory. We show in Section 2.1 that these Dirac operators are direct integrals of families of operators with compact resolvents. Section 2.2 introduces the complex Bloch variety and the *complex Fermi curve* together with the first steps of the asymptotic analysis of these varieties. After considering several symmetries and reality conditions in Section 2.3 we present in Section 2.4 a remarkable relation of different representations of these Dirac operators and the corresponding spectral projections. The *Singularity conditions* (Section 4.1 as well as Section 5.1) and the generalized Cauchy's integral formula (Section 3.4.1) are based on this structure. In the last Subsection 2.5 we leave for a moment the main road to the proof and apply some

techniques of the ‘finite gap’ theory of integrable systems. This discussion motivates the complicated theorems in the following Section 3.

Section 3 essentially contains the proof of the *Differentiable structure* of the moduli space and the *Compactness*. The former property is proven in three steps in Section 3.1–3.2 and Section 3.5. In the first step the tangent mapping of the phase space to the spectral curves and its dual (with respect to the symplectic form and Serre duality) is constructed. In the second step an application of Deformation theory establishes the *Differentiable structure* for the moduli spaces with bounded genus. Finally, the asymptotic analysis extends this to the whole moduli space. The latter property is proven first for the moduli spaces with bounded genus (Section 3.3) and independently in the most involved Section 3.4.

After these preparations the proof of the Willmore conjecture is contained in Section 4. In Section 4.1 we prove that all *Weierstraß curves* 1.4 satisfy the *Weak Singularity condition* 1.5. Then the dividing property of relative minimizers follows in Section 4.2. Finally the classification of all local minimizers below 8π is presented in Section 4.3.

The last Section contains an investigation of the *Singularity condition* 5.3 (Section 5.1) and a proof that all conformal classes contain a minimum of finite type (Section 5.3).

Whenever possible we have tried to quote new standard textbooks for the tools we have used. These textbooks contain the references to the original articles.

Acknowledgements: This work is dedicated to my teacher Robert Schrader. His constant support gave me the opportunity to attack such a difficult problem without the safety of having a permanent position. In the beginning, Iskander Taimanov participated in this project and later accompanied it with several discussions. The collaboration inside the SFB 288 with the mathematicians Alexander Bobenko, Frank Duzaar, Daniel Grieser, Frank Hausser, Franz Pedit and Ulrich Pinkall was very stimulating and helpful. So results from ‘quaternionic function theory’ provided useful checks and led to new ideas. During several visits in Berlin Piotr Grinevich was a good adviser. Elmar Vogt helped to recognize an impasse. David Jerison pointed out to the author, that the arguments of [Wo, Proposition 2.6], where the analogous but weaker Carleman inequality for the gradient term of the Laplace operator is treated, carry over to the Dirac operator. Sonya Miller, Robert Schrader and Elmar Vogt helped to improved my English style.

2 The Bloch theory

Originally the Bloch theory investigates the spectral theory of a Schrödinger operator with a periodic potential. Bearing in mind that the Dirac operator is a natural relativistic modification of the Schrödinger operator we continue to use this notion of a Bloch theory in the context of Dirac operators with periodic potentials. The resolvent of an unbounded operator is a bounded operator, which contains all relevant information of the spectral theory of the former operator. Hence we first introduce the resolvents of the corresponding Dirac operators on the torus.

2.1 The resolvent of the Dirac operator

Let Λ be a lattice in $\mathbb{R}^2$ and V and W two periodic functions from $\mathbb{R}^2/\Lambda$ into $\mathbb{C}$. In this section we investigate the resolvent of the Dirac operator

$$D = \begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}, \text{ with } \partial = \frac{1}{2} \left(\frac{\partial}{\partial x_1} - \sqrt{-1} \frac{\partial}{\partial x_2} \right), \bar{\partial} = \frac{1}{2} \left(\frac{\partial}{\partial x_1} + \sqrt{-1} \frac{\partial}{\partial x_2} \right).$$

By abuse of notation the operator of multiplication with some function is denoted by the same symbol as the function. The operators ∂ and $\bar{\partial}$ coincide with holomorphic and anti-holomorphic derivation with respect to $z = x_1 + \sqrt{-1}x_2$. Therefore, we use sometimes the coordinates z and $\bar{z}$ instead of x . Consequently $\Lambda_{\mathbb{C}}$ denotes the corresponding sublattice of $\mathbb{C}$:

$$\Lambda_{\mathbb{C}} = \{ \gamma_1 + \sqrt{-1}\gamma_2 \in \mathbb{C} \mid \gamma \in \Lambda \}.$$

We use the natural scalar product on $\mathbb{C}^2$:

$$g(x, x') = x_1 x'_1 + x_2 x'_2 \quad \forall x, x' \in \mathbb{C}^2.$$

For all $k \in \mathbb{C}^2$ the map $\gamma \mapsto \exp(2\pi\sqrt{-1}g(\gamma, k))$ is a one-dimensional representation of Λ . Two of these representations are equivalent if and only if the difference of the corresponding elements in $\mathbb{C}^2$ is some element of the dual lattice

$$\Lambda^* := \{ \kappa \in \mathbb{R}^2 \mid g(\gamma, \kappa) \in \mathbb{Z} \forall \gamma \in \Lambda \}.$$

Hence the set of one-dimensional representations of Λ may be identified with $\mathbb{C}^2/\Lambda^*$. All these representations $[k] \in \mathbb{C}^2/\Lambda^*$ induce some line bundle on the torus $\mathbb{R}^2/\Lambda$, whose cross sections are functions ψ on $\mathbb{R}^2$ with the property

$$\psi(x + \gamma) = \exp(2\pi\sqrt{-1}g(\gamma, k)) \psi(x) \quad \text{for all } x \in \mathbb{R}^2 \text{ and all } \gamma \in \Lambda.$$

In the sequel we will often consider operators, which act on the space of cross sections of these bundles. These cross sections may be described in two different ways. We may either use a Fundamental domain or a Trivialization.

Fundamental domain 2.1. *We chose some fundamental domain*

$$\Delta = \{ x \in \mathbb{R}^2 \mid x = x' + \hat{q}_{\Delta} \hat{\gamma}_{\Delta} + \check{q}_{\Delta} \check{\gamma}_{\Delta} \text{ with } (\hat{q}_{\Delta}, \check{q}_{\Delta}) \in [0, 1]^2 \},$$

of the action of Λ on $\mathbb{R}^2$. Here $\hat{\gamma}_{\Delta}$ and $\check{\gamma}_{\Delta}$ are generators of Λ and x' is some element of $\mathbb{R}^2$. In the sequel we will sometimes pick out finitely many special elements of $\mathbb{R}^2/\Lambda$, and investigate the behaviour of some functions nearby these elements. In this case we will always assume that x' is chosen in such a way that the corresponding elements of Δ are contained in the interior of Δ . The sections of the line bundles corresponding to some $[k] \in \mathbb{C}^2/\Lambda^*$ may be described by usual functions on Δ , which on the boundary of Δ obey the conditions introduced above. Consequently the Hilbert space of cross sections of these line bundles are all equal to $L^2(\Delta)$. In this case the dual generators of the dual lattice are denoted by $\hat{\kappa}_{\Delta}$ and $\check{\kappa}_{\Delta}$,

with $g(\hat{\gamma}_\Delta, \hat{\kappa}_\Delta) = 1 = g(\check{\gamma}_\Delta, \check{\kappa}_\Delta)$ and $g(\hat{\gamma}_\Delta, \check{\kappa}_\Delta) = 0 = g(\check{\gamma}_\Delta, \hat{\kappa}_\Delta)$. Therefore, we have natural coordinates of the space $\mathbb{R}^2$ denoted by $\hat{q}_\Delta = g(x, \hat{\kappa}_\Delta)$ and $\check{q}_\Delta = g(x, \check{\kappa}_\Delta)$, and the dual space denoted by $\hat{p}_\Delta = g(\hat{\gamma}_\Delta, k)$ and $\check{p}_\Delta = g(\check{\gamma}_\Delta, k)$. Moreover, we assume that the orientation of the generators of Λ is positive, which means that the real number $\hat{\gamma}_{\Delta,1}\check{\gamma}_{\Delta,2} - \hat{\gamma}_{\Delta,2}\check{\gamma}_{\Delta,1}$ is positive and therefore equal to the Euclidean volume $\text{vol}(\mathbb{R}^2/\Lambda)$ of the torus. Due to this assumption the modular group $SL(2, \mathbb{Z})$ acts freely and transitively on the set of all possible choices of such generators. If we want to emphasize the dependence of some objects on this choice, we will decorate this object with an index Δ , but in general we shall omit this index.

Trivialization 2.2. The functions ψ_k are global non-vanishing sections of the line bundles corresponding to $[k]$. Hence all cross sections may be described by usual functions times these trivializations. However, these Trivializations are different for different representatives k and $k + \kappa$ of the class $[k]$. Therefore the representatives of the cross sections of the line bundles corresponding to $[k] \in \mathbb{C}^2/\Lambda^*$ belong to a vector bundle over $\mathbb{C}^2/\Lambda^*$. If for example we consider the Hilbert spaces of L^2 -sections, then the cross sections belong to the induced Hilbert bundle on the quotient $\mathbb{C}^2/\Lambda^*$, of the unitary representation $\kappa \mapsto \psi_\kappa$ of Λ^* on $L^2(\mathbb{R}^2/\Lambda)$. The sections $\psi(k)$ of this induced bundle have to obey the relation $\psi(k + \kappa) = \psi_{-\kappa}\psi(k)$. If we use this approach the elements of this Hilbert bundle are called periodic parts of the eigenfunctions, and the corresponding operators, which act on this Hilbert bundle, will be decorated by some tilde.

In the sequel the dependence on k , which is easier to describe with the help of the *Fundamental domain*, will be more important than the dependence on Λ , which is easier to describe with the help of the *Trivialization*. Hence we will mostly use the *Fundamental domain*. Obviously the operator of multiplication with ψ_k transforms the periodic parts of the eigenfunctions defined with the help of the *Trivialization* into the elements of the Hilbert space $L^2(\Delta)$ over the *Fundamental domain*. Therefore, the operators on the former Hilbert bundle are transformed into the operators on the latter Hilbert space by taking the conjugate with the operator ψ_k .

In addition to the hermitian scalar product the Hilbert spaces $L^2(\Delta)$ and $L^2(\mathbb{R}^2/\Lambda)$ have a natural complex conjugation denoted by $\psi \mapsto \bar{\psi}$ and a symmetric bilinear form denoted by

$$\langle \phi, \psi \rangle = \int_{\Delta} \phi(x)\psi(x)d^2x.$$

Consequently, the hermitian scalar product is denoted by $\langle \bar{\phi}, \psi \rangle$. Let $\partial_{[k]}$ and $\bar{\partial}_{[k]}$ be for all $k \in \mathbb{C}^2$ the unique closures of the unbounded operators ∂ and $\bar{\partial}$ on $L^2(\Delta)$ with the domain of smooth functions ψ , so that

$$\psi(x + \gamma) = \exp(2\pi\sqrt{-1}g(\gamma, k))\psi(x) \quad \text{for all } x \in \mathbb{R}^2 \text{ and all } \gamma \in \Lambda.$$

Obviously, these operators depend only on the representative $[k] \in \mathbb{C}^2/\Lambda^*$. The eigenfunctions of these operators are given by ψ_k with eigenvalues

$$\partial_{[k]}\psi_k = \pi(k_2 + \sqrt{-1}k_1)\psi_k \quad \text{and} \quad \bar{\partial}_{[k]}\psi_k = \pi(-k_2 + \sqrt{-1}k_1)\psi_k,$$

where k runs over all elements of the preimage of $[k]$ under the natural map $\mathbb{C}^2 \rightarrow \mathbb{C}^2/\Lambda^*$. Let $D(V, W, k)$ denote the operator

$$D(V, W, k) = \begin{pmatrix} V & \partial_{[k]} \\ -\bar{\partial}_{[k]} & W \end{pmatrix}.$$

The exact definition as a closed operator on some Banach space is part of the following

Theorem 2.3. *The resolvent $R(V, W, k, \lambda) = (\lambda \mathbf{1} - D(V, W, k))^{-1}$ defines a meromorphic map from $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda) \times \mathbb{C}^2 \times \mathbb{C}$ into the set of compact operators on $L^2(\Delta) \times L^2(\Delta)$. Moreover, for all $(V_0, W_0, k_0, \lambda_0) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda) \times \mathbb{C}^2 \times \mathbb{C}$ there exists some open neighbourhood $\mathcal{U} \subset L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda) \times \mathbb{C}^2$ of (V_0, W_0, k_0) and some holomorphic function $\check{P}$ from $\mathcal{U}$ into the finite rank projections on $L^2(\Delta) \times L^2(\Delta)$, such that the following equation holds:*

$$(\mathbf{1} - \check{P}(V, W, k)) R(V, W, k, \lambda) = R(V, W, k, \lambda) (\mathbf{1} - \check{P}(V, W, k))$$

for all $(V, W, k) \in \mathcal{U}$ and all λ in some small neighbourhood of λ_0 . Both functions are holomorphic on this domain. Finally, for all $2 \leq q < \infty$ the images of the values of the map $\check{P}$ are contained in $L^q(\Delta) \times L^q(\Delta)$.

Proof. We will prove the equivalent statements for the resolvent $\tilde{R}(V, W, k, \lambda)$ of the operator $\tilde{D}(V, W, k) = \begin{pmatrix} \psi_{-k} & 0 \\ 0 & \psi_{-k} \end{pmatrix} D(V, W, k) \begin{pmatrix} \psi_k & 0 \\ 0 & \psi_k \end{pmatrix}$, which is an operator on $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$. First we prove two lemmata.

Lemma 2.4. *Let $1 < p < 2$ and $q = \frac{2p}{2-p}$. Then there exists some constant S_p depending only on p with the following property: For any fixed $S'_p > S_p$ and for all $k \in \mathbb{C}^2$ there exists some $\delta > 0$ such that for all real $\lambda > \delta$ the operator $\tilde{R}(0, 0, k, \sqrt{-1}\lambda)$ from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ into $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$ has norm smaller than S'_p .*

Remark 2.5. *The constant $\sqrt{2}S_p$ is the Sobolev constant of the embedding $W^{1,p}(\mathbb{R}^2/\Lambda) \hookrightarrow L^{\frac{2p}{2-p}}(\mathbb{R}^2/\Lambda)$. With the optimal Sobolev constants in [Str, Chapter I. Section 4.6] we may set $S_p = \frac{p}{\sqrt{\pi(2-p)}}$.*

Proof. Let $\mathcal{G}_\lambda(x) = \frac{1}{2\pi} \mathcal{K}_0(\lambda \sqrt{g(x, x)})$ be the Green's function of $(\lambda^2 - \Delta)^{-1}$ on $\mathbb{R}^2$ ([St, Chapter V §3.1 and §6.5] and [G-J, Section 7.2]), where $\mathcal{K}_0$ denotes the modified Bessel function of zero-order. Then the resolvent $\left(\frac{\sqrt{-1}\lambda}{\bar{\partial}} \frac{-\partial}{\sqrt{-1}\lambda} \right)^{-1}$ has on $\mathbb{R}^2$ the integral kernel

$$\begin{pmatrix} a_\lambda(x - x') & b_\lambda(x - x') \\ c_\lambda(x - x') & d_\lambda(x - x') \end{pmatrix} d^2 x' = -4 \begin{pmatrix} \sqrt{-1}\lambda \mathcal{G}_{2\lambda}(x - x') & \partial \mathcal{G}_{2\lambda}(x - x') \\ -\bar{\partial} \mathcal{G}_{2\lambda}(x - x') & \sqrt{-1}\lambda \mathcal{G}_{2\lambda}(x - x') \end{pmatrix} d^2 x'.$$

The arguments given in the proof of [G-J, Proposition 7.2.1] also show that

- (i) $|\mathcal{K}_0(|t|)| \leq \mathbf{O}(1) \exp(-|t|)/|t|$ and $|\mathcal{K}'_0(|t|)| \leq \mathbf{O}(1) \exp(-|t|)/|t|$ for t bounded away from zero.

(ii) $\frac{1}{2\pi}\mathcal{K}_0(|t|) \sim -\ln(|t|)$ in some neighbourhood of $t = 0$.

(iii) $\frac{1}{2\pi}\mathcal{K}'_0(|t|) \sim -1/|t|$ in some neighbourhood of $t = 0$.

Hence the infinite sum

$$\begin{pmatrix} a_{\lambda,\Lambda}(x) & b_{\lambda,\Lambda}(x) \\ c_{\lambda,\Lambda}(x) & d_{\lambda,\Lambda}(x) \end{pmatrix} = \sum_{\gamma \in \Lambda} \begin{pmatrix} a_{\lambda}(x + \gamma) & b_{\lambda}(x + \gamma) \\ c_{\lambda}(x + \gamma) & d_{\lambda}(x + \gamma) \end{pmatrix}$$

converges on $\mathbb{R}^2$ to the integral kernel of $\tilde{\mathbf{R}}(0, 0, 0, \sqrt{-1}\lambda)$. Moreover, due to the Hardy–Littlewood–Sobolev theorem [St, Chapter V. §1.2 Theorem 1] the resolvent $\begin{pmatrix} \sqrt{-1}\lambda & \partial \\ -\partial & \sqrt{-1}\lambda \end{pmatrix}^{-1}$ of the Dirac operator on $\mathbb{R}^2$ is a bounded operator from $L^p(\mathbb{R}^2) \times L^p(\mathbb{R}^2)$ into $L^q(\mathbb{R}^2) \times L^q(\mathbb{R}^2)$. Using the equalities

$$\lambda^{\frac{2}{p}} \left(\int_{\mathbb{R}^2} |\psi(\lambda x)|^p d^2x \right)^{1/p} = \left(\int_{\mathbb{R}^2} |\psi(x)|^p d^2x \right)^{1/p} \quad \text{and}$$

$$\begin{pmatrix} a_{\lambda}(x - x') & b_{\lambda}(x - x') \\ c_{\lambda}(x - x') & d_{\lambda}(x - x') \end{pmatrix} d^2x' = \begin{pmatrix} a_1(\lambda x - \lambda x') & b_1(\lambda x - \lambda x') \\ c_1(\lambda x - \lambda x') & d_1(\lambda x - \lambda x') \end{pmatrix} \frac{d^2\lambda x'}{\lambda}$$

we conclude that the norm of this operator does not depend on λ . Moreover, this resolvent considered as an operator on $L^p(\mathbb{R}^2) \times L^p(\mathbb{R}^2)$ or $L^q(\mathbb{R}^2) \times L^q(\mathbb{R}^2)$ converges to zero, if λ tends to infinity. Due to the definition of the integral kernel of $\tilde{\mathbf{R}}(0, 0, 0, \sqrt{-1}\lambda)$ this implies the equality

$$\begin{pmatrix} a_{\lambda,\Lambda}(x) & b_{\lambda,\Lambda}(x) \\ c_{\lambda,\Lambda}(x) & d_{\lambda,\Lambda}(x) \end{pmatrix} = \lambda \begin{pmatrix} a_{1,\lambda\Lambda}(\lambda x) & b_{1,\lambda\Lambda}(\lambda x) \\ c_{1,\lambda\Lambda}(\lambda x) & d_{1,\lambda\Lambda}(\lambda x) \end{pmatrix},$$

where $\lambda\Lambda$ denotes the lattice $\lambda\Lambda = \{\lambda\gamma \mid \gamma \in \Lambda\}$. Moreover, for all S'_p larger than the Sobolev constant S_p there exists some $\delta > 0$ such that for all real $\lambda > \delta$, the resolvent $\tilde{\mathbf{R}}(0, 0, 0, \sqrt{-1}\lambda)$ is an operator from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ into $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$, whose norm is smaller than S'_p . This shows the claim for $k = 0$. Finally, this resolvent $\tilde{\mathbf{R}}(0, 0, 0, \sqrt{-1}\lambda)$ considered as an operator on $L^p(\mathbb{R}^2) \times L^p(\mathbb{R}^2)$ or $L^q(\mathbb{R}^2) \times L^q(\mathbb{R}^2)$ converges to zero, if λ tends to infinity. Obviously the difference $\tilde{\mathbf{D}}(0, 0, k) - \tilde{\mathbf{D}}(0, 0, 0)$ is the bounded operator $\begin{pmatrix} 0 & \pi(k_2 + \sqrt{-1}k_1) \\ \pi(k_2 - \sqrt{-1}k_1) & 0 \end{pmatrix}$. Hence the Neumann series

$$\begin{aligned} \tilde{\mathbf{R}}(0, 0, k, \sqrt{-1}\lambda) - \tilde{\mathbf{R}}(0, 0, 0, \sqrt{-1}\lambda) &= \\ &= \sum_{l=1}^{\infty} \tilde{\mathbf{R}}(0, 0, 0, \sqrt{-1}\lambda) \left(\begin{pmatrix} 0 & \pi(k_2 + \sqrt{-1}k_1) \\ \pi(k_2 - \sqrt{-1}k_1) & 0 \end{pmatrix} \tilde{\mathbf{R}}(0, 0, 0, \sqrt{-1}\lambda) \right)^l \end{aligned}$$

of operators from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ into $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$ converges to zero for $\lambda \rightarrow \infty$. **q.e.d.**

In the following lemma we make use of a family of norms on $L^2(\mathbb{R}^2/\Lambda)$. The family is parameterized by a positive real number ε . We consider the balls $B(x, \varepsilon)$ in $\mathbb{R}^2/\Lambda$, which for large ε becomes equal to the whole torus independent of x . But we are mainly interested in small ε . More precisely, if 2ε is smaller than the lengths of all non-zero periods, the corresponding balls have volume $\pi\varepsilon^2$. For all $U \in L^2(\mathbb{R}^2/\Lambda)$, the function

$$x \mapsto \left\| U|_{B(x, \varepsilon)} \right\|_2$$

is a continuous function on $\mathbb{R}^2/\Lambda$. Obviously, the supremum of this function in dependence of U , denoted by $\|U\|_{\varepsilon, 2}$, defines a norm on $L^2(\mathbb{R}^2/\Lambda)$, which is equivalent to the usual L^2 -norm. In fact, Fubini's theorem [R-S-I, Theorem I.22] and the invariance of the Lebesgue measure under translations [Ro-2, Chapter 14 Section 6 24. Proposition] yields the identity

$$\begin{aligned} \int_{\mathbb{R}^2/\Lambda} \int_{B(x, \varepsilon)} f(x') d^2x' d^2x &= \int_{\mathbb{R}^2/\Lambda} \int_{B(0, \varepsilon)} f(x+x') d^2x' d^2x = \\ \int_{B(0, \varepsilon)} \int_{\mathbb{R}^2/\Lambda} f(x+x') d^2x d^2x' &= \int_{B(0, \varepsilon)} \int_{\mathbb{R}^2/\Lambda} f(x) d^2x d^2x' = \\ &= \text{vol}(B(0, \varepsilon)) \int_{\mathbb{R}^2/\Lambda} f(x) d^2x \quad \forall f \in L^1(\mathbb{R}^2/\Lambda). \end{aligned}$$

For $f(x)$ equal to the square of the former function we obtain that the usual L^2 -norm of the former function is equal to $\sqrt{\text{vol}(B(x', \varepsilon))}$ times $\|U\|_2$. Hence we have

$$\sqrt{\frac{\text{vol}(B(x', \varepsilon))}{\text{vol}(\mathbb{R}^2/\Lambda)}} \|U\|_2 \leq \|U\|_{\varepsilon, 2} \leq \|U\|_2.$$

Due to [Yo, Appendix to Chapter V. Theorem 1.] the convex (weakly closed) subsets

$$\{U \in L^2(\mathbb{R}^2/\Lambda) \mid \|U\|_{\varepsilon, 2} \leq C\}$$

of $L^2(\mathbb{R}^2/\Lambda)$ are compact with respect to the weak topology for all $\varepsilon > 0$ and $C > 0$. Moreover, for any $U \in L^2(\mathbb{R}^2/\Lambda)$ the limit $\lim_{\varepsilon \downarrow 0} \|U\|_{\varepsilon, 2}$ is zero, but not uniformly on bounded sets of $L^2(\mathbb{R}^2/\Lambda)$. Hence for all U and all $C > 0$ there exists some small $\varepsilon > 0$, such that $\|U\|_{\varepsilon, 2} \leq C$.

Lemma 2.6. *All $C_p < S_p^{-1}$ (compare with Lemma 2.4 and Remark 2.5) have the following property: For all $\varepsilon > 0$ and open bounded subsets $\mathcal{O}$ of $\mathbb{C}^2$ there exists some $\delta > 0$ such that for all $\lambda > \delta$ the mapping $(V, W, k) \mapsto R(V, W, k, \sqrt{-1}\lambda)$ is a holomorphic weakly continuous map from $\{(V, W) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda) \mid \|V\|_{\varepsilon, 2} \leq C_p \text{ and } \|W\|_{\varepsilon, 2} \leq C_p\} \times \mathcal{O}$ into the compact operators on $L^2(\Delta) \times L^2(\Delta)$.*

Proof. We decompose the free resolvent $R(0, 0, k, \lambda) = R_{\varepsilon' \text{-near}}(0, 0, k, \lambda) + R_{\varepsilon' \text{-distant}}(0, 0, k, \lambda)$ into a sum of two operators, where the integral kernel of the first summand is the product of the integral kernel of $R(0, 0, k, \lambda)$, with the function depending on $x - x'$, which is equal to zero, if $\|x - x'\| > \varepsilon'$, and otherwise equal to 1. Consequently the integral kernel of the second summand is the product of the integral kernel of $R(0, 0, k, \lambda)$ with the function depending on $x - x'$, which is equal to zero, if $\|x - x'\| \leq \varepsilon'$, and otherwise equal to 1. Due to Hölder's inequality [R-S-I, Theorem III.1 (c)] the operators of multiplication with V and W are bounded operators from $L^q(\mathbb{R}^2/\Lambda)$ into $L^p(\mathbb{R}^2/\Lambda)$. Now Lemma 2.4 implies that all fixed $C_p < S_p^{-1}$ and all $(V, W, k) \in \{(V, W) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda) \mid \|V\|_{\varepsilon, 2} < C_p \text{ and } \|W\|_{\varepsilon, 2} < C_p\} \times \mathcal{O}$ and all large real λ the norm of the operator $\begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} R_{\varepsilon' \text{-near}}(0, 0, k, \sqrt{-1}\lambda)$ on $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ is for small $0 < \varepsilon' < \varepsilon$ bounded by some constant smaller than $\left(\frac{\text{vol}(B(0, \varepsilon))}{\text{vol}(B(0, \varepsilon - \varepsilon'))}\right)^{1/p} = \left(1 - \frac{\varepsilon'}{\varepsilon}\right)^{-2/p}$. In fact, for all $x \in \mathbb{R}^2/\Lambda$ the $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ -norm of the restriction of $\begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} R_{\varepsilon' \text{-near}}(0, 0, k, \sqrt{-1}\lambda)\psi$ to $B(x, \varepsilon - \varepsilon')$ is smaller than the norm of the restriction of ψ to $B(x, \varepsilon)$. Therefore, the bound on the operator $\begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} R_{\varepsilon' \text{-near}}(0, 0, k, \sqrt{-1}\lambda)$ follows from the identity

$$\int_{\mathbb{R}^2/\Lambda} \left\| f|_{B(x, \varepsilon)} \right\|_p^p dx = \text{vol}(B(0, \varepsilon)) \|f\|_p^p \quad \forall f \in L^p(\mathbb{R}^2/\Lambda),$$

which follows again from Fubini's theorem [R-S-I, Theorem I.22] and the invariance of the Lebesgue measure under translations [Ro-2, Chapter 14 Section 6 24. Proposition]. With very small ε' the norm of this operator $\begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} R_{\varepsilon' \text{-near}}(0, 0, k, \sqrt{-1}\lambda)$ on $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ is smaller than one. On the other hand we have seen in the proof of Lemma 2.4 that under the natural isometries of $L^p(\mathbb{R}^2/\Lambda)$ and $L^q(\mathbb{R}^2/\Lambda)$ onto $L^p(\mathbb{R}^2/\lambda\Lambda)$ and $L^q(\mathbb{R}^2/\lambda\Lambda)$, respectively, the resolvent $R(0, 0, k, \sqrt{-1})$ considered as an operator from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ to $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$ is transformed into the resolvent $R(0, 0, k, \sqrt{-1}\lambda)$ considered as an operator from $L^p(\mathbb{R}^2/\lambda\Lambda) \times L^p(\mathbb{R}^2/\lambda\Lambda)$ to $L^q(\mathbb{R}^2/\lambda\Lambda) \times L^q(\mathbb{R}^2/\lambda\Lambda)$. Moreover, the asymptotic behaviour (i) of the Green's function of the Laplace operator mentioned in the proof of Lemma 2.4 implies that for large λ , the integral kernel of this resolvent becomes arbitrary small, if the distance of $x - x'$ in $\mathbb{R}^2/\lambda\Lambda$ becomes large. Hence for all $(V, W, k) \in \{(V, W) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda) \mid \|V\|_{\varepsilon, 2} < C_p \text{ and } \|W\|_{\varepsilon, 2} < C_p\} \times \mathcal{O}$ the norm of the operator $\begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} R_{\varepsilon' \text{-distant}}(0, 0, k, \sqrt{-1}\lambda)$ on $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ converges in the limit $\lambda \rightarrow \infty$ uniformly to zero. Consequently the Neumann series

$$\tilde{R}(V, W, k, \sqrt{-1}\lambda) = \sum_{l=0}^{\infty} \tilde{R}(0, 0, k, \sqrt{-1}\lambda) \left(\begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \tilde{R}(0, 0, k, \sqrt{-1}\lambda) \right)^l$$

converges to a holomorphic function with values in the bounded operators from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ to $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$. The latter Banach space is naturally contained in the former Banach space. Therefore, it remains to show, firstly that the values of this function are compact operators on the former Banach space, and secondly that these functions are weakly continuous.

In doing so we consider again the free resolvent $\tilde{R}(0, 0, k, \sqrt{-1}\lambda)$ as an operator from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ to $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$, but this time we assume the inequality $q < \frac{2p}{2-p}$, and therefore $1/p = 1/q + 1/r$ with some $r > 2$. Due to the Hausdorff–Young inequality [R-S-II, Theorem IX.8] the (not normalized) Fourier transform

$$F : \psi \mapsto \left(\hat{\psi}(\kappa) \right)_{\kappa \in \Lambda^*}, \text{ with } \psi(x) = \sum_{\kappa \in \Lambda^*} \hat{\psi}(\kappa) \exp(2\pi\sqrt{-1}g(x, \kappa)).$$

is a bounded map from $L^p(\mathbb{R}^2/\Lambda)$ into $\ell_{\frac{p}{p-1}}(\Lambda^*)$ and the inverse F^{-1} from $\ell_{\frac{q}{q-1}}(\Lambda^*)$ into $L^q(\mathbb{R}^2/\Lambda)$. If λ does not belong to the spectrum of the free Dirac operator $\tilde{D}(0, 0, k)$, then the Fourier transform $F\tilde{R}(0, 0, k, \sqrt{-1}\lambda)F^{-1}$ acts on $\ell_{\frac{p}{p-1}}(\Lambda^*) \times \ell_{\frac{p}{p-1}}(\Lambda^*)$ as the point-wise multiplication with the sequence of matrices

$$\begin{pmatrix} \sqrt{-1}\lambda & -\pi(k_2 + \kappa_2 + \sqrt{-1}(k_1 + \kappa_1)) \\ -\pi(k_2 + \kappa_2 - \sqrt{-1}(k_1 + \kappa_1)) & \sqrt{-1}\lambda \end{pmatrix}^{-1}.$$

Obviously, for all $k \in \mathbb{C}^2$, $\varepsilon > 0$ and $r > 2$ there exists some $\delta > 0$ such that all real $\lambda > \delta$ fulfill the following estimate:

$$\left(\sum_{\kappa \in \Lambda^*} \left\| \begin{pmatrix} \sqrt{-1}\lambda & -\pi(k_2 + \kappa_2 + \sqrt{-1}(k_1 + \kappa_1)) \\ -\pi(k_2 + \kappa_2 - \sqrt{-1}(k_1 + \kappa_1)) & \sqrt{-1}\lambda \end{pmatrix}^{-1} \right\|^r \right)^{1/r} < \infty.$$

With the help of Hölder's inequality [R-S-I, Theorem III.1 (c)] we conclude that for all $k \in \mathbb{C}^2$ there exists some $\delta > 0$ such that for all real $\lambda > \delta$ the resolvent $\tilde{R}(0, 0, k, \sqrt{-1}\lambda)$ is a bounded operator from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ into $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$. Let $\tilde{P}_n$ be the natural projection onto the span of all ψ_κ with $g(\kappa, \kappa) \leq n$. The same argument shows that for all $l \in \mathbb{N}$ and the same V and W as before the sequence

$$\tilde{R}(0, 0, k, \sqrt{-1}\lambda) \left(\tilde{P}_n \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \tilde{P}_n \tilde{R}(0, 0, k, \sqrt{-1}\lambda) \right)^l$$

converges in the limit $n \rightarrow \infty$ uniformly with respect to V and W to the operator

$$\tilde{R}(0, 0, k, \sqrt{-1}\lambda) \left(\begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \tilde{R}(0, 0, k, \sqrt{-1}\lambda) \right)^l$$

on $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$. Since all elements of the sequences are finite rank operators, the limits are compact operators [R-S-I, Theorem VI.12]. Obviously these sequences are sequences of weakly continuous functions in dependence of (V, W) . Since the uniform limits of continuous functions on a compact Hausdorff space are again a continuous functions, the limits are weakly continuous on the sets

$$\{(V, W) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda) \mid \|V\|_{\varepsilon, 2} \leq C_p \text{ and } \|W\|_{\varepsilon, 2} \leq C_p\} \times \mathcal{O}.$$

Due to the considerations above the Neumann series converges uniformly on these sets, and the lemma follows. **q.e.d.**

Continuation of the proof of Theorem 2.3. In the last lemma we actually proved that for all $(V_0, W_0, k_0) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda) \times \mathbb{C}^2$, there exists a complex number λ_1 and some open neighbourhood $\mathcal{U}$ of (V_0, W_0, k_0) , so that the map $(V, W, k) \mapsto \tilde{R}(V, W, k, \lambda_1)$ is a holomorphic map from $\mathcal{U}$ into the compact operators on $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$. Thus, due to the Riesz–Schauder theorem [R-S-I, Theorem VI.15] the spectrum of the operator $\tilde{R}(V, W, k, \lambda_1)$ is discrete and the only limit point of the spectrum is zero. Let $\mathcal{S} \subset \mathbb{C}$ denote the set

$$\mathcal{S} = \left\{ \lambda_1 - \frac{1}{\lambda} \mid \lambda \text{ is an eigenvalue of } \tilde{R}(V_0, W_0, k_0, \lambda_1) \right\}.$$

Then for all $\lambda \in \mathbb{C} \setminus \mathcal{S}$, the operator

$$\tilde{R}(V_0, W_0, k_0, \lambda) = \frac{\tilde{R}(V_0, W_0, k_0, \lambda_1)}{\lambda_1 - \lambda} \left(\frac{1}{\lambda_1 - \lambda} - \tilde{R}(V_0, W_0, k_0, \lambda_1) \right)^{-1}$$

is compact and depends holomorphically on λ .

Let $\lambda_0 \in \mathcal{S}$ be a pole of this map and choose a positive number $\varepsilon > 0$, such that there exists no other pole in the disc $\{\lambda \mid |\lambda - \lambda_0| < 2\varepsilon\}$. Then there exists an open neighbourhood $\mathcal{U} \subset L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda) \times \mathbb{C}^2$ of (V_0, W_0, k_0) such that $\tilde{R}(V, W, k, \lambda)$ is compact for all $(V, W, k) \in \mathcal{U}$ and all λ on the circle $\{\lambda \mid |\lambda - \lambda_0| = \varepsilon\}$. Now we define a map $\check{P}$ from $\mathcal{U}$ into the set of compact operators on $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ by

$$\check{P}(V, W, k) = -\frac{1}{2\pi\sqrt{-1}} \oint_{\{\lambda \mid |\lambda - \lambda_0| = \varepsilon\}} \tilde{R}(V, W, k, \lambda) d\lambda.$$

The combination of the results from [R-S-I, VI.5] and [R-S-IV, Appendix to XII.1] implies that for all $(V, W, k) \in \mathcal{U}$, $\check{P}(V, W, k)$ is a finite-dimensional projector, which commutes with $\tilde{D}$. Also the eigenvalues of the restriction to the corresponding finite-dimensional subspace of $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ are elements of the disc $\{\lambda \mid |\lambda - \lambda_0| < \varepsilon\}$. Furthermore, all eigenfunctions, which correspond to eigenvalues inside of this disc, are elements of the finite-dimensional subspace mentioned above. It remains to prove the last statement of the theorem. In Lemma 2.6 we actually proved that for all $1 < p < 2$ and all $p \leq q < \frac{2p}{2-p}$ the resolvent $\tilde{R}(V, W, k, \sqrt{-1}\lambda)$ is a compact operator from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ into $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$. This implies that $\check{P}(V, W, k)$ is a finite-dimensional projector on $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$, for all $2 \leq q < \infty$. **q.e.d.**

Remark 2.7. Due to [St, Chapter V §.3.4 Lemma 3.] the operators $\partial(1 - \partial\bar{\partial})^{-1/2}$ and $\bar{\partial}(1 - \partial\bar{\partial})^{-1/2}$ are bounded operators on $L^p(\mathbb{R}^2/\Lambda)$ with $1 < p < 2$. Therefore any function $f \in L^p(\mathbb{R}^2/\Lambda)$ with either $\partial f \in L^p(\mathbb{R}^2/\Lambda)$ or $\bar{\partial}f \in L^p(\mathbb{R}^2/\Lambda)$ belongs to the Sobolev space $W^{1,p}(\mathbb{R}^2/\Lambda)$ [St, Chapter V §.3.4 Theorem 3.]. Hence our arguments show that the eigenfunctions belong to the intersection $\bigcap_{1 < p < 2} W^{1,p}(\Delta) \times W^{1,p}(\Delta)$ of Sobolev spaces.

Remark 2.8. Unfortunately the sets described in Lemma 2.6, by which the whole complex Bloch varieties depend weakly continuous on the potentials, do not contain balls, whose radius is larger or equal to S_p^{-1} (compare with Remark 2.5). This failure will cause the most difficulties of this approach. So let us elaborate a bit on this failure. For all elements (V_0, W_0) of the sets

$$\{(V, W) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda) \mid \|V\|_{\varepsilon,2} \leq C_p \text{ and } \|W\|_{\varepsilon,2} \leq C_p\} \text{ with } C_p < S_p^{-1}$$

the balls $B(V_0, C'_p) \times B(W_0, C'_p)$, with $C'_p < S_p^{-1}$ are contained in another set of this form

$$\{(V, W) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda) \mid \|V\|_{\varepsilon',2} \leq C''_p \text{ and } \|W\|_{\varepsilon',2} \leq C''_p\} \text{ with } C''_p < S_p^{-1}$$

and with some small $\varepsilon' > 0$. Moreover, all bounded subsets of $L^r(\mathbb{R}^2/\Lambda) \times L^r(\mathbb{R}^2/\Lambda)$ with $2 < r$ are contained in a set of this form. Finally, we shall see in Lemma 3.19 that a weakly convergent subsequence is contained in a set of this form, if the weak limit of the measures $V(x)\bar{V}(x)d^2x$ and $W(x)\bar{W}(x)d^2x$ exist and contain no point measures with mass larger or equal to S_p^{-2} . If the norms of these sequences are bounded, this can fail only at finitely many points of $\mathbb{R}^2/\Lambda$. In this case the limits of the corresponding resolvents are the resolvents of some perturbation of the Dirac operator acting on eigenfunctions with poles (or on line bundles of non-vanishing degree) as described in Section 3.4.

We close this section with a proof of the *strong unique continuation property* of the eigenfunctions of the Dirac operators. We use the classical Carleman method, which is based on an improved Sobolev inequality (compare with [Ca] and [Wo, Proposition 1.3]).

Carleman inequality 2.9. *There exists some constant S_p (compare with Lemma 2.4 and Remark 2.5) depending only on $1 < p < 2$, such that for all $n \in \mathbb{Z}$ and all $\psi \in C_0^\infty(\mathbb{C} \setminus \{0\}) \times C_0^\infty(\mathbb{C} \setminus \{0\})$ the following inequality holds:*

$$\| |z|^{-n} \psi \|_{\frac{2p}{2-p}} \leq S_p \| |z|^{-n} \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} \psi \|_p.$$

The literature [Je, Ki-1, Ma, Ki-2] deals with the much more difficult higher-dimensional case and does not treat our case. David Jerison pointed out to the author, that the arguments of [Wo, Proposition 2.6], where the analogous but weaker statement about the gradient term of the Laplace operator is treated, carry over to the Dirac operator.

Proof. Dolbeault's Lemma [Gu-Ro, Chapter I Section D 2. Lemma] implies for all smooth ψ with compact support the equality

$$\psi(z) = \int_{\mathbb{C}} \begin{pmatrix} 0 & (z'-z)^{-1} \\ (\bar{z}-\bar{z}')^{-1} & 0 \end{pmatrix} \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} \psi(z') \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}}.$$

In fact, the components of the difference of the left hand side minus the right hand side are holomorphic and anti-holomorphic functions on $\mathbb{C}$, respectively, which vanish at $z = \infty$. In particular, the integrals $\int_{\mathbb{C}} z^n \bar{\partial} \psi_1 d\bar{z} \wedge dz$ and $\int_{\mathbb{C}} \bar{z}^n \partial \psi_2 d\bar{z} \wedge dz$ with $n \in \mathbb{N}_0$ are proportional

to the Taylor coefficients of ψ at ∞ , which vanish. Moreover, if the support of ψ does not contain 0 and therefore also a small neighbourhood of 0, then the integrals $\int_{\mathbb{C}} z^{-n} \bar{\partial} \psi_1 d\bar{z} \wedge dz$ and $\int_{\mathbb{C}} \bar{z}^{-n} \partial \psi_2 d\bar{z} \wedge dz$ with $n \in \mathbb{N}$ are proportional to the Taylor coefficients of ψ at 0, which in this case also vanish. These cancellations follow also from partial integration. We conclude that for all $n \in \mathbb{Z}$

$$\psi(z) = \int_{\mathbb{C}} \begin{pmatrix} 0 & \left(\frac{z}{z'}\right)^n \frac{1}{z'-z} \\ \left(\frac{\bar{z}}{\bar{z}'}\right)^n \frac{1}{\bar{z}-\bar{z}'} & 0 \end{pmatrix} \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} \psi(z') \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}}.$$

In fact, for negative n the left hand side minus the left hand side of the foregoing formula is equal to the Taylor polynomial of ψ at ∞ up to order $|n|$, and for positive n equal to the Taylor polynomial of ψ at 0 up to order $n-1$. Finally, the Hardy–Littlewood–Sobolev theorem [St, Chapter V. §1.2 Theorem 1] implies that the operator with integral kernel

$$\begin{pmatrix} 0 & \left(\frac{|z'|z}{|z|\bar{z}'}\right)^n \frac{1}{z'-z} \\ \left(\frac{|z|\bar{z}}{|z'|\bar{z}'}\right)^n \frac{1}{\bar{z}-\bar{z}'} & 0 \end{pmatrix} \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}}$$

from $L^p(\mathbb{C}) \times L^p(\mathbb{C})$ into $L^{\frac{2p}{2-p}}(\mathbb{C}) \times L^{\frac{2p}{2-p}}(\mathbb{C})$ is bounded by some constant S_p not depending on n , and maps $|z|^{-n} \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} \psi$ onto $|z|^{-n} \psi$. **q.e.d.**

Due to a standard argument (e. g. [So, Proof of Theorem 5.1.4] and [Wo, Section Carleman Method]) this *Carleman inequality* implies the

Strong unique continuation property 2.10. *Let V and W be potentials in $L^2_{\text{loc}}(\mathcal{O})$ on an open connected set $0 \ni \mathcal{O} \subset \mathbb{C}$ and $\psi \in W^{1,p}_{\text{loc}}(\mathcal{O}) \times W^{1,p}_{\text{loc}}(\mathcal{O})$ an element of the kernel of $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ on $\mathcal{O}$ with $1 < p < 2$. If the $L^{\frac{2p}{2-p}}$ -norm of the restriction of ψ to the balls $B(0, \varepsilon)$ converges in the limit $\varepsilon \downarrow 0$ faster to zero than any power of ε :*

$$\left(\int_{B(0, \varepsilon)} (|\psi_1|^{\frac{2p}{2-p}} + |\psi_2|^{\frac{2p}{2-p}}) d^2x \right)^{\frac{2-p}{2p}} \leq \mathbf{O}(\varepsilon^n) \quad \forall n \in \mathbb{N},$$

then ψ vanishes identically on $\mathcal{O}$.

Proof. The question is local so we may assume that V and W are elements of L^2 rather than L^2_{loc} . We fix ε small enough that $\max \{ \|V\|_{L^2(B(z, 2\varepsilon))}, \|W\|_{L^2(B(z, 2\varepsilon))} \} \leq 1/(2S_p)$ for all z , where S_p is the constant of the *Carleman inequality*. Let $\phi \in C^\infty$ be 1 on $B(0, \varepsilon)$ and 0 on $\mathbb{C} \setminus B(0, 2\varepsilon)$. A limiting argument using the infinite order vanishing of ψ and the equality $\begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} \psi = -\begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \psi$ shows that the proof of the *Carleman inequality* is also true for $\phi\psi$. So

$$\| |z|^{-n} \phi \psi \|_{\frac{2p}{2-p}} \leq S_p \| |z|^{-n} \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} \phi \psi \|_p \leq S_p \| |z|^{-n} \phi \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} \psi \|_p + S_p \| |z|^{-n} E \|_p.$$

Here E (for error) $= \begin{pmatrix} \psi_2 \partial \phi \\ -\psi_1 \bar{\partial} \phi \end{pmatrix}$ is an $L^p \times L^p$ function supported in $\{x \mid \varepsilon \leq |z| \leq 2\varepsilon\}$. Using the equality $\begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} \psi = -\begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \psi$ and Hölder's inequality [R-S-I, Theorem III.1 (c)] yields

$$\begin{aligned} \| |z|^{-n} \phi \psi \|_{\frac{2p}{2-p}} &\leq S_p \| |z|^{-n} \phi \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \psi \|_p + S_p \| |z|^{-n} E \|_p \\ &\leq S_p \max \{ \|V\|_{L^2(B(0,2\varepsilon))}, \|W\|_{L^2(B(0,2\varepsilon))} \} \| |z|^{-n} \phi \psi \|_{\frac{2p}{2-p}} + S_p \| |z|^{-n} E \|_p. \end{aligned}$$

By the choice of ε the first term can be absorbed into a factor 2

$$\| |z|^{-n} \phi \psi \|_{\frac{2p}{2-p}} \leq 2S_p \| |z|^{-n} E \|_p.$$

Now comes the crucial observation: E is supported in $\{z \mid \varepsilon \leq |z| \leq 2\varepsilon\}$, so

$$\| |z|^{-n} \phi \psi \|_{\frac{2p}{2-p}} \leq 2S_p \varepsilon^{-n} \|E\|_p \quad \text{and} \quad \left\| \left(\frac{\varepsilon}{|z|} \right)^n \phi \psi \right\|_{\frac{2p}{2-p}} \leq 2S_p \|E\|_p.$$

Using the limit $n \rightarrow \infty$ we conclude that $\phi \psi$ vanishes on $B(0, \varepsilon)$, and therefore also ψ . In other words, the set $\{z \mid \psi \text{ vanishes to infinite order at } z\}$ is open, and in fact contains a ball of fixed radius ε centered at any of its points. So this set must be all of $\mathcal{O}$ and the proof is complete. **q.e.d.**

2.2 The Bloch variety and the Fermi curve

For two given periodic potentials V and W the corresponding Dirac operator commutes with all shifts by the lattice vectors. Hence these shifts and the Dirac operator may be diagonalized simultaneously. The common spectrum of all these operators is called a Bloch variety. The proof in [R-S-IV, Theorem XIII.97], which applies to Schrödinger operators with periodic potential, carries over to the Dirac operators and shows that a Dirac operator with periodic potentials V and W is the direct integral of the operators $\tilde{D}(V, W, k)$ where k runs through $\mathbb{R}^2/\Lambda^*$ (compare with [Ku]). The Bloch variety may be considered as the set of points $([k], \lambda) \in \mathbb{R}^2/\Lambda^* \times \mathbb{C}$, so that λ is an eigenvalue of $D(V, W, k)$. Let us call the collection of all points $(k, \lambda) \in \mathbb{C}^2 \times \mathbb{C}$, so that λ is an eigenvalue of $D(V, W, k)$, complex Bloch variety and denote this set by $\mathcal{B}(V, W)$. Due to Theorem 2.3 for all $(k_0, \lambda_0) \in \mathbb{C}^2 \times \mathbb{C}$ there exists some open neighbourhood $\mathcal{U} \times \{\lambda \mid |\lambda - \lambda_0| < \varepsilon\} \subset \mathbb{C}^2 \times \mathbb{C}$ of (k_0, λ_0) and a holomorphic $l \times l$ matrix valued function A on $\mathcal{U}$, such that $\mathcal{B}(V, W) \cap (\mathcal{U} \times \{\lambda \mid |\lambda - \lambda_0| < \varepsilon\})$ is the zero set of the function

$$\mathcal{U} \times \{\lambda \mid |\lambda - \lambda_0| < \varepsilon\} \rightarrow \mathbb{C}, \quad (k, \lambda) \mapsto \det(\lambda \mathbf{1} - A(k)).$$

Hence $\mathcal{B}(V, W)$ is a subvariety of $\mathbb{C}^2 \times \mathbb{C}$. The action of the dual lattice Λ^* on $\mathbb{C}^2$ given by $\kappa.k = \kappa + k$ for all $\kappa \in \Lambda^*$ and all $k \in \mathbb{C}^2$, lifts to an action of the dual lattice Λ^* on this subspace $\mathcal{B}(V, W)$ of $\mathbb{C}^2 \times \mathbb{C}$. The set of orbits of this action is equal to the subspace $\mathcal{B}(V, W)/\Lambda^*$ of all elements $([k], \lambda) \in \mathbb{C}^2/\Lambda^* \times \mathbb{C}$, such that λ is an eigenvalue of the operator $D(V, W, k)$.

We call the zero energy level of the complex Bloch variety the *complex Fermi curve* and denote it by

$$\mathcal{F}(V, W) = \{k \in \mathbb{C}^2 \mid (k, 0) \in \mathcal{B}(V, W)\}.$$

Sometimes we shall not distinguish between the values of k and the corresponding elements of $\mathcal{F}$. However, mostly k denotes the corresponding $\mathbb{C}^2$ -valued function on the *complex Fermi curves*. Due to Theorem 2.3 the *complex Fermi curve* $\mathcal{F}(V, W)$ is locally the zero set of one holomorphic function. Hence it is a pure one-dimensional subvariety of $\mathbb{C}^2$, which is invariant under the action of Λ^* . Strictly speaking, the *complex Fermi curve* is the quotient space $\mathcal{F}(V, W)/\Lambda^*$, but we shall also call $\mathcal{F}(V, W)$ a *complex Fermi curve*.

These *complex Fermi curves* are of special interest for two reasons: Firstly the corresponding eigenfunctions belong to the kernel of the Dirac operator and therefore give rise to local immersions of surfaces into the three-dimensional Euclidean space due to the local Weierstraß representation. Secondly, these *complex Fermi curves* are the spectral curves of the corresponding integrable system described by the Davey–Stewartson equation.

Due to a general feature of differential operators, in the high energy limit the higher-order derivatives dominate the lower order derivatives. Moreover, typically the complex Bloch variety converges in the high energy limit to the complex Bloch variety of the free Dirac operator and has an asymptotic expansion for smooth potentials, which can be calculated explicitly (compare with [Sch, Chapter 1]). In the remainder of this section we shall investigate the asymptotic behaviour of the *complex Fermi curves* for general potentials $V, W \in L^2(\mathbb{R}^2/\Lambda)$.

For this purpose we use a covariance property of the *complex Fermi curves* under some unitary transformations of the potentials. By definition of the operators $\partial_{[k]}$ and $\bar{\partial}_{[k]}$ they transform under translations by some k' as follows:

$$\begin{aligned} \partial_{[k+k']} &= \psi_{-k'} \partial_{[k]} \psi_{k'} & \text{if } k'_2 + \sqrt{-1}k'_1 &= 0, \\ \bar{\partial}_{[k+k']} &= \psi_{-k'} \bar{\partial}_{[k]} \psi_{k'} & \text{if } k'_2 - \sqrt{-1}k'_1 &= 0. \end{aligned}$$

For all κ there exist unique pairs $(k_\kappa^-, k_\kappa^+) \in \mathbb{C}^2 \times \mathbb{C}^2$, whose first element obeys the former condition and the second element the latter condition, and whose difference $k_\kappa^+ - k_\kappa^-$ is equal to κ .

We conclude that for all $\kappa \in \Lambda^*$ the Dirac operators transform as follows:

$$D(V, W, k + k_\kappa^\pm) = \begin{pmatrix} \psi_{-k_\kappa^-} & 0 \\ 0 & \psi_{-k_\kappa^+} \end{pmatrix} D(\psi_{-\kappa} V, \psi_\kappa W, k) \begin{pmatrix} \psi_{k_\kappa^+} & 0 \\ 0 & \psi_{k_\kappa^-} \end{pmatrix}.$$

This implies the following

Lemma 2.11. *For all $\kappa \in \Lambda^*$ the translation $k \mapsto k + k_\kappa^+ = k + k_\kappa^- + \kappa$ induces a biholomorphic isomorphism of the complex Fermi curves $\mathcal{F}(V, W) \simeq \mathcal{F}(\psi_{-\kappa} V, \psi_\kappa W)$. **q.e.d.***

For all potentials $V, W \in L^2(\mathbb{R}^2/\Lambda)$ there exists an $\varepsilon > 0$ such that $\|V\|_{\varepsilon, 2} < S_p^{-1}$ and $\|W\|_{\varepsilon, 2} < S_p^{-1}$. Consequently, for all $\kappa \in \Lambda^*$ also the potentials $\psi_{-\kappa} V$ and $\psi_\kappa W$ obey these estimates. Moreover, the sequence of potentials $(\psi_{-\kappa} V, \psi_\kappa W)$ converges in the limit $g(\kappa, \kappa) \rightarrow \infty$ weakly to the zero potentials in $L^2(\mathbb{R}^2/\Lambda)$. More precisely, if we identify

the Banach space $L^1(\mathbb{R}^2/\Lambda) \simeq L^1(\Delta)$ with the corresponding natural subspace of $L^1(\mathbb{R}^2)$, then due to the Riemann–Lebesgue Lemma [R-S-II, Theorem IX.7] the Fourier coefficients indexed by $\kappa \in \Lambda^*$ of any element in $L^1(\mathbb{R}^2/\Lambda)$ converges in the limit $g(\kappa, \kappa) \rightarrow \infty$ to zero. Therefore, for any $V, W \in L^2(\mathbb{R}^2/\Lambda)$ and any weakly open neighbourhood $\mathcal{U}$ of $0 \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ there exists some $\delta > 0$ such that for all $\kappa \in \Lambda^*$ with $g(\kappa, \kappa) > \delta^{-2}$ the pair of potentials $(\psi_{-\kappa}V, \psi_{\kappa}W)$ belongs to $\mathcal{U}$. We conclude from Lemma 2.6 that for all k in a bounded open subsets $\mathcal{O} \subset \mathbb{C}^2 \setminus \mathcal{F}(0, 0)$ the resolvents $R(\psi_{-\kappa}V, \psi_{\kappa}W, k, 0)$ converge in the limit $g(\kappa, \kappa) \rightarrow \infty$ uniformly to $R(0, 0, k, 0)$. Obviously the eigenvalues of the operators $\begin{pmatrix} 0 & \pm \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix} \hat{D}(V, W, k)$ are completely determined by the *complex Fermi curve* $\mathcal{F}(V, W)$. Conversely, one resolvent of the family of resolvents $R(V, W, k, 0)$ indexed by $k \in \mathbb{C}^2 \setminus \mathcal{F}(V, W)$ determines the spectrum of the operators $\begin{pmatrix} 0 & \pm \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix} \hat{D}(V, W, k)$ and therefore also all other resolvents of this family. Therefore, the intersections of the *complex Fermi curves* $\mathcal{F}(\psi_{-\kappa}V, \psi_{\kappa}W)$ with all bounded open subsets $\mathcal{O} \subset \mathbb{C}^2$ converge in the limit $g(\kappa, \kappa) \rightarrow \infty$ to $\mathcal{O} \cap \mathcal{F}(0, 0)$. Moreover, the corresponding eigenfunctions converge to the corresponding eigenfunctions of the free Dirac operator. An application of Lemma 2.11 yields that the intersections of $\mathcal{F}(V, W) \cap (\mathcal{O} + k_{\kappa}^+) = (\mathcal{F}(V, W) \cap (\mathcal{O} + k_{\kappa}^-)) + \kappa$ converges to $\mathcal{F}(0, 0) \cap (\mathcal{O} + k_{\kappa}^+) = (\mathcal{F}(0, 0) \cap (\mathcal{O} + k_{\kappa}^-)) + \kappa$. An easy calculation shows that the free *complex Fermi curve* is equal to

$$\mathcal{F}(0, 0) = \bigcup_{\kappa \in \Lambda^*} \kappa + \{k \in \mathbb{C}^2 \mid k_2 + \sqrt{-1}k_1 = 0\} \bigcup_{\kappa \in \Lambda^*} \kappa + \{k \in \mathbb{C}^2 \mid k_2 - \sqrt{-1}k_1 = 0\}.$$

Therefore, the free *complex Fermi curve* $\mathcal{F}(0, 0)/\Lambda^*$ is isomorphic to the two subvarieties $\{k \mid k_1 \pm \sqrt{-1}k_2 = 0\}$ of $\mathbb{C}^2$, glued at infinitely many ordinary double points indexed by $\kappa \in \Lambda^*$ of the form $(k_{\kappa}^-, k_{\kappa}^+)$ with

$$\begin{aligned} k_{\kappa}^- &= (-\kappa_1/2 - \sqrt{-1}\kappa_2/2, -\kappa_2/2 + \sqrt{-1}\kappa_1/2) \text{ and} \\ k_{\kappa}^+ &= (\kappa_1/2 - \sqrt{-1}\kappa_2/2, \kappa_2/2 + \sqrt{-1}\kappa_1/2) = k_{\kappa}^- + \kappa. \end{aligned}$$

Also on these components the eigenfunctions of the free Dirac operator are equal to

$$\psi = \psi_k \begin{pmatrix} 0 \\ 1 \end{pmatrix} \text{ on } \{k \mid k_1 - \sqrt{-1}k_2 = 0\} \text{ and } \psi = \psi_k \begin{pmatrix} 1 \\ 0 \end{pmatrix} \text{ on } \{k \mid k_1 + \sqrt{-1}k_2 = 0\}.$$

respectively. For positive ε and δ let $\mathcal{V}_{\varepsilon, \delta}^{\pm}$ denote the intersection of $\mathcal{F}(V, W)$ with the following open subsets $\mathcal{U}_{\varepsilon, \delta}^{\pm}$ of $\mathbb{C}^2$:

$$\mathcal{U}_{\varepsilon, \delta}^{\pm} = \{k \in \mathbb{C}^2 \mid |k_1 \pm \sqrt{-1}k_2| < \varepsilon, \|k\| > 1/\delta \text{ and } \|k - k_{\kappa}^{\pm}\| > \varepsilon \forall \kappa \in \Lambda^* \setminus \{0\}\}.$$

Here $\|k\|$ denotes the natural hermitian norm $\|k\| = \sqrt{g(k, k)}$ on $\mathbb{C}^2$. Since for small ε and δ these open sets $\mathcal{U}_{\varepsilon, \delta}^{\pm}$ intersect any Λ^* -orbit at most once, we may consider these open sets as subsets of $\mathbb{C}^2/\Lambda^*$. Since the quotient of $\mathbb{C}^2$ modulo the sublattice $\{k_{\kappa}^+ + \kappa' \mid \kappa, \kappa' \in \Lambda^*\} = \{k_{\kappa}^- + \kappa' \mid \kappa, \kappa' \in \Lambda^*\}$ is compact, the combination of Lemma 2.6 and Lemma 2.11 implies the following

Theorem 2.12. *For all pairs of potentials $V, W \in L^2(\mathbb{R}^2/\Lambda)$ and all $\varepsilon > 0$ there exists a $\delta > 0$ so that the two open sets $\mathcal{V}_{\varepsilon, \delta}^{\pm}$ are connected complex one-dimensional submanifolds of $\mathbb{C}^2$ with one-dimensional kernels of the corresponding Dirac operators $\mathcal{D}(V, W, k)$. Moreover, the relative complement of $\mathcal{V}_{\varepsilon, \delta}^+ \cup \mathcal{V}_{\varepsilon, \delta}^-$ in $\mathcal{F}(V, W)/\Lambda^*$ decomposes into a compact set contained in $\{k \in \mathbb{C}^2 \mid \|k\| \leq 1/\delta\}/\Lambda^*$ and infinitely many small handles indexed by $\Lambda_{\delta}^* = \{\kappa \in \Lambda^* \mid \|k_{\kappa}^+\| > 1/\delta\}$. The handle with index κ is contained in $\{k \in \mathbb{C}^2 \mid \|k - k_{\kappa}^+\| \leq \varepsilon\} = \{k \in \mathbb{C}^2 \mid \|k - k_{\kappa}^-\| \leq \varepsilon\} + \kappa$ and connects the small disc around k_{κ}^+ excluded from $\mathcal{V}_{\varepsilon, \delta}^+$ with the small disc around k_{κ}^- excluded from $\mathcal{V}_{\varepsilon, \delta}^-$. Finally, for all $q < \infty$ the Fourier components $\hat{\psi}(\kappa)$ of the periodic parts*

$$\tilde{\psi} = \exp(-2\pi\sqrt{-1}g(x, k))\psi = \sum_{\kappa \in \Lambda^*} \hat{\psi}(\kappa) \exp(2\pi\sqrt{-1}g(x, \kappa))$$

obey for some ε (depending on q) on these parts of $\mathcal{F}(V, W)$ the following estimates:

$$\left\| \tilde{\psi} - \begin{pmatrix} 0 \\ \hat{\psi}_2(0) \end{pmatrix} \right\|_q < \varepsilon |\hat{\psi}_2(0)| \text{ on } \mathcal{V}_{\varepsilon, \delta}^-, \quad \left\| \tilde{\psi} - \begin{pmatrix} \hat{\psi}_1(0) \\ 0 \end{pmatrix} \right\|_q < \varepsilon |\hat{\psi}_1(0)| \text{ on } \mathcal{V}_{\varepsilon, \delta}^+, \text{ and}$$

$$\left\| \tilde{\psi} - \begin{pmatrix} \hat{\psi}_1(0) \\ \hat{\psi}_2(-\kappa) \exp(-2\pi\sqrt{-1}g(x, \kappa)) \end{pmatrix} \right\|_q < \varepsilon \sqrt{|\hat{\psi}_2(0)|^2 + |\hat{\psi}_1(-\kappa)|^2}$$

on the small handle with index κ with respect to the wave vector k , which coincides at the border to $\mathcal{V}_{\varepsilon, \delta}^+$ with the wave vector of $\mathcal{V}_{\varepsilon, \delta}^+$, respectively. **q.e.d.**

The complex Fermi curves $\mathcal{F}(V, W)$ are locally finite sheeted coverings over $\hat{p} \in \mathbb{C}$ or $\check{p} \in \mathbb{C}$. If we enlarge the imaginary part of $\hat{p}$ (or $\check{p}$) we may connect any element of $\mathcal{F}(V, W)$ by a path either with some $\mathcal{V}_{\varepsilon, \delta}^-$ or with some $\mathcal{V}_{\varepsilon, \delta}^+$. Therefore, Theorem 2.12 implies

Corollary 2.13. *The regular part of $\mathcal{F}(V, W)/\Lambda^*$ has at most two connected components. Each component contains one of the two sets $\mathcal{V}_{\varepsilon, \delta}^{\pm}$ described in Theorem 2.12.* **q.e.d.**

This implies that the normalization of a complex Fermi curve also has at most two connected components, and each component contains one of the smooth open subsets $\mathcal{V}_{\varepsilon, \delta}^{\pm}$. The fixed energy level of the complex Bloch variety $\mathcal{B}(V, W)$ corresponding to the eigenvalue λ is equal to the complex Fermi curve $\mathcal{F}(V + \lambda, W + \lambda)$. Since the subset of the complex Bloch variety corresponding to degenerated eigenvalues is a subvariety, Theorem 2.12 implies

Corollary 2.14. *The complement of the subvariety of degenerated eigenvalues is open and dense in $\mathcal{B}(V, W)$.* **q.e.d.**

We shall see in Section 3.3 that if the normalization of a complex Fermi curve $\mathcal{F}(V, W)/\Lambda^*$ has disconnected normalization, then the first integral $4 \int_{\mathbb{R}^2/\Lambda} V(x)W(x)d^2x$ is a multiple of 4π . Therefore, Theorem 2.12 implies finally that the regular part of the complex Bloch variety $\mathcal{B}(V, W)/\Lambda^*$ is connected (i. e. the complex Bloch variety $\mathcal{B}(V, W)/\Lambda^*$ is irreducible).

2.3 Several reductions

In the last two sections we investigated the spectrum of the Dirac operator with two periodic complex-valued potentials V and W and introduced the *complex Fermi curve* being the spectral curve of the integrable system described by the Davey–Stewartson equation. The Weierstraß representation of immersions into $\mathbb{R}^3$ yields a one-to-one correspondence of Dirac operators $\begin{pmatrix} U & \partial \\ -\bar{\partial} & U \end{pmatrix}$ with real potentials U on $\mathbb{R}^2/\Lambda$, whose kernel contains non-trivial elements obeying the *Periodicity condition* 1.1. Hence we shall reduce the integrable system, whose spectral curves are the *complex Fermi curves* $\mathcal{F}(V, W)/\Lambda^*$, to the sub-system, whose spectral curves are the *complex Fermi curves* $\mathcal{F}(U, U)/\Lambda^*$ of a real potential U . Typically the reduced system is the set of fixed points of one or several involutions. Moreover, these involutions induce involutions of the corresponding spectral curves, and the spectral curves of the fixed points of the former involutions are invariant under the latter involutions.

In the sequel we will also consider the transposed Dirac operator $D^t(V, W, k)$. We will mostly denote the eigenfunction of the Dirac operator by $\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$ and the eigenfunction of the transposed Dirac operator by $\phi = \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}$. The transposition is always the one induced by the symmetric bilinear form

$$\langle\langle \phi, \psi \rangle\rangle = \int_{\Delta} (\phi_1(x)\psi_1(x) + \phi_2(x)\psi_2(x)) d^2x$$

on $L^2(\Delta) \times L^2(\Delta)$. Consequently the pointwise complex conjugation on this Hilbert space is used. Obviously the differential operators $\partial_{[k]}$ and $\bar{\partial}_{[k]}$ satisfy the relations

$$\partial_{[k]}^t = -\partial_{[-k]}, \bar{\partial}_{[k]}^t = -\bar{\partial}_{[-k]} \text{ and } \partial_{[k]}^* = -\bar{\partial}_{[\bar{k}]}, \bar{\partial}_{[k]}^* = -\partial_{[\bar{k}]}.$$

Hence we have

$$\begin{aligned} D^t(V, W, k) &= J^{-1}D(W, V, -k)J, \\ D^*(V, W, k) &= D(\bar{V}, \bar{W}, \bar{k}), \text{ and} \\ \bar{D}(V, W, k) &= J^{-1}D(\bar{W}, \bar{V}, -\bar{k})J, \end{aligned}$$

where J denotes the operator $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. This implies the following

Lemma 2.15. (i) *The holomorphic involution $\sigma : \mathbb{C}^3 \rightarrow \mathbb{C}^3, (k, \lambda) \mapsto (-k, \lambda)$, induces a biholomorphic isomorphism of the complex Bloch varieties $\mathcal{B}(V, W) \simeq \mathcal{B}(W, V)$.*

(ii) *The anti-holomorphic involution $\rho : \mathbb{C}^3 \rightarrow \mathbb{C}^3, (k, \lambda) \mapsto (\bar{k}, \bar{\lambda})$, induces a anti-biholomorphic isomorphism of the complex Bloch varieties $\mathcal{B}(V, W) \simeq \mathcal{B}(\bar{V}, \bar{W})$.*

(iii) *The anti-holomorphic involution $\eta : \mathbb{C}^3 \rightarrow \mathbb{C}^3, (k, \lambda) \mapsto (-\bar{k}, \bar{\lambda})$, induces a anti-biholomorphic isomorphism of the complex Bloch varieties $\mathcal{B}(V, W) \simeq \mathcal{B}(\bar{W}, \bar{V})$. **q.e.d.***

- Corollary 2.16.** (i) *The complex Bloch variety $\mathcal{B}(U, U)$ has a holomorphic involution $\sigma : (k, \lambda) \mapsto (-k, \lambda)$. Moreover, if ψ is an eigenfunction corresponding to the element (k, λ) of $\mathcal{B}(U, U)$ then $\phi = J\psi$ is an eigenfunction of the transposed Dirac operator corresponding to the element $\sigma(k, \lambda) = (-k, \lambda)$, and vice versa.*
- (ii) *If V and W are real valued periodic functions, the complex Bloch variety $\mathcal{B}(V, W)$ has an anti-holomorphic involution $\rho : (k, \lambda) \mapsto (\bar{k}, \bar{\lambda})$. Moreover, if ψ is an eigenfunction corresponding to the element (k, λ) of $\mathcal{B}(V, W)$ then $\phi = \bar{\psi}$ is an eigenfunction of the transposed Dirac operator corresponding to the element $\rho(k, \lambda) = (\bar{k}, \bar{\lambda})$, and vice versa.*
- (iii) *The complex Bloch variety $\mathcal{B}(U, \bar{U})$ has an anti-holomorphic involution $\eta : (k, \lambda) \mapsto (-\bar{k}, \bar{\lambda})$. Moreover, if ψ is an eigenfunction corresponding to the element (k, λ) of $\mathcal{B}(U, \bar{U})$ then $J\bar{\psi}$ is an eigenfunction of the complex conjugate of the Dirac operator corresponding to the element $\eta(k, \lambda) = (\bar{k}, \bar{\lambda})$, and vice versa. **q.e.d.***

The holomorphic involution σ induces a holomorphic involution of $\mathcal{F}(U, U)$, $\mathcal{F}(U, U)/\Lambda^*$, and of the normalization of $\mathcal{F}(U, U)/\Lambda^*$. All of them are also denoted by σ . If both potentials V and W are real, then the anti-holomorphic involution ρ induces an involution of $\mathcal{F}(V, W)$, $\mathcal{F}(V, W)/\Lambda^*$ and the normalization of $\mathcal{F}(V, W)/\Lambda^*$. Again these involutions are denoted by ρ . We call the fixed points of the involution ρ the real part of the *complex Fermi curve*. Consequently the real part of the normalization of the *complex Fermi curve* consists of the fixed points of the corresponding involution on the normalization.

- Corollary 2.17.** (i) *If V and W are real potentials, then all elements of the preimage of $\mathcal{F}(V, W)/\Lambda^* \cap \mathbb{R}^2/\Lambda^*$ under the normalization map, which are not fixed points of the involution ρ , are isolated points of the preimage of $\mathcal{F}(V, W)/\Lambda^* \cap \mathbb{R}^2/\Lambda^*$ under the normalization map.*
- (ii) *The involution η of the normalization of $\mathcal{F}(U, \bar{U})/\Lambda^*$ does not have any fixed point. More precisely, for all elements y of the normalization of $\mathcal{F}(U, U)/\Lambda^*$ the eigenfunctions corresponding to y and $\eta(y)$ are linearly independent.*
- (iii) *If $\mathcal{F}(U, \bar{U})/\Lambda^*$ contains an element of the form $[k]$ with $2\Re(k) \in \Lambda^*$, then the dimension of the eigenspace corresponding to this element is even.*

Proof. All elements of the real part of $\mathcal{F}(V, W)/\Lambda^*$ are obviously fixed points of the involution ρ . In particular, all regular points of the real part of $\mathcal{F}(V, W)/\Lambda^*$ are fixed points of ρ . Since the singular points of $\mathcal{F}(V, W)/\Lambda^*$ are isolated, this implies (i). Let y be some element of the normalization of $\mathcal{F}(U, \bar{U})/\Lambda^*$, which is a fixed point of the involution η . Due to Lemma 2.16 the eigenfunction ψ corresponding to this point has to be a multiple of $J\bar{\psi}$. But the equation $\psi = \alpha J\bar{\psi}$ implies $\psi = \alpha \bar{\alpha} J(J\bar{\psi})$. This would imply $\alpha \bar{\alpha} = -1$, which is impossible. This proves (ii). All elements of $\mathcal{F}(U, \bar{U})/\Lambda^*$ of the form $[k]$ with $2\Re(k) \in \Lambda^*$ are fixed points of the involution η . Due to Lemma 2.16 the eigenspace corresponding to this element $[k]$ is invariant under the anti-unitary map $\psi \mapsto J\bar{\psi}$. The square of this map is equal to $-\mathbf{1}$. Hence for all elements ψ of the eigenspace corresponding to $[k]$ the subspace spanned by ψ and $J\bar{\psi}$

is invariant under this map and two-dimensional, because $J\bar{\psi} = \alpha\psi$ would imply $\alpha\bar{\alpha} = -1$, which is impossible. Since the eigenspace corresponding to $[k]$ has to be a direct sum of subspaces of this form, its dimension is even. **q.e.d.**

Besides these involutions there exists some transformations of the potentials, which do not change the complex Bloch variety. In particular, all automorphisms of the group $\mathbb{R}^2/\Lambda$ induce transformations of the potentials, which do not change the complex Bloch variety. Moreover, the translations also induce transformations of the potentials, which do not change the complex Bloch variety (the corresponding transformations of the eigen bundle plays an important role in the inverse spectral theory (see e. g. [Kr-1, D-K-N])). Furthermore, for all $z \in \mathbb{C}^*$ the Dirac operator transforms like

$$D(z^2V, z^{-2}W, k) = \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix} D(V, W, k) \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix}.$$

Therefore, the *complex Fermi curves* $\mathcal{F}(zV, z^{-1}W)$ are independent of $z \in \mathbb{C}^*$. This action of the group $\mathbb{C}^*$ on the *isospectral sets* (i. e. the set of potentials, which have the same *complex Fermi curves*) is only the simplest example of a hierarchy of isospectral transformations [G-S-2], which may be constructed with the tools of soliton theory [Kr-1, D-K-N].

Finally, we remark that all affine symmetries of the torus $\mathbb{R}^2/\Lambda$ induce symmetries of the corresponding complex Bloch varieties. Besides the translations there exist in general only one, namely the parity transformation:

$$\mathbb{R}^2/\Lambda \rightarrow \mathbb{R}^2/\Lambda, x \mapsto -x.$$

This symmetry induces an isometry on $L^p(\mathbb{R}^2/\Lambda)$ and $L^p(\Delta)$, which is denoted by $\mathcal{P}$. By definition the operators $\partial_{[k]}$ and $\bar{\partial}_{[k]}$ transform as follows under this symmetry:

$$\mathcal{P}\partial_{[k]}\mathcal{P} = -\partial_{[-k]} = \partial_{[k]}^t \text{ and } \mathcal{P}\bar{\partial}_{[k]}\mathcal{P} = -\bar{\partial}_{[-k]} = \bar{\partial}_{[k]}^t.$$

This implies the following relations:

$$\begin{aligned} \begin{pmatrix} 0 & \mathcal{P} \\ \mathcal{P} & 0 \end{pmatrix} D(V, W, k) \begin{pmatrix} 0 & \mathcal{P} \\ \mathcal{P} & 0 \end{pmatrix} &= D^t(\mathcal{P}(W), \mathcal{P}(V), k), \\ \begin{pmatrix} 0 & \mathcal{P} \\ \mathcal{P} & 0 \end{pmatrix} \bar{D}(V, W, k) \begin{pmatrix} 0 & \mathcal{P} \\ \mathcal{P} & 0 \end{pmatrix} &= D(\mathcal{P}(\bar{W}), \mathcal{P}(\bar{V}), \bar{k}), \\ \begin{pmatrix} \mathcal{P} & 0 \\ 0 & -\mathcal{P} \end{pmatrix} D(V, W, k) \begin{pmatrix} \mathcal{P} & 0 \\ 0 & -\mathcal{P} \end{pmatrix} &= D(\mathcal{P}(V), \mathcal{P}(W), -k), \\ \begin{pmatrix} \mathcal{P} & 0 \\ 0 & -\mathcal{P} \end{pmatrix} \bar{D}(V, W, k) \begin{pmatrix} \mathcal{P} & 0 \\ 0 & -\mathcal{P} \end{pmatrix} &= D^t(\mathcal{P}(\bar{V}), \mathcal{P}(\bar{W}), -\bar{k}). \end{aligned}$$

Hence we obtain the following

Lemma 2.18. *The involutions $\mathbf{1}$, ρ , σ and η induce isomorphisms of the complex Bloch varieties $\mathcal{B}(V, W) \simeq \mathcal{B}(\mathcal{P}(W), \mathcal{P}(V))$, $\mathcal{B}(V, W) \simeq \mathcal{B}(\mathcal{P}(\bar{W}), \mathcal{P}(\bar{V}))$, $\mathcal{B}(V, W) \simeq \mathcal{B}(\mathcal{P}(V), \mathcal{P}(W))$ and $\mathcal{B}(V, W) \simeq \mathcal{B}(\mathcal{P}(\bar{V}), \mathcal{P}(\bar{W}))$, respectively. More precisely, if ψ is an eigenfunction of $D(V, W, k)$ with eigenvalue λ , then $\begin{pmatrix} 0 & \mathcal{P} \\ \mathcal{P} & 0 \end{pmatrix} \bar{\psi}$ and $\begin{pmatrix} \mathcal{P} & 0 \\ 0 & -\mathcal{P} \end{pmatrix} \psi$ are eigenfunctions of the operators $D(\mathcal{P}(\bar{W}), \mathcal{P}(\bar{V}), \bar{k})$ and $D(\mathcal{P}(V), \mathcal{P}(W), -k)$ with eigenvalues $\bar{\lambda}$ and λ , respectively. Moreover, $\begin{pmatrix} 0 & \mathcal{P} \\ \mathcal{P} & 0 \end{pmatrix} \psi$ and $\begin{pmatrix} \mathcal{P} & 0 \\ 0 & -\mathcal{P} \end{pmatrix} \bar{\psi}$ are transposed eigenfunctions of the operators $D(\mathcal{P}(W), \mathcal{P}(V), k)$ and $D(\mathcal{P}(\bar{V}), \mathcal{P}(\bar{W}), -\bar{k})$ with eigenvalues λ and $\bar{\lambda}$, respectively. In particular, if the pair of potentials is invariant under the transformation $(V, W) \mapsto (\mathcal{P}(W), \mathcal{P}(V))$, $(V, W) \mapsto (\mathcal{P}(\bar{W}), \mathcal{P}(\bar{V}))$, $(V, W) \mapsto (\mathcal{P}(V), \mathcal{P}(W))$ and $(V, W) \mapsto (\mathcal{P}(\bar{V}), \mathcal{P}(\bar{W}))$, then the complex Bloch variety is invariant under the corresponding involution $\mathbf{1}$, ρ , σ and η , respectively. **q.e.d.***

2.4 Spectral projections

In Section 2.2 the *complex Fermi curve* was introduced as a subvariety of $\mathbb{C}^2$. In this section we will work out a slightly different representation of this *complex Fermi curve*, which in Section 4.1 and Section 5.1 will turn out to be the key for the understanding of the *Periodicity condition* 1.1. As a preparation we investigate the spectral projections of the Dirac operator in some more detail.

Due to Theorem 2.3 and Corollary 2.14 locally there exist two meromorphic functions ψ and ϕ on the complex Bloch variety $\mathcal{B}(V, W)/\Lambda^*$, which map the elements $([k], \lambda)$ onto an eigenfunction of the Dirac operator $D(V, W, k)$ and an eigenfunction of the transposed Dirac operator $D^t(V, W, k)$, respectively. Moreover, these functions are unique up to multiplication by some invertible meromorphic function. Now we define a meromorphic function on $\mathcal{B}(V, W)/\Lambda^*$ with values in the finite rank operators on $L^2(\Delta) \times L^2(\Delta)$ by

$$P([k], \lambda) : \begin{pmatrix} \chi_1 \\ \chi_2 \end{pmatrix} \mapsto \frac{\langle \phi([k], \lambda), \chi \rangle}{\langle \phi([k], \lambda), \psi([k], \lambda) \rangle} \begin{pmatrix} \psi_1([k], \lambda) \\ \psi_2([k], \lambda) \end{pmatrix}.$$

Obviously this definition does not depend on the normalization of the functions ψ and ϕ . In fact, if we multiply both functions with some non-vanishing meromorphic functions, then the operator P does not change. Moreover, due to Corollary 2.14 the denominator does not vanish identically on the complex Bloch variety. Thus P is a well defined global meromorphic function on $\mathcal{B}(V, W)/\Lambda^*$. We will see that in some sense this projection can be considered as a projection-valued regular form.

On varieties with singularities the suitable generalization of holomorphic forms are the regular forms ([Se, Chapter IV §3.] and [Kun]). In general the corresponding sheaf is not the generalized cotangent sheaf [G-P-R, Chapter II. §1.2.]. Of particular interest are the regular forms of degree equal to the dimension of the variety. The sheaf of these regular forms is called dualizing sheaf [G-P-R, Chapter II. §5.3.] . In our case there exists a simple definition of the regular forms of degree equal to the dimension of the variety. In order to keep this paper self-contained we include a proof of the following lemma, which is a special case of a more general statement [G-P-R, Chapter II. Lemma 5.23].

Remark 2.19. *In the sequel we shall meet quite often complex spaces, which are locally bi-holomorphic to finite sheeted coverings over open subsets of $\mathbb{C}^n$. If we restrict these coverings to the preimage of small open balls, then different sheets without branch points are not connected with each other. However, in arbitrary small neighbourhoods of branch points several sheets are connected. In the sequel we shall call those sheets, whose restrictions to arbitrary small neighbourhoods of a given element contain this element, the local sheets, which contain this element.*

Lemma 2.20. *Let $R(z_0, \dots, z_n)$ be a holomorphic function on some open subset $U \subset \mathbb{C}^{n+1}$, whose partial derivative $\partial R / \partial z_0$ is not identically zero on the connected components of the subvariety of U defined by the equation $R(z_0, \dots, z_n) = 0$. Therefore this subvariety locally may be considered as a covering space over $(z_1, \dots, z_n) \in \mathbb{C}^n$. The following conditions on a meromorphic function f on this subvariety are equivalent:*

- (i) *The function f is holomorphic.*
- (ii) *For all holomorphic functions g the local sum of $gf/(\partial R/\partial z_0)$ over all sheets of the subvariety considered as a covering space over $(z_1, \dots, z_n) \in \mathbb{C}^n$, which contain an arbitrary element (compare with Remark 2.19), is a holomorphic function.*

Proof. Let us choose some element u of the subvariety. Due to the Weierstraß Preparation theorem [G-P-R, Chapter I. Theorem 1.4] locally there exists a polynomial Q with respect to z_0 , whose coefficients are holomorphic functions depending on $z_1, \dots, z_n$, such that R/Q is locally holomorphic. Moreover, the degree d of Q may be chosen to be equal to the number of all sheets of the subvariety considered as a covering space over $(z_1, \dots, z_n) \in \mathbb{C}^n$, which contain the element u . Near this element u the condition (ii) is obviously equivalent to an analogous condition, where R is replaced by Q . Let $Q(z_0)$ be an arbitrary polynomial of degree d , whose coefficients are complex numbers, and let $u_{0,1}, \dots, u_{0,d}$ be the zeroes of this polynomial. For an arbitrary polynomial $g(z_0)$ of degree less than d and with complex-valued coefficients we have the identity

$$g(z_0) = \sum_{i=1}^d \frac{g(u_{0,i})}{\partial Q(u_{0,i})/\partial z_0} \prod_{j \neq i} (z_0 - u_{0,j}).$$

In fact, if all zeroes of Q are pairwise different, the polynomials on both sides take the same values at all these zeroes and therefore have to be equal. In particular, the sum $\sum_i^d \frac{g(u_{0,i})}{\partial Q(u_{0,i})/\partial z_0}$ is equal to the coefficient of the monomial z_0^{d-1} in $g(z_0)$. For the monomials $g(z_0) = z_0^l$ this implies that

$$\sum_{i=1}^d \frac{u_{0,i}^l}{\partial Q(u_{0,i})/\partial z_0} = \begin{cases} 0 & \text{if } l < d-1 \\ 1 & \text{if } l = d-1. \end{cases}$$

We conclude that the same is true if the coefficients of Q are holomorphic functions depending on $z_1, \dots, z_n$, such that the discriminant of Q does not vanish identically. Due to the Weierstraß Preparation theorem [G-P-R, Chapter I. Theorem 1.4] each meromorphic function on the subvariety defined by the equation $Q = 0$ may be written locally as a polynomial with respect to z_0 of degree $d-1$, whose coefficients are meromorphic functions depending on $z_1, \dots, z_n$. Hence the local sum of this function over all sheets of the subvariety considered as a covering space over $(z_1, \dots, z_n) \in \mathbb{C}^n$, which contain u , is locally a holomorphic function, if and only if the coefficient corresponding to z_0^{d-1} of the polynomial is locally holomorphic. This shows that (i) and (ii) are equivalent. **q.e.d.**

If besides the partial derivative $\partial R/\partial z_0$ some other partial derivative $\partial R/\partial z_i$ is also not identically zero on the connected components of the subvariety of U defined by the equation $R(z_0, \dots, z_n) = 0$, then we may consider this subvariety locally either as a covering space over $(z_1, \dots, z_n) \in \mathbb{C}^n$ or as a covering space over $(z_0, \dots, z_{i-1}, z_{i+1}, \dots, z_n) \in \mathbb{C}^n$. The equation $R(z_0, \dots, z_n) = 0$ implies the relation $(\partial R/\partial z_0)dz_0 + \dots + (\partial R/\partial z_n)dz_n = 0$. We conclude that

the n -form $\frac{1}{\partial R/\partial z_0} dz_1 \wedge \dots \wedge dz_n$ is equal to the n -form $\frac{(-1)^i}{\partial R/\partial z_i} dz_0 \wedge \dots \wedge dz_{i-1} \wedge dz_{i+1} \wedge \dots \wedge dz_{n-1}$. Consequently we call a n -form

$$\omega = \frac{f(z_0, \dots, z_n)}{\partial R(z_0, \dots, z_n)/\partial z_0} dz_1 \wedge \dots \wedge dz_n =$$

$$(-1)^i \frac{f(z_0, \dots, z_n)}{\partial R(z_0, \dots, z_n)/\partial z_i} dz_0 \wedge \dots \wedge dz_{i-1} \wedge dz_{i+1} \wedge \dots \wedge dz_{n-1}$$

regular, if and only if f satisfies one of the equivalent conditions (i), (ii) or the analogous condition to (ii), when z_0 is replaced by z_i ([Se, Chapter IV §3.], [Kun] and [G-P-R, Chapter II. Definition 5.22]).

Lemma 2.21. *The function P has the following properties:*

- (i) *The values of P are projections of rank one.*
- (ii) *If $([k], \lambda)$ and $([k], \lambda')$ are two elements of the complex Bloch variety with $\lambda \neq \lambda'$, then $P([k], \lambda)P([k], \lambda') = 0 = P([k], \lambda')P([k], \lambda)$.*
- (iii) *Due to Theorem 2.3 the complex Bloch variety is locally a Weierstraß covering (see e.g. [G-P-R, Chapter I. §12.3]) over $k \in \mathbb{C}^2$. The local sum of P over all sheets of this covering, which contain some element $([k], \lambda) \in \mathcal{B}(V, W)/\Lambda^*$ (compare with Remark 2.19), is a holomorphic function on some neighbourhood of $k \in \mathbb{C}^2$. Moreover, the values of this function are projections, whose rank is equal to the number of sheets, over which that sum is taken. Finally, the value of this function at $([k], \lambda)$ is equal to the spectral projection $\tilde{P}([k], \lambda)$ of the Dirac operator $D(V, W, k)$ onto the generalized eigenspace associated with eigenvalue λ (see e. g. [R-S-IV, Appendix to XII.1]).*
- (iv) *The projection-valued form $Pdk_1 \wedge dk_2$ is regular.*

Proof. The first statement is obvious. Due to Theorem 2.3 locally we may replace the Dirac operator $D(V, W, k)$ by some holomorphic matrix-valued function $A(k)$ on some open subset $k \in \mathbb{C}^2$. Due to Corollary 2.14 this function generically has pairwise different eigenvalues. If A is a $l \times l$ matrix with pairwise different eigenvalues $\lambda_1, \dots, \lambda_l$, then the eigenvectors $\psi_1, \dots, \psi_l$ to the eigenvalues $\lambda_1, \dots, \lambda_l$ form a basis of $\mathbb{C}^2$. Moreover, the dual basis $\phi_1, \dots, \phi_l$ gives the eigenvectors of A^t to the eigenvalues $\lambda_1, \dots, \lambda_l$. In fact, if $\tilde{\phi}_i$ is some eigenvector of A^t with eigenvalue λ_i , then we have $\tilde{\phi}_i^t \psi_j (\lambda_i - \lambda_j) = \left(A^t \tilde{\phi}_i \right)^t \psi_j - \tilde{\phi}_i^t A \psi_j = 0$ for all $1 \leq i, j \leq l$ and $\tilde{\phi}_i$ has to be proportional to ϕ_i . We conclude that for all $1 \leq i \neq j \leq l$ the eigen projections

$$P_i = \frac{\psi_i \phi_i^t}{\phi_i^t \psi_i} \quad i = 1, \dots, l$$

satisfy the relations $P_i P_j = 0 = P_j P_i$. This proves (ii). On the other hand $P_1 + \dots + P_l$ is equal to the identity matrix and (iii) is proven. Due to Theorem 2.3 the complex Bloch variety is locally a Weierstraß covering (see e.g. [G-P-R, Chapter I. §12.3]) over $k \in \mathbb{C}^2$. Hence any

holomorphic function f on the complex Bloch variety is locally equal to some polynomial with respect to λ , whose coefficients are holomorphic functions on some open subset of $k \in \mathbb{C}^2$. Again due to Theorem 2.3 locally the sum over the sheets of this Weierstraß covering of $\mathbf{P}$ times any holomorphic function is a holomorphic function on some open subset of $k \in \mathbb{C}^2$ with values in the finite rank operators on $L^2(\Delta) \times L^2(\Delta)$. Hence the function $\mathbf{P}$ satisfies condition (ii) of Lemma 2.20. **q.e.d.**

Since the Dirac operator is a holomorphic unbounded-operator-valued function depending on the wave vectors $k \in \mathbb{C}^2$, the complex Bloch variety has naturally the structure of a covering space over $k \in \mathbb{C}^2$. But in order to understand the *complex Fermi curve*, which is our main interest, it would be more appropriate to consider the complex Bloch variety as a covering space over λ and some component of k , because such a covering would just have to be restricted to the plane $\lambda = 0$ to obtain the *complex Fermi curve*. In the remainder of this section this point of view will be worked out. The starting point is the following observation, which generalizes the equivalence of the *Periodicity condition* 1.1. Here we use the notations introduced in the context of the *Fundamental domain* 2.1, but we shall omit the index Δ .

Lemma 2.22. *Let $\psi(x) = \begin{pmatrix} \psi_1(x) \\ \psi_2(x) \end{pmatrix}$ be an eigenfunction corresponding to an element (k, λ) of $\mathcal{B}(V, W)$ and $\phi(x) = \begin{pmatrix} \phi_1(x) \\ \phi_2(x) \end{pmatrix}$ an eigenfunction of the transposed Dirac operator corresponding to an element $(k', \lambda) \in \mathcal{B}(V, W)$. Then $\phi_2 \psi_1 dz + \phi_1 \psi_2 d\bar{z}$, with $dz = dx_1 + \sqrt{-1}dx_2$ and $d\bar{z} = dx_1 - \sqrt{-1}dx_2$, is a closed form. If the two components $\hat{p} = g(\hat{\gamma}, k)$ and $\hat{p}' = g(\hat{\gamma}, k')$ of k and k' coincide, then the integral of this form from some point $x \in \mathbb{R}^2$ to $x + \hat{\gamma}$ does not depend on x , and is equal to*

$$\frac{(\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2) \langle \phi_2, \psi_1 \rangle + (\hat{\gamma}_1 - \sqrt{-1}\hat{\gamma}_2) \langle \phi_1, \psi_2 \rangle}{\hat{\gamma}_1 \check{\gamma}_2 - \hat{\gamma}_2 \check{\gamma}_1}.$$

If in addition the elements $[k]$ and $[k']$ of $\mathbb{C}^2/\Lambda^$ are different, then both integrals vanish.*

Proof. Since ψ is an eigenfunction of the Dirac operator and ϕ is a eigenfunction of the transposed Dirac operator with eigenvalues zero, we have

$$\begin{aligned} V\psi_1 + \partial\psi_2 &= \lambda\psi_1 & V\phi_1 + \bar{\partial}\phi_2 &= \lambda\phi_1 \\ -\bar{\partial}\psi_1 + W\psi_2 &= \lambda\psi_2 & -\partial\phi_1 + W\phi_2 &= \lambda\phi_2. \end{aligned}$$

Thus

$$d(\phi_2 \psi_1 dz + \phi_1 \psi_2 d\bar{z}) = ((\bar{\partial}\phi_2) \psi_1 + \phi_2 (\bar{\partial}\psi_1) - (\partial\phi_1) \psi_2 - \phi_1 (\partial\psi_2)) d\bar{z} \wedge dz = 0.$$

If the two components $\hat{p}$ and $\hat{p}'$ of k and k' coincide, then this closed form is invariant under the shift by the period $\hat{\gamma}$. This implies that the integral of this form along some path from some arbitrary point $x \in \mathbb{R}^2$ to $x + \hat{\gamma}$, does not depend on x . In particular, this integral is

equal to the integral

$$\int_{[0,1]^2} ((\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2) \phi_2(x)\psi_1(x) + (\hat{\gamma}_1 - \sqrt{-1}\hat{\gamma}_2) \phi_1(x)\psi_2(x)) d\hat{q} \wedge d\check{q} = \frac{(\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2) \langle \phi_2, \psi_1 \rangle + (\hat{\gamma}_1 - \sqrt{-1}\hat{\gamma}_2) \langle \phi_1, \psi_2 \rangle}{\hat{\gamma}_1\check{\gamma}_2 - \hat{\gamma}_2\check{\gamma}_1}.$$

If the components with respect to the other period $\check{\gamma}$ are different $\check{p} \neq \check{p}' \pmod{\mathbb{Z}}$, then the shift by $\check{\gamma}$ changes the form by the factor $\exp(2\pi\sqrt{-1}(\check{p} - \check{p}')) \neq 1$. Hence the integral of this form along $\hat{\gamma}$ has to vanish. **q.e.d.**

This lemma motivates the following definition: For all $\mu \in \mathbb{C}^2$, let $\langle \langle \cdot, \cdot \rangle \rangle_\mu$ be the following bilinear form on the Hilbert spaces $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ and $L^2(\Delta) \times L^2(\Delta)$, respectively:

$$\langle \langle \phi, \psi \rangle \rangle_\mu := (\mu_1 + \sqrt{-1}\mu_2) \langle \phi_2, \psi_1 \rangle + (\mu_1 - \sqrt{-1}\mu_2) \langle \phi_1, \psi_2 \rangle.$$

The complex Bloch variety is locally a finite-sheeted covering over $(\hat{p}, \lambda) \in \mathbb{C}^2$, and the zeroes of the differential $d\hat{p} \wedge d\lambda$ are the branch points of this covering. Now we define another projection, which is related to this covering. Again we use the notations introduced in the context of the *Fundamental domain* 2.1. Furthermore, we again use the meromorphic functions $\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$ and $\phi = \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}$ on $\mathcal{B}(V, W)/\Lambda^*$. Let $P_{\hat{\gamma}}$ be the meromorphic map from $\mathcal{B}(V, W)/\Lambda^*$ into the finite rank operators on the Hilbert space $L^2(\Delta) \times L^2(\Delta)$, which for all elements $([k], \lambda)$ of the complex Bloch variety is defined by

$$P_{\hat{\gamma}}([k], \lambda) : \begin{pmatrix} \chi_1 \\ \chi_2 \end{pmatrix} \longmapsto \frac{\langle \langle \phi([k], \lambda), \chi \rangle \rangle_{\hat{\gamma}}}{\langle \langle \phi([k], \lambda), \psi([k], \lambda) \rangle \rangle_{\hat{\gamma}}} \begin{pmatrix} \psi_1([k], \lambda) \\ \psi_2([k], \lambda) \end{pmatrix}.$$

Obviously this definition does not depend on the normalization of the functions ψ and ϕ . In fact, if we multiply both functions with some non-vanishing meromorphic functions, then the operator $P_{\hat{\gamma}}$ does not change. Moreover, due to proof of Theorem 2.12 and Corollary 2.13 the denominator does not vanish identically on the complex Bloch variety. Thus $P_{\hat{\gamma}}$ is a well defined global meromorphic function on $\mathcal{B}(V, W)/\Lambda^*$.

Lemma 2.23. *The function $P_{\hat{\gamma}}$ has the following properties:*

- (i) *The values of $P_{\hat{\gamma}}$ are projections of rank one.*
- (ii) *If $([k], \lambda)$ and $([k'], \lambda)$ are two different elements of the complex Bloch variety with $g(\hat{\gamma}, k) = g(\hat{\gamma}, k') \pmod{\mathbb{Z}}$, then $P_{\hat{\gamma}}([k], \lambda)P_{\hat{\gamma}}([k'], \lambda) = 0 = P_{\hat{\gamma}}([k'], \lambda)P_{\hat{\gamma}}([k], \lambda)$.*
- (iii) *Locally the complex Bloch variety is a finite-sheeted covering over $(\hat{p}, \lambda) \in \mathbb{C}^2$. The local sum of $P_{\hat{\gamma}}$ over all sheets, which contain one element $([k], \lambda)$ (compare with Remark 2.19), is a holomorphic function on some open subset of $(\hat{p}, \lambda) \in \mathbb{C}^2$, with values in the finite rank projections on $L^2(\Delta) \times L^2(\Delta)$. Moreover, the rank of any of these projections is equal to the number of sheets, over which the sum is taken.*

- (iv) The 2-form $P_{\hat{\gamma}} d\hat{p} \wedge d\lambda$ is a regular form on the complex Bloch variety and the one-form $P_{\hat{\gamma}} d\hat{p}$ is a regular form on the complex Fermi curve.
- (v) Due to (iii) for all elements $([k], \lambda)$ of $\mathcal{B}(V, W)/\Lambda^*$ there exists a unique projection $\check{P}_{\hat{\gamma}}([k], \lambda)$, which is the value at $([k], \lambda)$ of the local sum of $P_{\hat{\gamma}}$ over all sheets of $\mathcal{B}(V, W)/\Lambda^*$, which contain $([k], \lambda)$. For two different elements $([k], \lambda)$ and $([k'], \lambda)$ of $\mathcal{B}(V, W)/\Lambda^*$ with equal components $g(\hat{\gamma}, k) = g(\hat{\gamma}, k') \pmod{\mathbb{Z}}$ the corresponding projections are disjoint: $\check{P}_{\hat{\gamma}}([k], \lambda)\check{P}_{\hat{\gamma}}([k'], \lambda) = 0 = \check{P}_{\hat{\gamma}}([k'], \lambda)\check{P}_{\hat{\gamma}}([k], \lambda)$.
- (vi) If χ is a proper eigenfunction of the Dirac operator corresponding to some element $([k], \lambda)$ of the complex Bloch variety, then the range of $\check{P}_{\hat{\gamma}}([k], \lambda)$ contains χ .

We remark that the range of the spectral projections contain all elements in the kernels of $(\lambda \mathbf{1} - D(V, W, k))^l$ with $l \in \mathbb{N}$.

Proof. The first statement is obvious. Statement (ii) is a direct consequence of Lemma 2.22. If we consider the Dirac operator as an operator acting on the Hilbert bundle introduced in the context of the *Trivialization* 2.2, then the partial derivative $\partial \tilde{D}(V, W, k)/\partial \check{p}$ is equal to

$$\frac{\partial \tilde{D}(V, W, k)}{\partial \check{p}} = \pi \begin{pmatrix} 0 & (\check{\kappa}_2 + \sqrt{-1}\check{\kappa}_1) \mathbf{1} \\ (\check{\kappa}_2 - \sqrt{-1}\check{\kappa}_1) \mathbf{1} & 0 \end{pmatrix}.$$

We conclude that

$$\frac{\partial \lambda}{\partial \check{p}} \langle \langle \phi([k], \lambda), \psi([k], \lambda) \rangle \rangle = \frac{\pi}{\hat{\gamma}_1 \check{\gamma}_2 - \hat{\gamma}_2 \check{\gamma}_1} \langle \langle \phi([k], \lambda), \psi([k], \lambda) \rangle \rangle_{\hat{\gamma}},$$

because $\hat{\kappa}$ and $\check{\kappa}$ form the dual basis of $\hat{\gamma}$ and $\check{\gamma}$, and therefore $\check{\kappa}$ is equal to $\frac{1}{\hat{\gamma}_1 \check{\gamma}_2 - \hat{\gamma}_2 \check{\gamma}_1} (-\hat{\gamma}_2, \hat{\gamma}_1)$. Since the zeroes and poles of the numerator of $P_{\hat{\gamma}}$ and P coincide, condition (iv) of Lemma 2.21 implies that $P_{\hat{\gamma}} d\hat{p} \wedge d\lambda = (\partial \lambda / \partial \check{p}) P_{\hat{\gamma}} d\hat{p} \wedge d\check{p}$ is a regular form on the complex Bloch variety with values in the projections of rank one on $L^2(\Delta) \times L^2(\Delta)$. This shows that the local sum of $P_{\hat{\gamma}}$ over all those sheets of $\mathcal{B}(V, W)/\Lambda^*$ considered as a covering space over $(\hat{p}, \lambda) \in \mathbb{C}^2$, which contain $([k], \lambda)$, is holomorphic. Finally, Lemma 2.22 implies that the values of this sum are projections, whose rank is equal to the number of sheets. Due to Theorem 2.12 and Corollary 2.13 the restrictions of the sheets of the complex Bloch variety considered as a covering space over $(\hat{p}, \lambda) \in \mathbb{C}^2$ to the *complex Fermi curve* are locally different. Hence the fourth statement (iv) is a consequence of (iii) and Lemma 2.20. Condition (v) is a direct consequence of (iii). As for the proof of (vi), let χ be an eigenfunction of the Dirac operator $D(V, W, k)$ with eigenvalue λ . The exterior derivative of the form $\phi_2([k'], \lambda') \chi_1 dz + \phi_1([k'], \lambda') \chi_2 d\bar{z}$ is equal to

$$d(\phi_2([k'], \lambda') \chi_1 dz + \phi_1([k'], \lambda') \chi_2 d\bar{z}) = (\lambda' - \lambda) (\phi_1([k'], \lambda') \chi_1 + \phi_2([k'], \lambda') \chi_2) d\bar{z} \wedge dz.$$

If furthermore $g(\hat{\gamma}, k) = g(\hat{\gamma}, k') \pmod{\mathbb{Z}}$, we may apply Lemma 2.22 and conclude

$$\begin{aligned} & \frac{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k' - k)) - 1}{\hat{\gamma}_1\check{\gamma}_2 - \hat{\gamma}_2\check{\gamma}_1} \langle \langle \phi([k'], \lambda'), \chi \rangle \rangle_{\hat{\gamma}} = \\ & (\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k' - k)) - 1) \int_x^{x+\hat{\gamma}} \phi_2([k'], \lambda') \chi_1 dz + \phi_1([k'], \lambda') \chi_2 d\bar{z} = \\ & 2\sqrt{-1}(\lambda' - \lambda) \langle \langle \phi([k'], \lambda'), \chi \rangle \rangle. \end{aligned}$$

Hence, due to the relation between the denominators of the projections $\mathbf{P}$ and $\mathbf{P}_{\hat{\gamma}}$ established in the proof of (iii), the restrictions of the following two forms to the subvariety $\hat{p}' = \hat{p} = g(\hat{\gamma}, k) \pmod{\mathbb{Z}}$ of $\mathcal{B}(V, W)/\Lambda^*$ coincide:

$$\frac{2\pi\sqrt{-1}d\check{p}'}{\exp(2\pi\sqrt{-1}(\check{p}' - \check{p})) - 1} \mathbf{P}([k'], \lambda') \chi = \frac{d\lambda'}{\lambda' - \lambda} \mathbf{P}_{\hat{\gamma}}([k'], \lambda') \chi.$$

The subvariety $\{(\hat{p}', \lambda') \in \mathbb{C}^2 \mid (\hat{p}'\hat{\kappa} + \check{p}\check{\kappa}, \lambda') \in \mathcal{B}(V, W)\}$ of the complex Bloch variety may be considered locally either as a covering space over $\hat{p}' \in \mathbb{C}$ or as a covering space over $\lambda' \in \mathbb{C}$. The residue of the left hand side is the value of the local sum of $\mathbf{P}\chi$ over all sheets, which contains $(\hat{p}, \lambda)$, of the subvariety considered as a covering over $\hat{p}' \in \mathbb{C}$, and the residue of the right hand side is the value of the local sum of $\mathbf{P}_{\hat{\gamma}}\chi$ over all sheets, which contains $(\hat{p}, \lambda)$, of the subvariety considered as a covering over $\lambda' \in \mathbb{C}$. Obviously the restrictions of the sheets of $\mathcal{B}(V, W)$ considered as a covering space over $k \in \mathbb{C}^2$ to the subvariety $g(\hat{\gamma}, k) = \text{constant}$ are locally different and also the restrictions of the sheets of $\mathcal{B}(V, W)$ considered as a covering space over $(\hat{p}, \lambda) \in \mathbb{C}^2$ to the subvariety $\hat{p} = \text{constant}$ are locally different. Hence the residue of the left hand side is equal to $\check{\mathbf{P}}([k], \lambda)\chi$ and the residue of the right hand side is equal to $\check{\mathbf{P}}_{\hat{\gamma}}([k], \lambda)\chi$. This proves (vi). **q.e.d.**

These Lemmata have some remarkable consequences. If we fix the component $\hat{p} = \hat{p}'$ of the wave vectors k and the energy $\lambda = \lambda'$, then all eigenfunctions χ of the Dirac operator with eigenvalue $\lambda = \lambda'$ and boundary condition $\chi(\hat{q}, \check{q}) = \exp(2\pi\sqrt{-1}\hat{q}\hat{p}') \chi(\hat{q} + 1, \check{q})$ are uniquely determined by their restriction to some line $\check{q} = \check{q}'$. In fact, if the component $\hat{p}$ of k is equal to $\hat{p}'$, then due to Lemma 2.22 the action of the projection $\check{\mathbf{P}}_{\hat{\gamma}}([k])$ defined in (v) of Lemma 2.23 on χ depends only on the restriction to $\check{q} = \check{q}'$.

For reasons of simplicity we restrict ourself in the following discussion to the *complex Fermi curve*, which is our main interest. The restrictions of the functions $\mathbf{P}$ and $\mathbf{P}_{\hat{\gamma}}$ to the *complex Fermi curve* are also denoted by $\mathbf{P}$ and $\mathbf{P}_{\hat{\gamma}}$, respectively. If we identify the Hilbert space $L^2(\mathbb{R}/\mathbb{Z}) \times L^2(\mathbb{R}/\mathbb{Z})$ with all L^2 functions on some line $\hat{q} \in \mathbb{R}/\mathbb{Z}, \check{q} = \check{q}'$, then the composition of the operator

$$\begin{pmatrix} \chi_1 \\ \chi_2 \end{pmatrix} \longmapsto \frac{\int_{\hat{q} \in \mathbb{R}/\mathbb{Z}, \check{q} = \check{q}'} \phi_2([k]) \chi_1 dz + \phi_1([k]) \chi_2 d\bar{z}}{\int_{\hat{q} \in \mathbb{R}/\mathbb{Z}, \check{q} = \check{q}'} \phi_2([k]) \psi_1([k]) dz + \phi_1([k]) \psi_2([k]) d\bar{z}} \begin{pmatrix} \psi_1([k]) \\ \psi_2([k]) \end{pmatrix}$$

from $L^2(\mathbb{R}/\mathbb{Z}) \times L^2(\mathbb{R}/\mathbb{Z})$ into $L^2(\Delta) \times L^2(\Delta)$ and the operator, which restricts the functions to the line $\hat{q} \in \mathbb{R}/\mathbb{Z}$, $\check{q} = \check{q}' + 1$ defines an operator on the Hilbert space $L^2(\mathbb{R}/\mathbb{Z}) \times L^2(\mathbb{R}/\mathbb{Z})$. This restriction is not defined for all L^2 -functions. But for all eigenfunctions of the Dirac operators it is. In fact, due to Remark 2.7 the eigenfunctions belong to the Sobolev spaces $W^{1,p}(\Delta) \times W^{1,p}(\Delta)$ with $1 < p < 2$. Due to the Sobolev embedding theorem [Ad, 5.4 Theorem] this implies that the restrictions of the eigenfunctions to a one-dimensional plane in $\mathbb{R}^2/\Lambda$ are L^q -functions for all $q < \infty$. The result is some finite rank operator-valued function on the *complex Fermi curve*, whose eigenvalues are equal to $\exp(2\pi\sqrt{-1}g(\check{\gamma}, k)) = \exp(2\pi\sqrt{-1}\check{p})$. Moreover, due to Lemma 2.23 the local sum of this function over all sheets of $\mathcal{F}(V, W)$ considered as a covering space over $\hat{p} \in \mathbb{C}$, which contain an element $k' \in \mathcal{F}(V, W)$ (compare with Remark 2.19), is locally a holomorphic function depending on $\hat{p}$. It can be proven that the sum of these operators over all sheets of $\mathcal{F}(V, W)$ considered as a covering space over $\hat{p} \in \mathbb{C}$ defines a holomorphic function from $\mathbb{C}$ into the unbounded closed operators on $L^2(\mathbb{R}/\mathbb{Z}) \times L^2(\mathbb{R}/\mathbb{Z})$, which is periodic with period 1. The *complex Fermi curve* is the complex Bloch variety of this entire function. Due to Lemma 2.22, the restriction of the projection $P_{\check{\gamma}}$ to the Hilbert space $L^2(\mathbb{R}/\mathbb{Z}) \times L^2(\mathbb{R}/\mathbb{Z})$ is equal to the composition of the operator

$$\begin{pmatrix} \chi_1 \\ \chi_2 \end{pmatrix} \mapsto \frac{\int_{\hat{q} \in \mathbb{R}/\mathbb{Z}, \check{q} = \check{q}'} \phi_2([k])\chi_1 dz + \phi_1([k])\chi_2 d\bar{z}}{\int_{\hat{q} \in \mathbb{R}/\mathbb{Z}, \check{q} = \check{q}'} \phi_2([k])\psi_1([k])dz + \phi_1([k])\psi_2([k])d\bar{z}} \begin{pmatrix} \psi_1([k]) \\ \psi_2([k]) \end{pmatrix}$$

from $L^2(\mathbb{R}/\mathbb{Z}) \times L^2(\mathbb{R}/\mathbb{Z})$ into $L^2(\Delta) \times L^2(\Delta)$ and the operator, which restricts the functions to the line $\hat{q} \in \mathbb{R}/\mathbb{Z}$, $\check{q} = \check{q}'$. These projections may be considered as the spectral projection of the operator-valued entire function depending on $\hat{p}$ like P is the spectral projection of the Dirac operator. In particular, the sum of $P_{\check{\gamma}}$ considered as an projection on $L^2(\mathbb{R}/\mathbb{Z}) \times L^2(\mathbb{R}/\mathbb{Z})$ over all sheets of $\mathcal{F}(V, W)$ considered as a covering space over $\hat{p} \in \mathbb{C}$ converges in the strong operator topology to the identity.

All operator-valued holomorphic functions depending on $\hat{p} \in \mathbb{C}$, which commute pointwise with this unbounded-operator-valued entire function, may be diagonalized simultaneously with this entire function. The eigenvalues of such commuting operator-valued functions defines the structure sheaf of a one-sheeted covering over the *complex Fermi curve*:

Structure sheaf 2.24. *Due to [Gr-Re, Chapter 8. §3.] there exists an up to isomorphism unique one-sheeted covering of the complex Fermi curve, whose direct image of the Structure sheaf yields the following coherent subsheaf of the normalization sheaf (which is the direct image of the structure sheaf of the normalization under the normalization map): A germ f of the normalization sheaf belongs to the direct image of the Structure sheaf, if and only if the local sum of $fP_{\check{\gamma}}$ over all local sheets of the covering map $\hat{p}$ of the complex Fermi curve, which contain the base point of the corresponding stalk (compare with Remark 2.19), is a germ of the sheaf of operator-valued holomorphic functions depending on $\hat{p} \in \mathbb{C}$.*

Obviously the normalization of this one sheeted covering is isomorphic to the normalization of the *complex Fermi curve*. To sum up the complex space with *Structure sheaf 2.24* is a one-sheeted covering over the *complex Fermi curve* considered as a subvariety of $\mathbb{C}^2/\Lambda^*$, and

the normalization is a one-sheeted covering over the former space. Therefore, we distinguish between the

geometric genus of the normalization, the

arithmetic genus of the *Structure sheaf* 2.24, and, finally, the

arithmetic genus of the *complex Fermi curve*.

We shall see in Proposition 3.52 that the last genus is always infinite and in some sense locally preserved. Moreover, in Lemma 2.26 we shall see that for pairs of potentials of the form $(U, \bar{U})$ the difference of the *arithmetic genus* of the *Structure sheaf* 2.24 minus the *geometric genus* is finite. Potentials, whose *Structure sheaf* 2.24 have finite arithmetic genus, are called *finite type potentials* (compare with Section 2.5).

It can be shown with similar arguments to those used in the proof of Lemma 2.23 (vi), that this *Structure sheaf* does not depend on the choice of $\hat{\gamma}$ and $\check{\gamma}$. This one-sheeted covering would be a more appropriate definition of the *complex Fermi curve*. For example, the degree of the dual eigen bundle is equal to the arithmetic genus of this curve plus one [Sch].

So we have investigated the structure of the *complex Fermi curve* as a covering space over some coordinate $\hat{p} = g(\hat{\gamma}, k)$. Let us generalize this construction to the more general case, when the period $\hat{\gamma}$ is replaced by some arbitrary element $\mu \in \mathbb{C}^2$. The starting point is again some generalization of the equivalence of the *Periodicity condition* 1.1. For this purpose, however, we use the *Trivialization* 2.2 and the eigenfunctions are described by their periodic parts in the corresponding Hilbert bundle.

Lemma 2.25. *Let $\psi = \begin{pmatrix} \psi_1(x) \\ \psi_2(x) \end{pmatrix}$ be an eigenfunction corresponding to an element (k, λ) of $\mathcal{B}(V, W)$ and $\phi = \begin{pmatrix} \phi_1(x) \\ \phi_2(x) \end{pmatrix}$ an eigenfunction of the transposed Dirac operator corresponding to an element $(k', \lambda) \in \mathcal{B}(V, W)$, and let $\tilde{\psi}(x) = \exp(-2\pi\sqrt{-1}g(x, k))\psi(x)$ and $\tilde{\phi}(x) = \exp(2\pi\sqrt{-1}g(x, k'))\phi(x)$ the corresponding periodic parts. Then the exterior derivative of the form $\tilde{\phi}_2\tilde{\psi}_1dz + \tilde{\phi}_1\tilde{\psi}_2d\bar{z}$ is equal to*

$$\pi \left((k_2 - k'_2 - \sqrt{-1}(k_1 - k'_1)) \tilde{\phi}_2\tilde{\psi}_1 + (k_2 - k'_2 + \sqrt{-1}(k_1 - k'_1)) \tilde{\phi}_1\tilde{\psi}_2 \right) d\bar{z} \wedge dz.$$

Therefore, the bilinear form $\langle\langle \tilde{\phi}, \tilde{\psi} \rangle\rangle_\mu$ vanishes, whenever $g(\mu, k - k') = 0$ but $k - k' \neq 0$.

Proof. Since ψ is an eigenfunction of the Dirac operator and ϕ is a eigenfunction of the transposed Dirac operator with eigenvalues zero, we have

$$\begin{aligned} V\tilde{\psi}_1 + (\pi(k_2 + \sqrt{-1}k_1) + \partial)\tilde{\psi}_2 &= \lambda\tilde{\psi}_1 & V\tilde{\phi}_1 + (\pi(k'_2 - \sqrt{-1}k'_1) + \bar{\partial})\tilde{\phi}_2 &= \lambda\tilde{\phi}_1 \\ (\pi(k_2 - \sqrt{-1}k_1) - \bar{\partial})\tilde{\psi}_1 + W\tilde{\psi}_2 &= \lambda\tilde{\psi}_2 & (\pi(k'_2 + \sqrt{-1}k'_1) - \partial)\tilde{\phi}_1 + W\tilde{\phi}_2 &= \lambda\tilde{\phi}_2. \end{aligned}$$

Thus the exterior derivative of $d(\tilde{\phi}_2\tilde{\psi}_1dz + \tilde{\phi}_1\tilde{\psi}_2d\bar{z})$ is equal to

$$\pi \left((k_2 - k'_2 - \sqrt{-1}(k_1 - k'_1)) \tilde{\phi}_2\tilde{\psi}_1 + (k_2 - k'_2 + \sqrt{-1}(k_1 - k'_1)) \tilde{\phi}_1\tilde{\psi}_2 \right) d\bar{z} \wedge dz.$$

Since both forms are periodic, the integral of the right hand side over the torus $\mathbb{R}^2/\Lambda$ vanishes. This integral is equal to $2\pi\sqrt{-1}\langle\langle\tilde{\phi}, \tilde{\psi}\rangle\rangle_\mu$ if μ is equal to $(k'_2 - k_2, k_1 - k'_1)$, which implies that $g(\mu, k - k') = 0$. **q.e.d.**

This lemma implies that whenever for some element $\mu \in \mathbb{C}^2$ the components $g(\mu, k)$ and $g(\mu, k')$ coincide, the bilinear form $\langle\langle\tilde{\phi}, \tilde{\psi}\rangle\rangle_\mu$ vanishes. But for this generalization we have to pay a price. Since instead of the eigenfunctions we used the periodic part of the eigenfunctions, we lose invariance under the action of the dual lattice. Now we can also define for general $\mu \in \mathbb{C}^2$ a projection valued function $\tilde{P}_\mu$, but this function should be considered to act not on the Hilbert space, but on the Hilbert bundle introduced in the *Trivialization* 2.2. Let $\tilde{P}_\mu$ be the meromorphic map from $\mathcal{B}(V, W)/\Lambda^*$ into the finite rank operators on the Hilbert space $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$, which for all elements $([k], \lambda)$ of the complex Bloch variety is defined by

$$\tilde{P}_\mu([k], \lambda) : \begin{pmatrix} \tilde{\chi}_1 \\ \tilde{\chi}_2 \end{pmatrix} \longmapsto \frac{\langle\langle\phi([k], \lambda), \psi_k \tilde{\chi}\rangle\rangle_\mu}{\langle\langle\phi([k], \lambda), \psi([k], \lambda)\rangle\rangle_\mu} \begin{pmatrix} \psi_{-k}\psi_1([k], \lambda) \\ \psi_{-k}\psi_2([k], \lambda) \end{pmatrix}.$$

Here ψ_k is the operator-valued holomorphic function on the complex Bloch variety, which is given by the multiplication with the function ψ_k defined in Section 2.1. Lemma 2.23 carries over to this function, in particular $\tilde{P}_\mu dg(\mu, k) \wedge d\lambda$ is a regular 2-form on the complex Bloch variety and all eigenfunctions χ , which correspond to some element (k, λ) of the complex Bloch variety belong to the range of the projection $\psi_k \check{\tilde{P}}_\mu(k, \lambda)$, which is analogous to the projection $\check{\tilde{P}}_{\tilde{\gamma}}(k, \lambda)$ defined in (v) of Lemma 2.23. It might happen, however, that the restriction of $\tilde{P}_\mu$ to the *complex Fermi curve* is not well defined, because the denominator might vanish identically on a connected component of the *complex Fermi curve*. In these cases, the second part of the statement (iv) of Lemma 2.23 cannot be generalized to this projection. But Theorem 2.12 and Corollary 2.13 imply that this can only happen, if μ is proportional to $(1, \pm\sqrt{-1})$ and if the *complex Fermi curve* is the same as the *complex Fermi curve* of the zero potential.

The restriction of this projection to the *complex Fermi curve* is again the projection of some operator-valued holomorphic function, whose complex Bloch variety is equal to the *complex Fermi curve* $\mathcal{F}(V, W)$. Let $\hat{\mu}$ and $\check{\mu}$ be an arbitrary bases of $\mathbb{C}^2$ and $\hat{\nu}$ and $\check{\nu}$ the dual basis:

$$g(\hat{\mu}, \hat{\nu}) = 1 = g(\check{\mu}, \check{\nu}) \text{ and } g(\hat{\mu}, \check{\nu}) = 0 = g(\check{\mu}, \hat{\nu}).$$

In analogy to the notation introduced in the context of the *Fundamental domain* 2.1 we parameterize the momentum space by $\hat{p} = g(\hat{\mu}, k)$ and $\check{p} = g(\check{\mu}, k)$, which is equivalent to $k = \hat{p}\hat{\nu} + \check{p}\check{\nu}$. If we assume that $g(\check{\nu}, \check{\nu})$ is not equal to zero, then the matrix $\begin{pmatrix} 0 & \check{\nu}_2 + \sqrt{-1}\check{\nu}_1 \\ \check{\nu}_2 - \sqrt{-1}\check{\nu}_1 & 0 \end{pmatrix}$ is invertible, and the inverse equals $\begin{pmatrix} 0 & 1/(\check{\nu}_2 - \sqrt{-1}\check{\nu}_1) \\ 1/(\check{\nu}_2 + \sqrt{-1}\check{\nu}_1) & 0 \end{pmatrix}$. The function $\psi = \psi_{\hat{p}\hat{\nu} + \check{p}\check{\nu}}\tilde{\psi}$ is an eigenfunction of the Dirac operator $D(V, W, \hat{p}\hat{\nu} + \check{p}\check{\nu})$ with eigenvalue zero if and only if

the function $\tilde{\psi}$ is an eigenfunction of the operator

$$\begin{aligned}\tilde{D}_{\hat{\mu}, \check{\mu}}(V, W, \hat{p}) &= \begin{pmatrix} 0 & \check{\nu}_2 + \sqrt{-1}\check{\nu}_1 \\ \check{\nu}_2 - \sqrt{-1}\check{\nu}_1 & 0 \end{pmatrix}^{-1} \left(D(V, W, 0) + \hat{p}\pi \begin{pmatrix} 0 & \hat{\nu}_2 + \sqrt{-1}\hat{\nu}_1 \\ \hat{\nu}_2 - \sqrt{-1}\hat{\nu}_1 & 0 \end{pmatrix} \right) \\ &= \begin{pmatrix} \frac{\hat{p}\pi(\check{\nu}_2 - \sqrt{-1}\check{\nu}_1) - \check{\partial}_0}{\check{\nu}_2 - \sqrt{-1}\check{\nu}_1} & \frac{W}{\check{\nu}_2 - \sqrt{-1}\check{\nu}_1} \\ \frac{V}{\check{\nu}_2 + \sqrt{-1}\check{\nu}_1} & \frac{\hat{p}\pi(\hat{\nu}_2 + \sqrt{-1}\hat{\nu}_1) + \partial_0}{\hat{\nu}_2 + \sqrt{-1}\hat{\nu}_1} \end{pmatrix}\end{aligned}$$

with eigenvalue $\check{p}\pi$. Thus the *complex Fermi curve* $\mathcal{F}(V, W)$ is equal to the set of all

$$\left\{ k = \hat{p}\hat{\nu} + \check{p}\check{\nu} \in \mathbb{C}^2 \mid \check{p}\pi \text{ is an eigenvalue of } \tilde{D}_{\hat{\mu}, \check{\mu}}(V, W, \hat{p}) \right\},$$

and due to Lemma 2.25 for all $k = \hat{p}\hat{\nu} + \check{p}\check{\nu} \in \mathcal{F}(V, W)$ the projection $\check{\mathbf{P}}_{\hat{\mu}}(k)$ is the spectral projection of $\tilde{D}_{\hat{\mu}, \check{\mu}}(V, W, \hat{p})$ onto the generalized eigenspace associated with eigenvalue $\check{p}\pi$, which is equal to the residue of the resolvent of $\tilde{D}_{\hat{\mu}, \check{\mu}}(V, W, \hat{p})$ at the eigenvalue $\check{p}\pi$ (see e. g. [R-S-IV, Appendix to XII.1]).

2.5 Complex Fermi curves of finite genus

If there exists some finite subset of Λ^* , such that for all κ in the complement of this finite subset the normalizations of the handle with index κ described in Theorem 2.12 decomposes into two connected components, one of which is a disc excluded from $\mathcal{V}_{\varepsilon, \delta}^+$ around the point k_{κ}^+ and the other is a disc excluded from $\mathcal{V}_{\varepsilon, \delta}^-$ around the point k_{κ}^- , then the normalization of $\mathcal{F}(V, W)/\Lambda^*$ is biholomorphic to a compact Riemann surface with two points at infinity removed. In fact, in this case the normalization of $\mathcal{F}(V, W)/\Lambda^*$ contains two open subsets, which are biholomorphic to the open subsets

$$\begin{aligned}&\{k \in \mathbb{C}^2 \mid k_1 - \sqrt{-1}k_2 = 0 \text{ and } \|k\| > 1/\delta\} \text{ and} \\ &\{k \in \mathbb{C}^2 \mid k_1 + \sqrt{-1}k_2 = 0 \text{ and } \|k\| > 1/\delta\}\end{aligned}$$

of the normalization of $\mathcal{F}(0, 0)$, with some $\varepsilon > 0$. Due to Theorem 2.12 the complement of these two open subsets of the normalization of $\mathcal{F}(V, W)/\Lambda^*$ is compact. Hence we may compactify the normalization of $\mathcal{F}(U, U)/\Lambda^*$ by adding to the first subset the point $\infty^- = \lim_{t \rightarrow \infty} (t, -\sqrt{-1}t)$ and to the second subset the point $\infty^+ = \lim_{t \rightarrow \infty} (t, \sqrt{-1}t)$.

If we normalize the eigenfunctions ψ of the Dirac operator by some linear condition, then the eigenfunction is a meromorphic function on $\mathcal{F}(V, W)/\Lambda^*$. The following normalization of the eigenfunction $\psi(x) = \begin{pmatrix} \psi_1(x) \\ \psi_2(x) \end{pmatrix}$ and the eigenfunction of the transposed Dirac operator $\phi(x) = \begin{pmatrix} \phi_1(x) \\ \phi_2(x) \end{pmatrix}$ is convenient:

$$\psi_1(x=0) + \psi_2(x=0) = \sqrt{2} \text{ and } \phi_1(x=0) - \phi_2(x=0) = \sqrt{2},$$

respectively. If h_0 denotes the matrix $h_0 = 1/\sqrt{2} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$, then it may be written in the form $(h_0\psi(x=0))_1 = 1$ and $(Jh_0\phi(x=0))_1 = 1$. Although these eigenfunctions are always

meromorphic functions on the normalization of $\mathcal{F}(V, W)/\Lambda^*$, it might happen that this normalization is biholomorphic to a compact Riemann surface with two points removed, but the eigenfunctions have an infinite number of singularities on the normalization of $\mathcal{F}(V, W)/\Lambda^*$ [Sch]. In this case they are not meromorphic functions on the compact Riemann surface with two essential singularities at the two points at infinity. Consequently we call a pair of potentials (V, W) to be of finite type, if they obey the following conditions:

Finite type potentials (i) The normalization of $\mathcal{F}(V, W)/\Lambda^*$ can be compactified to a compact Riemann surface by adding the two points ∞^- and ∞^+ at infinity.

Finite type potentials (ii) The pullback of the normalized eigenfunctions under the normalization map have only a finite number of poles on the normalization of $\mathcal{F}(V, W)/\Lambda^*$.

The entries of the normalized eigenfunctions generate a coherent subsheaf of the sheaf of meromorphic functions. Due to an improved version of statement (iv) in Lemma 2.23 (compare with [Sch, Chapter 9]) the number of poles of the normalized eigenfunction is equal to the arithmetic genus of the *Structure sheaf* 2.24 plus one (i. e. the arithmetic genus of the modified *Structure sheaf* 2.24, where the two infinities are identified to an ordinary double point). Therefore, condition *Finite type potentials* (ii) is equivalent to the condition that the *arithmetic genus* of the *Structure sheaf* 2.24 is finite. Moreover, condition *Finite type potentials* (i) is equivalent to the condition that the *geometric genus* is finite. Therefore, condition *Finite type potentials* (ii) implies condition *Finite type potentials* (i). We should remark that these eigenfunctions may be characterized by some abstract properties and are called *Baker–Akhiezer functions* (compare with [D-K-N, Chapter 2 §2.] and conditions *Baker–Akhiezer function* (i)–(ii) in the proof of Lemma 2.30). Obviously these functions uniquely determine the corresponding potentials, which therefore have to be analytic. Finally, all sheaves of this form may be characterized (compare with [Sch]).

Lemma 2.26. *If a pair of potentials of the form $(U, \bar{U})$ obeys the condition Finite type potentials (i), then it satisfies also the condition Finite type potentials (ii).*

Proof. For small $\varepsilon > 0$ the open subsets $\mathcal{V}_{\varepsilon, \delta}^\pm$ of $\mathcal{F}(0, 0)$ are one-sheeted coverings over some open subset of $k_1 \in \mathbb{C}$. Thus, due to the proof of Theorem 2.12, the same is true for arbitrary potentials V and W . Furthermore, with the exception of finitely many $\kappa \in \Lambda^*$ all handles with index κ are two-sheeted coverings over $\hat{p} \in \mathbb{C}$. Hence the only possible singularities of these handles are double points. If the normalization of $\mathcal{F}(U, \bar{U})/\Lambda^*$ is a compact Riemann surface with two points removed, then with the exception of finitely many $\kappa \in \Lambda^*$ the normalization of the handles with index κ have two connected components. Since the involution η permutes the points $k_\kappa^\pm$, this involution also permutes the two connected components of the normalization of these handles. This implies that with the exception of finitely many $\kappa \in \Lambda^*$ the only possible singularities of the handle with index κ are double points, which are invariant under η . Now Corollary 2.17 and condition (iii) of Lemma 2.23 imply that the projection valued function $P_{\hat{\gamma}}$ has no poles on these handles. Thus due to the Maximum principle (see e.g. [Ah, Chapter 4. Theorem 12.]), the estimates of the eigenfunctions from Theorem 2.12 on the open sets $\mathcal{V}_{\varepsilon, \delta}^\pm$ extend to these handles. This proves that the normalized eigenfunctions have only a finite number of poles on the normalization of $\mathcal{F}(U, \bar{U})/\Lambda^*$. **q.e.d.**

2.5.1 The inverse problem

As we have seen, in some cases we may associate to potentials a compact Riemann surface, and moreover we found several properties of these Riemann surface. Let us now try to invert this procedure and to associate to some Riemann surface with some specified properties a pair of potentials, such that the compactification of the normalization of the corresponding *complex Fermi curve* is equal to the given Riemann surface. First we introduce the data of tuples $(Y, \infty^-, \infty^+, k)$. Here Y is a compact pure one-dimensional complex space with two marked non-singular points $\infty^\pm$ with the property that each connected component of the subspace of nonsingular points contains at least one of these points, and k is a multi-valued meromorphic function from Y into $\mathbb{C}^2$, which is holomorphic on $Y \setminus \{\infty^-, \infty^+\}$. The complex space $Y \setminus \{\infty^-, \infty^+\}$ will be a candidate for the one-sheeted covering of a *complex Fermi curve*, whose structure sheaf is defined in *Structure sheaf* 2.24. In the sequel it mostly can be chosen to be a compact smooth Riemann surface. Furthermore, we assume the following conditions:

Quasi-momenta (i) The function k has poles of first-order at the two marked points and for some branch of this function the $\mathbb{C}$ -valued functions $k_1 - \sqrt{-1}k_2$ and $k_1 + \sqrt{-1}k_2$ vanish at ∞^- and ∞^+ , respectively.

Quasi-momenta (ii) The difference of two arbitrary branches of the function k is some element of Λ^* . Conversely, for all $\kappa \in \Lambda^*$ and all branches of k , $k + \kappa$ is some other branch of k .

Here we do not assume that we may pass from one branch to all other branches by continuation along closed cycles of Y (compare with Remark 4.17).

For all such data $(Y, \infty^-, \infty^+, k)$ the image of $Y \setminus \{\infty^-, \infty^+\}$ under the mapping k is a pure one-dimensional complex subvariety of $\mathbb{C}^2$, which is invariant under translations by elements of Λ^* . As shown in the following lemma these images are *complex Fermi curves* and consequently denoted by $\mathcal{F}(Y, \infty^-, \infty^+, k)$ or just $\mathcal{F}$. The *complex Fermi curves* we are interested in are endowed with the anti-holomorphic involutions ρ and η and the holomorphic involution σ . In the sequel we therefore impose sometimes the additional conditions:

Quasi-momenta (iii) The complex space Y is endowed with an anti-holomorphic involution η without fixed points. Moreover, any branch of the transformed function η^*k is equal to some branch of the function $-\bar{k}$.

Quasi-momenta (iv) The complex space Y is endowed with an holomorphic involution σ . Moreover, any branch of the transformed function σ^*k is equal to some branch of the function $-k$.

Lemma 2.27. *Let Y be a smooth connected compact smooth Riemann surface with two marked points ∞^- and ∞^+ and let k be a multi-valued meromorphic function, which is holomorphic on $Y \setminus \{\infty^-, \infty^+\}$. If $(Y, \infty^-, \infty^+, k)$ satisfies the conditions Quasi-momenta (i)–(iii), then there exists an analytic complex potential U , whose complex Fermi curve is equal*

to $\mathcal{F}(Y, \infty^-, \infty^+, k)$. Moreover, the pullback of the eigenfunction under the normalization map takes at all pairwise different elements of $Y \setminus \{\infty^-, \infty^+\}$ linearly independent values. Finally, for any choice of finitely many points $y_1, \dots, y_l$ of $Y \setminus \{\infty^-, \infty^+\}$ there exists one of these potentials, such that the values of the pullback of the corresponding eigenfunction at these points $y_1, \dots, y_l$ has no zeroes considered as a function from Δ to $\mathbb{C}^2$.

Due to [G-S-1, Lemma 2.3] the assumption of connectedness of Y is necessary. In fact, if Y has two connected components, the forms $dk_1 \pm \sqrt{-1}dk_2$ are holomorphic on the component containing $\infty^\pm$, respectively. This implies that the intersection form of those two cycles does not vanish, whose intersection forms with any other cycle is equal to the integral of $\hat{p}$ and $\check{p}$ along this cycle.

Proof. We use the methods of inverse spectral theory of Lax operators (see e. g. [Kr-1, D-K-N]). For this purpose it is convenient to introduce the compact complex space $\tilde{Y}$, which is obtained from Y by identifying the two marked points $\infty^\pm$ to an ordinary double point. Furthermore, we shall introduce the real two-dimensional subgroup of the Picard group corresponding to the translations by $\mathbb{R}^2/\Lambda$. For all $x \in \mathbb{R}^2$, let $\tilde{\mathcal{L}}(x)$ denote the locally free sheaf of rank one on $\tilde{Y}$, which is defined by a co-cycle with respect to the covering $\tilde{Y} = \tilde{Y} \setminus \{\infty^-, \infty^+\} \cup$ some small neighbourhood of the double point (∞^-, ∞^+) . On the intersection of the two open sets the transition function of this co-cycle is equal to

$$\exp(2\pi\sqrt{-1}g(x, k)) = \exp(2\pi\sqrt{-1}(g(x, \hat{\kappa})\hat{p} + g(x, \check{\kappa})\check{p})) = \exp(2\pi\sqrt{-1}(\hat{q}\hat{p} + \check{q}\check{p}))$$

for any $k \in \mathbb{C}^2$. If x belongs to the lattice Λ , then this transition function extends to a global non-vanishing holomorphic function on $\tilde{Y} \setminus \{\infty^-, \infty^+\}$, and the corresponding sheaf is equal to the structure sheaf of $\tilde{Y}$. Due to the general construction (see e. g. [D-K-N, Sch]) for any divisor $\tilde{D}$ on $\tilde{Y}$, which has the following properties, there is associated a unique *Baker–Akhiezer function*.

Divisor (i) $\tilde{D}$ is an integral divisor, whose support is contained in $\tilde{Y} \setminus \{\infty^-, \infty^+\}$. Moreover, the degree of $\tilde{D}$ is equal to the genus of Y plus one (i. e. the arithmetic genus of $\tilde{Y}$).

Divisor (ii) The difference $\eta(\tilde{D}) - \tilde{D}$ is the principal divisor of a function f , which takes the values ± 1 at the marked points $\infty^\pm$. In particular, this function obeys $f\eta^*f = -1$.

Divisor (iii) For all $x \in \mathbb{R}^2$ the sheaf $\tilde{\mathcal{L}}(x) \otimes \mathcal{O}_{\tilde{D}}$ on $\tilde{Y}$ has no non-trivial global section, which vanishes at the double point (∞^-, ∞^+) .

For given $\tilde{D}$ let D denote the corresponding divisors on Y . With the help of the Riemann–Roch theorem [Fo, Theorem 16.9] it is quite easy to see that for any divisor D on Y , which has the following properties *Divisor* (i')–(iii'), there exists a family $\tilde{D}_t$ parameterized by $t \in S^1$ of divisors on $\tilde{Y}$, fulfilling conditions *Divisor* (i)–(ii), whose corresponding divisors D_t are all isomorphic to D . The degree of freedom of this family corresponds to the isospectral transformations described at the end of Section 2.3.

Divisor (i') The degree of D is equal to the genus of Y plus one.

Divisor (ii') The difference $\eta(D) - D$ is a principal divisor of a function f , which obeys $f\eta^*\bar{f} < 0$.

Divisor (iii') The sheaf $\mathcal{O}_D$ has no global non-trivial section, which vanishes at the two marked points ∞^- and ∞^+ .

Remark 2.28. *Complex line bundles, whose divisors obey condition (ii) are called quaternionic, since the space of sections is a quaternionic space [Hi, p. 667]. In order to distinguish this notion of quaternionic line bundles from the notion of quaternionic line bundles occurring in ‘quaternionic function theory’, we shall emphasize that the first notion describes a subclass of complex line bundles. Consequently we shall speak of quaternionic complex line bundles.*

For all $x \in \mathbb{R}^2$ let $\mathcal{L}(x)$ denotes the analogous locally free sheaf of rank one on Y . Obviously these sheaves correspond to real line bundles of degree zero. In particular, the tensor product of these line bundles with any quaternionic complex line bundle gives again quaternionic complex line bundles. Hence we have to show that there exists a divisor D , whose degree is equal to the genus of Y plus one, such that the sheaves $\mathcal{L}(x) \otimes \mathcal{O}_D$ have no non-trivial global section, which vanishes at the two marked points ∞^- and ∞^+ . Obviously all these sheaves correspond to quaternionic complex line bundles. Hence the dimension of the space of global sections is divisible by two. Due to [G-H, Proposition 3.3] the real part of the component of the Picard group of Y corresponding to degree $g \pm 1$ has a component, which contains only quaternionic complex line bundles. Due to Marten’s theorem [A-C-G-H, Chapter IV (5.1) Theorem] the subvariety of the Picard group of divisors of degree $g - 1$, whose space of sections is at least two-dimensional, has co-dimension of at least three. In Marten’s theorem it is assumed that the genus is not smaller than three. If the genus is smaller than three, this statement is trivial, because in these cases no line bundle, whose degree is equal to the genus minus one, can have a two-dimensional space of global sections. Therefore, the subvariety of the real part of the Picard group of Y corresponding to quaternionic complex line bundles of degree $g - 1$, which have non-trivial global sections, has co-dimension larger than two. We conclude that there exists a divisor $\tilde{D}$ fulfilling conditions *Divisor* (i)–(iii), and consequently a corresponding *Baker–Akhiezer function* and a corresponding pair of potentials of the form $(U, \bar{U})$ whose *complex Fermi curve* is equal to $\mathcal{F}(Y, \infty^-, \infty^+, k)$.

If the eigenfunctions corresponding to two different elements $y \neq y'$ of $Y \setminus \{\infty^-, \infty^+\}$ are linearly dependent, then the function k has to take the same values mod Λ^* at these elements. Due to Lemma 2.25 this implies that the projection $P_{\hat{\gamma}} d\hat{p}$ is not an operator-valued holomorphic form on $Y \setminus \{\infty^-, \infty^+\}$. More precisely, the local sum over all sheets of Y considered as a covering over $\hat{p} \in \mathbb{C}$, which contain y or y' (compare with Remark 2.19), respectively, will then not be a holomorphic function with respect to $\hat{p}$. On the other hand the corresponding *Structure sheaf* 2.24 is equal to the normalization sheaves. Therefore, due to an improved version of statement (iv) in Lemma 2.23 (compare with [Sch, Chapter 9]), this projection valued form $P_{\hat{\gamma}} d\hat{p}$ is holomorphic, and the values of the *Baker–Akhiezer function* at different points are linearly independent.

Finally, we remark that due to the construction of the *Baker–Akhiezer function*, the eigenfunction corresponding to some element $y \in Y \setminus \{\infty^-, \infty^+\}$ considered as a function from Δ to $\mathbb{C}^2$ has a zero at $x \in \Delta$, if and only if

- (i) the element y belongs to the support of $\tilde{D}(x)$, and
- (ii) the unique global section f of $\mathcal{O}_{\tilde{D}(x)}$ on Y , which takes the values ± 1 at the two marked points $\infty^\pm$, is also a section of $\mathcal{O}_{\tilde{D}(x)-y}$.

Here $\tilde{D}(x)$ denotes the unique integral divisor of $\tilde{Y}$, whose sheaf $\mathcal{O}_{\tilde{D}(x)}$ is isomorphic to $\tilde{\mathcal{L}}(x) \otimes \mathcal{O}_{\tilde{D}}$. Due to condition *Divisor* (ii) this implies that $\eta(y)$ also belongs to the support of $\tilde{D}(x)$, and that f is even a global section of $\mathcal{O}_{\tilde{D}(x)-y-\eta(y)}$. In particular, the corresponding divisor $D(x)-y-\eta(y)$ of Y is an integral divisor of degree $g-1$, whose space of global sections is at least two-dimensional. Again Marten’s theorem [A-C-G-H, Chapter IV (5.1) Theorem] shows that for generic y this is not the case for all $x \in \mathbb{R}^2$. q.e.d.

2.5.2 The Bäcklund transformations

The one-sheeted coverings of the *complex Fermi curve* of these potentials, whose structure sheaf is defined in *Structure sheaf* 2.24, are equal to the normalization of $Y \setminus \{\infty^-, \infty^+\}$. This construction can be generalized to all one-sheeted coverings of $\mathcal{F}(Y, \infty^-, \infty^+, k)/\Lambda^*$ of finite arithmetic genus, on which the involution η does not have any fixed points (i. e. all double points of the form $(y, \eta(y))$ of the *complex Fermi curve* have to be removed). Moreover, the *Bäcklund transformation* provides a tool for transforming a potential corresponding to a one-sheeted covering of the *complex Fermi curve* into a potential corresponding to another one-sheeted covering, such that the former is a one-sheeted covering of the latter (compare with [E-K], [M-S, Chapter 6.] and [L-McL, Section 4]). We shall now derive the formulas of these *Bäcklund transformations* in the present context. First we remark that for all pairs (y, y') of the *complex Fermi curve* the transformation $D \mapsto D + \infty^- + \infty^+ - y - y'$ preserves the degree of the divisor and therefore should induce an isospectral transformation of the corresponding potentials, in case they exist. Moreover, due to the properties of the *Baker–Akhiezer function*, the n -th partial derivatives of the eigenfunctions yield cross sections of the sheaves $\mathcal{O}_{D(x)+n\infty^-+n\infty^+}$. Therefore, we may calculate the *Baker–Akhiezer functions* corresponding to the divisors $D(x) + \infty^- + \infty^+ - y - y'$ in terms of the *Baker–Akhiezer function* corresponding to the divisors $D(x)$. Since we are only interested in pairs of potentials of the form $(U, \bar{U})$ the pair (y, y') has to be invariant under η . In fact, the transformation $D \mapsto D + \infty^- + \infty^+ - y - \eta(y)$ preserves the divisors corresponding to quaternionic complex line bundles. Let ψ be the *Baker–Akhiezer function* of the pair of complex potentials $(U, \bar{U})$ and χ and $J\bar{\chi}$ the values of ψ at the pair $(y, \eta(y))$ of points of the corresponding *complex Fermi curve*. In the following considerations χ may be multiplied with some complex number without changing the transformation. The function

$$\psi' = \begin{pmatrix} \partial + a & b \\ c & \bar{\partial} + d \end{pmatrix} \psi$$

with periodic functions a, b, c, d on $\mathbb{R}^2/\Lambda$ has the correct asymptotic behaviour of the *Baker–Akhiezer function* corresponding to $D(x) + \infty^- + \infty^+ - y - \eta(y)$. If we impose the condition, that ψ' vanishes at y and $\eta(y)$, then these functions are given by

$$\begin{aligned} a &= -\frac{(\partial\chi_1)\bar{\chi}_1 + \chi_2(\partial\bar{\chi}_2)}{\chi_1\bar{\chi}_1 + \chi_2\bar{\chi}_2} & b &= \frac{\chi_1(\partial\bar{\chi}_2) - (\partial\chi_1)\bar{\chi}_2}{\chi_1\bar{\chi}_1 + \chi_2\bar{\chi}_2} \\ c &= -\bar{b} = \frac{\chi_2(\bar{\partial}\bar{\chi}_1) - (\bar{\partial}\chi_2)\bar{\chi}_1}{\chi_1\bar{\chi}_1 + \chi_2\bar{\chi}_2} & d &= \bar{a} = -\frac{\chi_1(\bar{\partial}\bar{\chi}_1) + (\bar{\partial}\chi_2)\bar{\chi}_2}{\chi_1\bar{\chi}_1 + \chi_2\bar{\chi}_2}. \end{aligned}$$

In order to verify that the corresponding *Baker–Akhiezer function* ψ' is indeed an eigenfunction of some Dirac operator we calculate the derivatives

$$\begin{aligned} \bar{\partial}a &= U\bar{U} - b\bar{b} & \bar{\partial}b &= -\partial\bar{U} - \bar{U}a + b\bar{a} \\ \partial c &= -\partial\bar{b} = \bar{\partial}U + U\bar{a} - a\bar{b} & \partial d &= \partial\bar{a} = U\bar{U} - b\bar{b}. \end{aligned}$$

Now a direct calculation shows that ψ' is indeed an eigenfunction corresponding to the pair of potentials $(U', \bar{U}') = (\bar{b}, b)$, if χ has no zero considered as a function from Δ to $\mathbb{C}^2$. In this case the *complex Fermi curve* $\mathcal{F}(U', \bar{U}')$ coincides with $\mathcal{F}(U, \bar{U})$. We should remark that the condition, that the value χ of the original *Baker–Akhiezer function* at y (or $\eta(y)$) has no zero considered as a function from Δ to $\mathbb{C}^2$, fits perfectly with Lemma 2.27. Obviously these formulas extend to the situation, where χ is any element of the kernel of the Dirac operator $D(U, \bar{U}, k)$ with $k \in \mathcal{F}(U, \bar{U})$, which has no zero considered as a function from Δ to $\mathbb{C}^2$. In case that k is a singularity of the *complex Fermi curve* there might exist several values of the pullback of the *Baker–Akhiezer function* to the normalization and more generally the eigenspace might have dimension larger than one. However, only those elements of the range corresponding spectral projection $\check{P}([k])$ may be used in order to obtain a periodic potential, which are proper eigenfunctions and have no zero considered as a function from Δ to $\mathbb{C}^2$.

In general these formulas yield for two given (not necessarily quasi-periodic) eigenfunctions χ and ψ of the Dirac operator $(\begin{smallmatrix} U & \partial \\ -\bar{\partial} & \bar{U} \end{smallmatrix})$ on some domain of $\mathbb{C}$ another eigenfunction ψ' corresponding to another Dirac operator $(\begin{smallmatrix} U' & \partial \\ -\bar{\partial} & \bar{U}' \end{smallmatrix})$. We would have obtained more general formulas, if we had started with two elements of the complex Bloch variety (y, y') , which are not invariant under η . In this context they are called Bäcklund transformations [E-K].

We shall derive also the formulas for the inverse transformation corresponding to the transformation $D \mapsto D + y + \eta(y) - \infty^- - \infty^+$. For this purpose we remark that the arguments of the proof of Lemma 2.23 (iv) shows that the divisor D of the *Baker–Akhiezer function* and the divisor D^t of the transposed *Baker–Akhiezer function* obey the relation $\mathcal{O}_D \otimes \mathcal{O}_{D^t} \simeq \mathcal{O}_{K+2\infty^-+2\infty^+}$, where $\mathcal{O}_K$ is the dualizing sheaf of the one-sheeted covering described in *Structure sheaf* 2.24 (for smooth one sheeted coverings K is the canonical divisor). More precisely, there exists a more restrictive version of this relation, which makes use of the corresponding one-sheeted coverings of the *complex Fermi curve*, whose infinities are identified to an ordinary double point (compare with [Sch]). Therefore, the transformation $D \mapsto D + y + \eta(y) - \infty^- - \infty^+$ corresponds to the transformation $D^t \mapsto D^t + \infty^- + \infty^+ - y - \eta(y)$, i. e. to the transformation described above applied to the transposed operators. However, we may use Lemma 2.22 in order to obtain a formula for the *Baker–Akhiezer functions* instead

of the transposed *Baker–Akhiezer functions*. Let ψ denote the *Baker–Akhiezer function* of the pair of potentials $(U, \bar{U})$, and let ϕ and $\bar{\phi}$ denote the values of the transposed *Baker–Akhiezer function* at the two elements k' and $-\bar{k}'$ of the *complex Fermi curve*. Again in the following formulas ϕ may be multiplied with some complex number without changing the transformation. Due to Lemma 2.22 the two functions

$$\frac{\int_x^{x+\hat{\gamma}} \phi_2 \psi_1 dz + \phi_1 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k - k')) - 1} \quad \text{and} \quad \frac{\int_x^{x+\hat{\gamma}} \bar{\phi}_1 \psi_1 dz - \bar{\phi}_2 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k + \bar{k}')) - 1}$$

define for all $x \in \mathbb{R}^2$ meromorphic functions on the *complex Fermi curve* with first-order poles at the elements k' and $-\bar{k}'$, respectively. Moreover, due to the normalization factor, in which k denotes the corresponding function on the *complex Fermi curve*, these functions are cross sections of the bundle on $\mathbb{R}^2/\Lambda$ corresponding to $[k - k']$ and $[k + \bar{k}']$, respectively. Furthermore, due to the normalization factor these functions do not depend on the period $\hat{\gamma}$. Their asymptotic behaviour near $\infty^\pm$ may be calculated as

$$\begin{aligned} \frac{\int_x^{x+\hat{\gamma}} \phi_2 \psi_1 dz + \phi_1 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k - k')) - 1} &= \begin{cases} \frac{(\hat{\gamma}_1 - \sqrt{-1}\hat{\gamma}_2)\phi_1(x) \exp(2\pi\sqrt{-1}g(x, k))}{2\pi\sqrt{-1}g(\hat{\gamma}, k)} (1 + \mathbf{O}(1/g(\hat{\gamma}, k))) & \text{at } \infty^- \\ \frac{(\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2)\phi_2(x) \exp(2\pi\sqrt{-1}g(x, k))}{2\pi\sqrt{-1}g(\hat{\gamma}, k)} (1 + \mathbf{O}(1/g(\hat{\gamma}, k))) & \text{at } \infty^+. \end{cases} \\ \frac{\int_x^{x+\hat{\gamma}} \bar{\phi}_1 \psi_1 dz - \bar{\phi}_2 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k + \bar{k}')) - 1} &= \begin{cases} \frac{-(\hat{\gamma}_1 - \sqrt{-1}\hat{\gamma}_2)\bar{\phi}_2(x) \exp(2\pi\sqrt{-1}g(x, k))}{2\pi\sqrt{-1}g(\hat{\gamma}, k)} (1 + \mathbf{O}(1/g(\hat{\gamma}, k))) & \text{at } \infty^- \\ \frac{(\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2)\bar{\phi}_1(x) \exp(2\pi\sqrt{-1}g(x, k))}{2\pi\sqrt{-1}g(\hat{\gamma}, k)} (1 + \mathbf{O}(1/g(\hat{\gamma}, k))) & \text{at } \infty^+. \end{cases} \end{aligned}$$

We conclude that

$$\psi' = \frac{\begin{pmatrix} \bar{\phi}_2 \\ \bar{\phi}_1 \end{pmatrix}}{\phi_1 \bar{\phi}_1 + \phi_2 \bar{\phi}_2} \frac{\int_x^{x+\hat{\gamma}} \phi_2 \psi_1 dz + \phi_1 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k - k')) - 1} + \frac{\begin{pmatrix} \phi_1 \\ -\phi_2 \end{pmatrix}}{\phi_1 \bar{\phi}_1 + \phi_2 \bar{\phi}_2} \frac{\int_x^{x+\hat{\gamma}} \bar{\phi}_1 \psi_1 dz - \bar{\phi}_2 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k + \bar{k}')) - 1}$$

has the properties of the *Baker–Akhiezer function* corresponding to the divisor $D + y + \eta(y) - \infty^- - \infty^+$. The derivatives of these functions are given by

$$\begin{aligned} \frac{\partial \int_x^{x+\hat{\gamma}} \phi_2 \psi_1 dz + \phi_1 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k - k')) - 1} &= \phi_2(x) \psi_1(x), & \frac{\partial \int_x^{x+\hat{\gamma}} \bar{\phi}_1 \psi_1 dz - \bar{\phi}_2 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k + \bar{k}')) - 1} &= \bar{\phi}_1(x) \psi_1(x), \\ \frac{\bar{\partial} \int_x^{x+\hat{\gamma}} \phi_2 \psi_1 dz + \phi_1 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k - k')) - 1} &= \phi_1(x) \psi_2(x), & \frac{\bar{\partial} \int_x^{x+\hat{\gamma}} \bar{\phi}_1 \psi_1 dz - \bar{\phi}_2 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k + \bar{k}')) - 1} &= -\bar{\phi}_2(x) \psi_2(x). \end{aligned}$$

Furthermore, the values of ψ' at the points k' and $-\bar{k}'$ satisfy the equations

$$\begin{pmatrix} U' & \partial \\ -\bar{\partial} & \bar{U}' \end{pmatrix} \frac{\begin{pmatrix} \bar{\phi}_2 \\ \bar{\phi}_1 \end{pmatrix}}{\phi_1 \bar{\phi}_1 + \phi_2 \bar{\phi}_2} = 0, \quad \begin{pmatrix} U' & \partial \\ -\bar{\partial} & \bar{U}' \end{pmatrix} \frac{\begin{pmatrix} \phi_1 \\ -\phi_2 \end{pmatrix}}{\phi_1 \bar{\phi}_1 + \phi_2 \bar{\phi}_2} = 0,$$

with the transformed potential

$$U' = \frac{(\partial\phi_2)\bar{\phi}_1 - \phi_2(\partial\bar{\phi}_1)}{\phi_1\bar{\phi}_1 + \phi_2\bar{\phi}_2}.$$

All these equations imply that ψ' is indeed the *Baker–Akhiezer function* of the pair of potentials $(U', \bar{U}')$, if ϕ considered as a function from Δ to $\mathbb{C}^2$ has no zero. Moreover, if in addition χ is an element of the kernel of $D(U, \bar{U}, k')$, such that the integral $\int_{x'}^x \phi_2 \chi_1 dz + \phi_1 \chi_2 d\bar{z}$ yields a periodic function with respect to x , then for all $x' \in \mathbb{R}^2$ the function

$$\chi' = \frac{\begin{pmatrix} \bar{\phi}_2 \\ \bar{\phi}_1 \end{pmatrix}}{\phi_1\bar{\phi}_1 + \phi_2\bar{\phi}_2} \int_{x'}^x \phi_2 \chi_1 dz + \phi_1 \chi_2 d\bar{z} + \frac{\begin{pmatrix} \phi_1 \\ -\phi_2 \end{pmatrix}}{\phi_1\bar{\phi}_1 + \phi_2\bar{\phi}_2} \frac{\int_x^{x+\hat{\gamma}} \bar{\phi}_1 \chi_1 dz - \bar{\phi}_2 \chi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k' + \bar{k}')) - 1}$$

belongs to the kernel of the transformed Dirac operator $D(U', \bar{U}', k')$. Due to Lemma 2.22 this condition on χ is equivalent to the condition $\langle \phi_2, \chi_1 \rangle = 0$ and $\langle \phi_1, \chi_2 \rangle = 0$.

A direct calculation shows that for any element y of the normalization the two *Bäcklund transformations* corresponding to the values of the pullback of the (transposed) *Baker–Akhiezer function* to the normalization are inverse to each other, in agreement with the corresponding transformations of the divisors: $D \mapsto D + \infty^- + \infty^+ - y - \eta(y)$ and $D \mapsto D + y + \eta(y) - \infty^- - \infty^+$. We collect our calculations and observations in the following

Lemma 2.29. *Let ψ be the Baker–Akhiezer function of the pair of potentials $(U, \bar{U})$. If χ belongs to the kernel of $D(U, \bar{U}, k)$ and has no zero considered as a function from Δ to $\mathbb{C}^2$, then the function*

$$\psi' = \left(\frac{1}{\chi_1 \bar{\chi}_1 + \chi_2 \bar{\chi}_2} \begin{pmatrix} -(\partial\chi_1)\bar{\chi}_1 + \chi_2(\partial\bar{\chi}_2) & \chi_1(\partial\bar{\chi}_2) - (\partial\chi_1)\bar{\chi}_2 \\ \chi_2(\bar{\partial}\bar{\chi}_1) - (\bar{\partial}\chi_2)\bar{\chi}_1 & -\chi_1(\bar{\partial}\bar{\chi}_1) + (\bar{\partial}\chi_2)\bar{\chi}_2 \end{pmatrix} + \begin{pmatrix} \partial & 0 \\ 0 & \bar{\partial} \end{pmatrix} \right) \psi$$

is the Baker–Akhiezer function of the pair of potentials $(U', \bar{U}')$ with

$$U' = \frac{(\bar{\partial}\chi_2)\bar{\chi}_1 - \chi_2(\bar{\partial}\bar{\chi}_1)}{\chi_1\bar{\chi}_1 + \chi_2\bar{\chi}_2},$$

and the complex Fermi curves $\mathcal{F}(U, \bar{U})$ and $\mathcal{F}(U', \bar{U}')$ coincide. In particular, if χ is the value of the pullback of ψ under the normalization map at some element y of the normalization, then the corresponding divisors obey the relation $D' \sim D + \infty^- + \infty^+ - y - \eta(y)$. Conversely, if ψ belongs to the kernel of $D^t(U, \bar{U}, k)$ and has no zero considered as a function from Δ to $\mathbb{C}^2$, then the function

$$\psi' = \frac{\begin{pmatrix} \bar{\phi}_2 \\ \bar{\phi}_1 \end{pmatrix}}{\phi_1\bar{\phi}_1 + \phi_2\bar{\phi}_2} \frac{\int_x^{x+\hat{\gamma}} \phi_2 \psi_1 dz + \phi_1 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k - k')) - 1} + \frac{\begin{pmatrix} \phi_1 \\ -\phi_2 \end{pmatrix}}{\phi_1\bar{\phi}_1 + \phi_2\bar{\phi}_2} \frac{\int_x^{x+\hat{\gamma}} \bar{\phi}_1 \psi_1 dz - \bar{\phi}_2 \psi_2 d\bar{z}}{\exp(2\pi\sqrt{-1}g(\hat{\gamma}, k + \bar{k}')) - 1}$$

is again the Baker–Akhiezer function of the pair of potentials $(U', \bar{U}')$ with

$$U' = \frac{(\partial\phi_2)\bar{\phi}_1 - \phi_2(\partial\bar{\phi}_1)}{\phi_1\bar{\phi}_1 + \phi_2\bar{\phi}_2}.$$

Again the complex Fermi curves $\mathcal{F}(U, \bar{U})$ and $\mathcal{F}(U', \bar{U}')$ coincide. If in particular ϕ is the value of the pullback of the transposed Baker–Akhiezer function under the normalization map at some element y of the normalization, then the corresponding divisors obey the relation $D' \sim D + y + \eta(y) - \infty^- - \infty^+$. Finally, the special cases of these two Bäcklund transformations, which correspond to the values of the pullback of the (transposed) Baker Akhiezer function at some element y of the normalization, are inverse to each other. **q.e.d.**

2.5.3 A compactification of the *isospectral sets*

For complex potentials U , the values of the *first integral* is equal to the square of the L^2 -norm times four. Hence, due to Lemma 2.6 the *isospectral set* of a *complex Fermi curve* corresponding to small values of the *first integral* is weakly compact. In the proof of Lemma 2.27 we actually showed that the isospectral set of complex potentials U is isomorphic to an open dense subset of one connected real component of the Picard group. Therefore, for small values of the *first integral* all elements of this connected real component of the Picard group obey condition *Divisor* (iii) (i. e. they are non special divisors of $\hat{Y}$). For two other classes of *complex Fermi curves* this is also true: in [Hi, Proposition 7.15] it is proven that for all *real- σ -hyperelliptic complex Fermi curves* (i. e. the normalization of the quotient Y_σ of Y modulo the holomorphic involution σ is isomorphic to $\mathbb{P}^1$ endowed with the anti-holomorphic involution with one real cycle) again all quaternionic complex line bundles of degree equal to the genus plus one are non-special. If a linear combination of the functions $\hat{p}$ and $\check{p}$ extends to a single-valued meromorphic functions on the Riemann surface Y , the same can be proven with the help of the arguments in [Sch, Theorem 8.3]. In all these cases the proof of Lemma 2.27 actually shows that the eigenfunctions corresponding to any element of the normalization of the *complex Fermi curve* has no zeroes considered as a $\mathbb{C}^2$ -valued function on Δ .

We shall see that not for all compact smooth Riemann surfaces Y obeying conditions *Quasi-momenta* (i)–(iii), the *isospectral sets* of all complex potentials U are compact. In contrast for example to the case where a linear combination of the functions $\hat{p}$ and $\check{p}$ extends to a single-valued meromorphic function, the L^2 -norms of the partial derivatives of the potential U , cannot be estimated in terms of the higher integrals. Moreover, later we shall consider non-connected compact Riemann surfaces Y fulfilling conditions *Quasi-momenta* (i)–(iii), which are not *complex Fermi curves*, i. e. the corresponding *isospectral sets* are empty. But the *isospectral sets* have a natural compactification, which we shall construct now.

Lemma 2.30. *Let Y be a compact smooth Riemann surface with two marked points ∞^- and ∞^+ and k a multi-valued meromorphic function, which is holomorphic on $Y \setminus \{\infty^-, \infty^+\}$. If Y, k, ∞^-, ∞^+ satisfies the conditions *Quasi-momenta* (i)–(iii), then the isospectral set has a natural compactification isomorphic to a real torus, whose dimension is equal to the*

genus of Y plus one. The elements of this compactified isospectral sets are integral divisors on $\mathbb{R}^2/\Lambda$ together with analytic complex potentials U on the complement of the support of the divisor. The corresponding Baker–Akhiezer functions yield a family of elements in the kernel of the restriction of the Dirac operator $D(U, \bar{U}, k)$ to the complement of the support of the divisor, which is parameterized by the corresponding complex Fermi curve. Finally, the first integral is equal to the square of the $L^2(\mathbb{R}^2/\Lambda)$ –norm of the potential times four plus 4π times the degree of the divisor.

The corresponding Baker–Akhiezer functions belong to the kernel of *Finite rank Perturbations* 3.22 of the Dirac operators $D(U, \bar{U}, k)$, whose support is the support of the divisor (compare with Section 3.4.1).

Proof. We extend the construction of the Baker–Akhiezer function in Lemma 2.27 to all divisors of $\tilde{Y}$, which obey conditions *Divisor* (i)–(ii). Here $\tilde{Y}$ is again obtained from Y by identifying the two marked points $\infty^\pm$ to an ordinary double point.

First we claim that these divisors obey condition *Divisor* (iii) for all $x \in \mathbb{R}^2/\Lambda$ with the exception of finitely many $x_1, \dots, x_L \in \mathbb{R}^2/\Lambda$. Due to the consideration in the proof of Lemma 2.27 it suffices to show, that for all divisors D on Y fulfilling conditions *Divisor* (i')–(ii'), there exists only finitely many $x_1, \dots, x_L \in \mathbb{R}^2/\Lambda$, for which the sheaf $\mathcal{L}(x) \otimes \mathcal{O}_{D-\infty^--\infty^+}$ has non-trivial global sections. Due to the Brill–Noether theory [A-C-G-H, Chapter IV] the set of all divisors on Y fulfilling condition *Divisor* (i'), which violate condition *Divisor* (iii'), is a subvariety of the $(g+1)$ –th symmetric product of Y . We conclude that for all divisors D on Y fulfilling conditions *Divisor* (i')–(ii') the set of all $x \in \mathbb{R}^2/\Lambda$, for which the sheaf $\mathcal{L}(x) \otimes \mathcal{O}_{D-\infty^--\infty^+}$ has non-trivial global sections is a real subvariety. Therefore, either this set is finite, or it contains locally an analytic one-dimensional submanifold. We exclude the latter case by contradiction. Without loss of generality we may assume that $x = 0$ belongs to the analytic submanifold, and that $\dim H^0(Y, \mathcal{L}(x) \otimes \mathcal{O}_{D-\infty^--\infty^+})$ is locally constant. In this case we have a flat family of locally free sheaves on Y parameterized by some open disc in $\mathbb{C}$. Due to [G-P-R, Chapter III. §4. Theorem 4.7] the direct image of this family has a section on the disc. The n –th derivatives of this section with respect to the deformation parameter yield global sections of $\mathcal{O}_{D+(n-1)\infty^--(n-1)\infty^+}$, which are not global sections of $\mathcal{O}_{D+(n-2)\infty^--(n-2)\infty^+}$. Since for all $n \in \mathbb{N}_0$ the divisor $D + (n-1)\infty^- + (n-1)\infty^+$ corresponds to a quaternionic complex line bundle, this implies that for all $n \in \mathbb{N}_0$

$$\dim H^0(Y, \mathcal{O}_{D+(n-1)\infty^--(n-1)\infty^+}) = 2n + \dim H^0(Y, \mathcal{O}_{D-\infty^--\infty^+}) \geq 2n + 2.$$

This contradicts to the Riemann–Roch theorem [Fo, Theorem 16.9 and Theorem 17.16].

We conclude that the Baker–Akhiezer function exists for all divisors $\tilde{D}$ on $\tilde{Y}$, which satisfy conditions *Divisor* (i)–(ii). But in general these Baker–Akhiezer functions have besides the poles on $\tilde{Y}$, which are independent of $x \in \mathbb{R}^2/\Lambda$, in addition finitely many poles $x_1, \dots, x_L$ on $\mathbb{R}^2/\Lambda$, which are independent of $y \in \tilde{Y}$. As in the usual case, on the complements of the second type of poles the asymptotic behaviour of the Baker–Akhiezer function determines a complex potential U , such that the value of the Baker–Akhiezer function at any element of $\tilde{Y} \setminus \{\infty^-, \infty^+\}$ restricted to this complement $(\mathbb{R}^2 \setminus \{x_1, \dots, x_L\})/\Lambda$ is an element of the

kernel of the Dirac operator $D(U, \bar{U}, k)$, where k is the value of the function k at the same element of $\tilde{Y} \setminus \{\infty^-, \infty^+\}$. Furthermore, the L^2 -norms of these potentials are bounded, since they are limits of potentials with bounded L^2 -norm.

If the divisor $\tilde{D}$ satisfies conditions *Divisor* (i)–(iii), then the *Baker–Akhiezer function* $\psi(x, y) = \begin{pmatrix} \psi_1(x, y) \\ \psi_2(x, y) \end{pmatrix}$ is the unique function with the following properties:

Baker–Akhiezer function (i) For all $x \in \mathbb{R}^2$ the restriction of $\psi(x, \cdot)$ to $Y \setminus \{\infty^-, \infty^+\}$ is a section of $\mathcal{O}_D$.

Baker–Akhiezer function (ii) For all $x \in \mathbb{R}^2$ the function $\exp(-2\pi\sqrt{-1}g(x, k))\psi(x, \cdot)$ is holomorphic in some neighbourhood of ∞^- and ∞^+ . At these points it takes the values $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$, respectively.

In some neighbourhood of $\infty^\pm$ we may expand the product $\exp(-2\pi\sqrt{-1}g(x, k))\psi(x, y)$ with respect to the local parameter $1/k_1$:

$$\exp(-2\pi\sqrt{-1}g(x, k))\psi(x, y) = \begin{cases} \begin{pmatrix} 0 \\ 1 \end{pmatrix} + \begin{pmatrix} \tilde{\psi}_{1,1}^-(x)/k_1 \\ \tilde{\psi}_{2,1}^-(x)/k_1 \end{pmatrix} + \mathbf{O}(1/k_1^2) & \text{at } \infty^- \\ \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} \tilde{\psi}_{1,1}^+(x)/k_1 \\ \tilde{\psi}_{2,1}^+(x)/k_1 \end{pmatrix} + \mathbf{O}(1/k_1^2) & \text{at } \infty^+. \end{cases}$$

Due to Lemma 3.12 the function k_2 has the asymptotic expansion

$$k_2 = \pm\sqrt{-1} \left(k_1 - \frac{\mathcal{W}}{8\pi^2 \text{vol}(\mathbb{R}^2/\Lambda)} 1/k_1 \right) + \mathbf{O}(1/k_1^2)$$

at $\infty^\pm$. Since the *Baker–Akhiezer function* lies in the kernel of the Dirac operator $D(U, \bar{U}, k)$, we obtain the equations

$$\begin{aligned} U(x) + 2\pi\sqrt{-1}\tilde{\psi}_{2,1}^+(x) &= 0 & U(x)\tilde{\psi}_{1,1}^-(x) + \frac{\sqrt{-1}\mathcal{W}}{8\pi \text{vol}(\mathbb{R}^2/\Lambda)} + \partial\tilde{\psi}_{2,1}^-(x) &= 0 \\ \bar{U}(x)\tilde{\psi}_{2,1}^+(x) - \frac{\sqrt{-1}\mathcal{W}}{8\pi \text{vol}(\mathbb{R}^2/\Lambda)} - \bar{\partial}\tilde{\psi}_{1,1}^+(x) &= 0 & \bar{U} - 2\pi\sqrt{-1}\tilde{\psi}_{1,1}^-(x) &= 0 \end{aligned}$$

These equations show that the *first integral* is equal to $\mathcal{W} = 4 \int_{\mathbb{R}^2/\Lambda} U(x)\bar{U}(x)d^2x$.

We shall modify this calculation of the *Baker–Akhiezer function* in the case of poles with respect to x . In this case the *Baker–Akhiezer function* satisfies conditions *Baker–Akhiezer function* (i)–(ii) only for all x in the complement of $(x_1 + \Lambda) \cup \dots \cup (x_L + \Lambda)$. This function has a unique meromorphic extension to the product of a small neighbourhood of $\mathbb{R}^2$ in $\mathbb{C}^2$ with $Y \setminus \{\infty^-, \infty^+\}$. Moreover, the product $\exp(-2\pi\sqrt{-1}g(x, k))\psi(x, y)$ has also a meromorphic extension to the product of a small neighbourhood of $\mathbb{R}^2$ in $\mathbb{C}^2$ with some neighbourhood of $\infty^\pm$. We shall see that the functions $\tilde{\psi}_{1,1}^+$ and $\tilde{\psi}_{2,1}^-$ have only first order poles at the points $x_1, \dots, x_L$. Dolbeault’s Lemma [Gu-Ro, Chapter I Section D 2. Lemma] and the equations

above imply that in this case the *first integral* is equal to $\mathcal{W} = 4 \int_{\Delta \setminus \{x_1, \dots, x_L\}} U(x) \bar{U}(x) d^2x$ minus $8\pi^2 \sqrt{-1}$ times the sum over all Laurent coefficients of $\tilde{\psi}_{1,1}^+(x)$ and $\tilde{\psi}_{2,1}^-(x)$ at $x_1, \dots, x_L$ corresponding to the singular term $((x - x_l)_1 \pm \sqrt{-1}(x - x_l)_2)^{-1}$. In order to calculate these Laurent coefficients it is convenient to consider the limit

$$\psi_l^\pm(x, k_1) = \lim_{\lambda \rightarrow \infty} \exp(-2\pi\sqrt{-1}g(x_l, k)) \psi(x_l + x/\lambda, k_1\lambda)$$

in the neighbourhood of $\infty^\pm$, respectively. Here we parameterize small neighbourhoods of $\infty^\pm$ by the local parameters $1/k_1$. More precisely, the *Baker–Akhiezer function* extends to a function on a neighbourhood of $\mathbb{R}^2 \times \infty^\pm$ in $\mathbb{C}^2 \times Y$ with prescribed essential singularity at $\infty^\pm$, which is meromorphic with respect to $x \in \mathbb{C}^2$. The functions $\psi_l^\pm(x, k_1)$ are the pullbacks of the products of $\exp(-2\pi\sqrt{-1}g(x_l, k))$ with the *Baker–Akhiezer function* to the blowing up of $\mathbb{C}^2 \times Y$ at the points $x_l \times \infty^\pm$ (compare with [Har, Chapter I Section 4]). The condition *Baker–Akhiezer function* (ii) determines the values of the products of these functions with $\exp(-2\pi\sqrt{-1}g(x, k))$ at $k_1 = \infty$. This implies that these products are meromorphic. Due to Lemma 2.27 the potential U is a limit of potentials with finite L^2 -norms. Therefore U has finite L^2 -norm. Moreover, the limits $\lim_{\lambda \rightarrow \infty} U(x_l + x/\lambda)/\lambda$ are equal to zero for $x \neq 0$. Furthermore, since on the complement of $(x_1 + \Lambda) \cup \dots \cup (x_L + \Lambda)$ the *Baker–Akhiezer functions* belong to the kernel of the Dirac operator, for $x \neq 0$ the functions $\psi_l^\pm$ belong to the kernel of $\begin{pmatrix} 0 & \partial \\ -\partial & 0 \end{pmatrix}$. This implies that the functions $\psi_l^\pm(x, k_1)$ are of the form

$$\psi_l^\pm(x, k_1) = q_l(1/z^\pm) \exp(z^\pm) \begin{pmatrix} (1 \pm 1)/2 \\ (1 \mp 1)/2 \end{pmatrix} \text{ with } z^\pm = 2\pi\sqrt{-1}k_1(x_1 \pm \sqrt{-1}x_2).$$

Here q_l is a rational function with respect to $1/z^\pm \in \mathbb{P}^1$, which is at $z^\pm = \infty$ equal to 1, and has poles only at $z^\pm = 0$.

Remark 2.31. *The non-trivial components of these functions can be considered as Baker–Akhiezer functions of rational solutions of the Kadomtsev–Petviashvili Equation. For example the non-trivial components of the functions $\psi_l^+(x, k)$ corresponding to the cases $\mathcal{N}_l = \{n \in \mathbb{Z} \mid -d \leq n\} \setminus \{-d+1, -d+3, \dots, d-1\}$ are the Baker–Akhiezer functions of the rational solutions of the KdV equation corresponding to the Lax operators $(\partial)^2 - d(d+1)/(x_1 + \sqrt{-1}x_2)^2$ [A-McK-M]. The blowing up extracts from the Baker–Akhiezer function exactly that part, which describes the singularity at x_l , and shifts the potential to infinity. Therefore, these Baker–Akhiezer functions are elements of the kernel of finite rank perturbations of the Dirac operator without potential on $\mathbb{P}^1$ (compare with Section 3.4.1). The corresponding spectral curves have disconnected normalization similar to the complex Fermi curves with disconnected normalization. The former Baker–Akhiezer functions, however, correspond to rational solutions of the Kadomtsev–Petviashvili Equation, and the latter to elliptic solutions.*

For all x_l let $\mathcal{N}_l \subset \mathbb{Z}$ be the unique subset of the integer numbers with the property that

$$\dim H^0(Y, \mathcal{O}_{D(x_l) + n(\infty^- + \infty^+)}) - \dim H^0(Y, \mathcal{O}_{D(x_l) + (n-1)(\infty^- + \infty^+)}) = \begin{cases} 2 & \text{if } n \in \mathcal{N}_l \\ 0 & \text{if } n \notin \mathcal{N}_l. \end{cases}$$

We should remark that the space of global sections of the sheaves $\mathcal{O}_{D(x_l)+n\infty^-+n\infty^+}$ have even dimension, since they correspond to quaternionic complex line bundles. Due to the Riemann–Roch theorem [Fo, Theorem 16.9 and Theorem 17.16] $\mathcal{N}_l$ has as many negative elements as there exists non-negative natural numbers, which does not belong to $\mathcal{N}_l$. Therefore, the Laurent expansion of $\psi_l^\pm$ at $z^\pm = \infty$ has no term of the form $(z^\pm)^n$, if $n \notin \mathcal{N}_l$. These properties uniquely determine the functions $\psi_l^\pm$. The corresponding rational functions q_l are polynomials with integer coefficients. In fact, if $-d_l$ denotes the smallest element of $\mathcal{N}_l$, then q_l has to be a polynomial of degree d_l , whose lowest coefficient is equal to one. Moreover, since $\mathcal{N}_l$ does not contain exactly d_l elements of the subset of all natural numbers, which are larger than $-d_l$, this polynomial has to obey exactly d_l equations. If $1, a_1, \dots, a_{d_l}$ are the coefficients of the polynomial q_l , then the polynomial

$$p_l(z) = (z+1)(z+2)\dots(z+d_l) + a_1((z+2)\dots(z+d_l)) + \dots + a_{d_l}$$

has to vanish at all $n \in \{n \in \mathbb{Z} \mid -d_l \leq n\} \setminus \mathcal{N}_l$. Since $\{n \in \mathbb{Z} \mid -d_l \leq n\} \setminus \mathcal{N}_l$ has exactly d_l elements, the polynomial p_l is the unique polynomial with highest coefficient 1, which vanishes at all elements of $\{n \in \mathbb{Z} \mid -d_l \leq n\} \setminus \mathcal{N}_l$. Obviously the coefficients of q_l are integer linear combinations of the coefficients of p_l . Therefore, the set $\mathcal{N}_l$ uniquely determines the polynomial q_l , and these polynomials have integer coefficients. Moreover, a_1 is equal to $-d_l(d_l+1)/2$ minus the sum over all elements of $\{n \in \mathbb{Z} \mid -d_l \leq n\} \setminus \mathcal{N}_l$. This natural number is equal to the sum over all elements in $\mathcal{N}_l \cap -\mathbb{N}$ minus the sum over all elements in $\mathbb{N}_0$, which does not belong to $\mathcal{N}_l$. We define the local contribution to the Willmore energy (compare with [P-P, B-F-L-P-P, F-L-P-P]) at x_l of the *Baker–Akhiezer function* $\mathcal{W}_{\text{sing}, x_l}(\psi)$ as this natural number times -4π , which is always non-negative, and zero if and only if $\mathcal{N}_l = \mathbb{N}_0$. We conclude that the *first integral* is equal to

$$\mathcal{W} = 4 \int_{\mathbb{R}^2/\Lambda} U(x) \bar{U}(x) d^2x + \mathcal{W}_{\text{sing}} \quad \text{with } \mathcal{W}_{\text{sing}} = \sum_{l=1}^L \mathcal{W}_{\text{sing}, x_l}(\cdot).$$

We consider the points $x_1, \dots, x_L$ as singularities of the holomorphic structure of the corresponding quaternionic line bundle in the sense of ‘quaternionic function theory’ (compare with [P-P, B-F-L-P-P, F-L-P-P]). In analogy to the *Local contribution to the arithmetic genus* 4.3 of singularities in the sense of complex function theory these singularities of ‘quaternionic function theory’ have in addition a local contribution to the Willmore energy. **q.e.d.**

Due to this formula the *isospectral sets* of all *complex Fermi curves* of finite *geometric genus*, whose *first integral* is smaller than 4π are compact (compare with Remark 2.5 and Remark 2.8). Moreover, due to the explicit calculation of $\mathcal{W}_{\text{sing}, x_l}(\cdot)$ in terms of $\mathcal{N}_l$ it is possible to classify all $\mathcal{N}_l$, with the property that the corresponding $\mathcal{W}_{\text{sing}, x_l}(\cdot) \leq 4n\pi$. For some given $\mathcal{N}_l$ the local contribution to the Willmore energy $\mathcal{W}_{\text{sing}, x_l}(\cdot)$ is equal to the sum over all elements of $\mathbb{N}_0$, which do not belong to $\mathcal{N}_l$, minus the sum over all elements of $\mathcal{N}_l \cap -\mathbb{N}$. For some given d_l the corresponding $\mathcal{W}_{\text{sing}, x_l}(\cdot)$ satisfies the inequality $\mathcal{W}_{\text{sing}, x_l}(\cdot) \geq 4d_l\pi$, and equality holds if and only if $\mathcal{N}_l = \mathbb{N} \cup \{-d_l\}$. The possible $\mathcal{N}_l$ ’s with some given $\mathcal{W}_{\text{sing}, x_l}(\cdot)$

decompose into groups with $m = 1, \dots, M$ negative elements, where

$$2M(M+1)\pi + 2M(M-1)\pi = 4M^2\pi \leq \mathcal{W}_{\text{sing},x_l}(\cdot).$$

Hence for each m with $4m^2\pi \leq \mathcal{W}_{\text{sing},x_l}(\cdot)$ the sum Σ_l^- of all negative elements of $\mathcal{N}_l$ and the sum Σ_l^+ of all elements of $\mathbb{N}_0$, which do not belong to $\mathcal{N}_l$ satisfy

$$\mathcal{W}_{\text{sing},x_l}(\cdot) = \Sigma_l^+ - \Sigma_l^- \quad \text{and} \quad -\frac{\mathcal{W}_{\text{sing},x_l}(\cdot)}{4\pi} + m(m-1)/2 \leq \Sigma_l^- \leq -m(m+1)/2.$$

Hence for each m with $4m^2\pi \leq \mathcal{W}_{\text{sing},x_l}(\cdot)$ there are exactly $\mathcal{W}_{\text{sing},x_l}(\cdot)/(4\pi) - m^2 + 1$ possible values of Σ_l^- and Σ_l^+ . Moreover, for some given m , Σ_l^- and Σ_l^+ , there are exactly so many $\mathcal{N}_l$'s, as the number of choices of m different elements of $-\mathbb{N}$, whose sum is equal to Σ_l^- , times the number of choices of m different elements of $\mathbb{N}_0$, whose sum is equal to Σ_l^+ . In the following table we state all possible $\mathcal{N}_l$'s, whose local contribution to the Willmore energy $\mathcal{W}_{\text{sing},x_l}(\cdot)$ is not larger than 20π . Instead of $\mathcal{N}_l$ we state all negative elements of $\mathcal{N}_l$ together with all elements of $\mathbb{N}_0$, which does not belong to $\mathcal{N}_l$:

$\mathcal{W}_{\text{sing},x_l}(\cdot)$	$m = 1$	$m = 2$
4π	$-1, 0$	
8π	$-1, 1 \quad -2, 0$	
12π	$-1, 2 \quad -2, 1 \quad -3, 0$	
16π	$-1, 3 \quad -2, 2 \quad -3, 1 \quad -4, 0$	$-2, -1, 0, 1$
20π	$-1, 4 \quad -2, 3 \quad -3, 2 \quad -4, 1 \quad -5, 0$	$-2, -1, 0, 2 \quad -3, -1, 0, 1$

On Riemann surfaces there exists a one-to-one correspondence between line bundles and divisors [Fo, Chapter III. §29.11], which is compatible with the degrees of line bundles and divisors. With the help of this correspondence those elements in the compactification of the *isospectral sets*, which do not belong to the *isospectral sets*, can be shown to correspond to Dirac operators with potentials acting on topological non-trivial bundles. These generalized Dirac operators occur also in ‘quaternionic function theory’ developed by F. Pedit and U. Pinkall, which describes immersions into $\mathbb{R}^4$.

If a *complex Fermi curve* has singularities, then the *isospectral set* contains besides the potentials considered in Lemma 2.27, whose corresponding *Structure sheaf* 2.24 is the normalization sheaf of the *complex Fermi curve* in addition potentials, whose corresponding *Structure sheaf* 2.24 differs from the normalization sheaf. Due to Corollary 2.17 (ii), on the one-sheeted coverings introduced in *Structure sheaf* 2.24 corresponding to potentials of the form $(U, \bar{U})$ the involution η does not have fixed points. Due to Lemma 2.26 all one-sheeted coverings of the *complex Fermi curve* of finite *geometric genus* have finite arithmetic genus. The proofs of Lemma 2.27 and Lemma 2.30 carry over to these one-sheeted coverings and show that for any such one-sheeted covering the real component of the Picard group, whose degree is equal to the arithmetic genus of this one-sheeted covering plus one, parameterizes a family of generalized potentials in the sense of Lemma 2.30. More precisely, sue to [Sch, Chapter 9] all sheaves, which correspond to some potential with a given *Structure sheaf* 2.24, are characterized as coherent subsheaves of the sheaf of meromorphic functions, whose degree is equal to the corresponding arithmetic genus, if the two infinities are identified to an

ordinary double point. But for our purpose it is enough to consider only those potentials, which correspond to locally free rank-one sheaves. For any data $(Y, \infty^-, \infty^+, k)$ fulfilling *Quasi-momenta* (i)–(iii) let $\mathcal{W}_{\text{sing,max}}(Y, \infty^-, \infty^+, k)$ denote the maximal contribution of the singularities to the Willmore energy of the corresponding compactified *isospectral set* described analogous to those described in Lemma 2.30, whose *Structure sheaf* 2.24 coincides with the structure sheaf of Y . Due to Lemma 2.30 this maximal contribution of the singularities to the Willmore energy obeys the inequality

$$\mathcal{W}_{\text{sing,max}}(Y, \infty^-, \infty^+, k) \leq \mathcal{W}(\mathcal{F}) = \mathcal{W}(Y, \infty^-, \infty^+, k).$$

A *Baker–Akhiezer function* has a pole at $x \in \mathbb{R}^2/\Lambda$, if and only if the divisor $D(x) - \infty^- - \infty^+$ is the pole divisor of a function f fulfilling $f\eta^*\bar{f} = -1$, whose degree is equal to the arithmetic genus g of Y minus a positive odd number. More precisely, the minimum of the degrees of all functions f fulfilling $f\eta^*\bar{f} = -1$, whose degree is equal to $g + 1$ modulo two, is larger or equal to $g + 1 - 2\mathcal{W}_{\text{sing,max}}(Y, \infty^-, \infty^+, k)/(4\pi)$. Moreover, the maximal contribution of the singularities to the Willmore energy does not vanish, if and only if there exists a function f fulfilling $f\eta^*\bar{f} = -1$, whose degree is equal to g minus a positive odd number.

If in addition $(Y, \infty^-, \infty^+, k)$ fulfills condition *Quasi-momenta* (iv), then the following condition implies that the corresponding (generalized) potentials are real:

Divisor (iv) There exists a function g on $\tilde{Y}$ which takes the same values at $\infty^\pm$, whose divisor is equal to the zero divisor of dk_2 minus $\tilde{D} + \sigma(\tilde{D})$.

Therefore, the real part of the compactification of the *isospectral sets* are real tori, which are isomorphic to the Prym varieties of $\tilde{Y}$ together with the holomorphic involution σ (see e. g. [L-B, Chapter 12]). In particular, if the *isospectral sets* are compact (e. g. if the *complex Fermi curve* is *real- σ -hyperelliptic*, or if a linear combination of $\hat{p}$ and $\check{p}$ extends to a single-valued meromorphic function, or if the *first integral* is smaller than 4π) then the real part of the *isospectral* is not empty. We do not know whether there exist *complex Fermi curves*, whose *isospectral sets* are not empty, but whose real parts are empty.

3 The moduli space

3.1 Variations of *complex Fermi curves*

In this section we want to investigate the behaviour of the map $(V, W) \mapsto \mathcal{F}(V, W)$ under small variations of the potentials. In particular, we will characterize those variations $(\delta V, \delta W)$ of the potentials, which induce no variation of the *complex Fermi curve*. For this purpose we use the *Fundamental domain* 2.1 and the corresponding notations. If δV and δW are variations of the potentials V and W , then we may define a one-form on the *complex Fermi curve* $\mathcal{F}(V, W)$ by the formula

$$\Omega_{V,W}(\delta V, \delta W) = \delta\check{p} \cdot d\hat{p} - \delta\hat{p} \cdot d\check{p}.$$

Since in general the *complex Fermi curves* of deformed potentials are also deformed, it is not clear a priori whether this definition makes sense. For example, either the condition $\delta\hat{p} = 0$ or the condition $\delta\check{p} = 0$ may be used to identify locally the *complex Fermi curve* of the deformed potential with the undeformed *complex Fermi curve*. In the first case we may consider $\check{p}$ locally as a function of $\hat{p}$ and a deformation parameter t and in the second case $\hat{p}$ locally as a function of $\check{p}$ and t . In the first case $\Omega_{V,W}(\delta V, \delta W)$ equals $\frac{\partial \check{p}}{\partial t} d\hat{p}$ and in the second case it equals $-\frac{\partial \hat{p}}{\partial t} d\check{p}$. But as shown in [G-S-1, Section 1.3] the form does not depend in which way we identify the different *complex Fermi curves*. In fact, due to Theorem 2.3 locally the *complex Fermi curves* may be described by an equation of the form $R(\hat{p}, \check{p}) = 0$. Moreover, if d is the number of sheets of the covering map $\hat{p}$, then due to the Weierstraß Preparation theorem [G-P-R, Chapter I. Theorem 1.4] R can be chosen to be a polynomial with respect to $\check{p}$ of degree d , whose coefficients are holomorphic functions depending on $\hat{p}$, and whose highest coefficient is equal to 1. The equations

$$\begin{aligned} dR(\hat{p}, \check{p}) &= \delta R(\hat{p}, \check{p}) + \frac{\partial R(\hat{p}, \check{p})}{\partial \hat{p}} \delta \hat{p} + \frac{\partial R(\hat{p}, \check{p})}{\partial \check{p}} \delta \check{p} = 0 \text{ and} \\ \frac{\partial R(\hat{p}, \check{p})}{\partial \hat{p}} d\hat{p} + \frac{\partial R(\hat{p}, \check{p})}{\partial \check{p}} d\check{p} &= 0 \end{aligned}$$

imply the equation

$$\delta \check{p} d\hat{p} - \delta \hat{p} d\check{p} = -\delta R(\hat{p}, \check{p}) \frac{d\hat{p}}{\frac{\partial R(\hat{p}, \check{p})}{\partial \check{p}}} = \delta R(\hat{p}, \check{p}) \frac{d\check{p}}{\frac{\partial R(\hat{p}, \check{p})}{\partial \hat{p}}}.$$

Hence due to Lemma 2.20 $\Omega_{V,W}(\delta V, \delta W)$ is a well defined regular form. If we consider λ locally as a function of $\hat{p}, \check{p}$, and t , then the choice $\delta\hat{p} = 0$ yields

$$\begin{aligned} \Omega_{V,W}(\delta V, \delta W) &= \frac{\partial \lambda}{\partial t} \frac{d\hat{p}}{\frac{\partial \lambda}{\partial \hat{p}}} = \text{tr} \left(P \begin{pmatrix} \delta V & 0 \\ 0 & \delta W \end{pmatrix} \right) \frac{d\hat{p}}{\frac{\partial \lambda}{\partial \hat{p}}} \\ &= \text{tr} \left(P_{\hat{\gamma}} \begin{pmatrix} 0 & \frac{\delta W}{\kappa_2 - \sqrt{-1}\kappa_1} \\ \frac{\delta V}{\kappa_2 + \sqrt{-1}\kappa_1} & 0 \end{pmatrix} \right) \frac{d\hat{p}}{\pi}. \end{aligned}$$

Here we have used the relation between P and $P_{\hat{\gamma}}$ established in the proof of Lemma 2.23.

Lemma 3.1. *Let $\delta V \in L^2(\mathbb{R}^2/\Lambda)$ and $\delta W \in L^2(\mathbb{R}^2/\Lambda)$ be variations of V and W , respectively. Then $\Omega_{V,W}(\delta V, \delta W) = \text{tr} \left(P_{\hat{\gamma}} \begin{pmatrix} 0 & \frac{\delta W}{\kappa_2 - \sqrt{-1}\kappa_1} \\ \frac{\delta V}{\kappa_2 + \sqrt{-1}\kappa_1} & 0 \end{pmatrix} \right) \frac{d\hat{p}}{\pi}$ is a regular one-form, on $\mathcal{F}(V, W)$, independent of the choice of $\hat{\gamma}$ and $\check{\gamma}$.*

Proof. By Lemma 2.23 (iv), $P_{\hat{\gamma}} d\hat{p}$ is a regular one-form on $\mathcal{F}(V, W)$. Thus also $\Omega_{V,W}(\delta V, \delta W)$ is a regular one-form. All matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ induce the modular transformations:

$$\begin{aligned} d\hat{p}_{\Delta'} &= a d\hat{p}_{\Delta} + b d\check{p}_{\Delta} & \frac{\partial}{\partial \hat{p}_{\Delta'}} &= d \frac{\partial}{\partial \hat{p}_{\Delta}} - c \frac{\partial}{\partial \check{p}_{\Delta}} \\ d\check{p}_{\Delta'} &= c d\hat{p}_{\Delta} + d d\check{p}_{\Delta} & \frac{\partial}{\partial \check{p}_{\Delta'}} &= -b \frac{\partial}{\partial \hat{p}_{\Delta}} + a \frac{\partial}{\partial \check{p}_{\Delta}} \end{aligned}$$

On the *complex Fermi curve* we have the following identity:

$$\frac{\partial \lambda}{\partial \hat{p}} d\hat{p} + \frac{\partial \lambda}{\partial \check{p}} d\check{p} = 0.$$

Therefore we obtain

$$\frac{d\hat{p}_{\Delta'}}{\frac{\partial \lambda}{\partial \hat{p}_{\Delta'}}} = \frac{a \cdot d\hat{p}_{\Delta} + b \cdot d\check{p}_{\Delta}}{a \frac{\partial \lambda}{\partial \hat{p}_{\Delta}} - b \frac{\partial \lambda}{\partial \check{p}_{\Delta}}} = \frac{a \frac{\partial \lambda}{\partial \check{p}_{\Delta}} d\hat{p}_{\Delta} + b \frac{\partial \lambda}{\partial \check{p}_{\Delta}} d\check{p}_{\Delta}}{\frac{\partial \lambda}{\partial \check{p}_{\Delta}} \left(a \frac{\partial \lambda}{\partial \hat{p}_{\Delta}} - b \frac{\partial \lambda}{\partial \check{p}_{\Delta}} \right)} = \frac{\left(a \frac{\partial \lambda}{\partial \check{p}_{\Delta}} - b \frac{\partial \lambda}{\partial \check{p}_{\Delta}} \right) d\hat{p}_{\Delta}}{\frac{\partial \lambda}{\partial \check{p}_{\Delta}} \left(a \frac{\partial \lambda}{\partial \hat{p}_{\Delta}} - b \frac{\partial \lambda}{\partial \check{p}_{\Delta}} \right)} = \frac{d\hat{p}_{\Delta}}{\frac{\partial \lambda}{\partial \hat{p}_{\Delta}}}.$$

On the other hand it follows from the proof of (iii) of Lemma 2.23 that $\Omega_{V,W}(\delta V, \delta W)$ is equal to

$$\Omega_{V,W}(\delta V, \delta W) = \text{tr} \left(P \begin{pmatrix} \delta V & 0 \\ 0 & \delta W \end{pmatrix} \right) \frac{d\hat{p}}{\frac{\partial \lambda}{\partial \hat{p}}},$$

and does not change under the modular transformations.

q.e.d.

This map $\Omega_{V,W}$ should be considered as the derivative of the map $(V, W) \mapsto \mathcal{F}(V, W)$. Due to a general feature of the integrable systems having Lax operators, the dual of this map is the infinitesimal action of the Lie algebra of the Picard group on the *isospectral sets*. More precisely, S erre duality is a pairing between the space of holomorphic one-forms and the Lie algebra of the Picard group [Fo,  17. and  21.6], and on the other hand the symplectic form is a pairing on the tangent space of the phase space of the integrable system. Therefore, the dual of the map $\Omega_{V,W}$ with respect to these pairings is a map from the Lie algebra of the Picard group into the tangent space of the phase space. Furthermore, the *isospectral sets* correspond to open dense subsets of one component of the Picard group (with specified degree). Therefore, the Picard group acts on the *isospectral sets*, and the derivative of this action yields a map from the Lie algebra of the Picard group into the tangent space of the phase space. The Hamiltonian structure fits together with the function theoretic properties of the spectral curves, and the second map is the dual of the first map. To work out this structure completely implies a lot of analytical work (compare with [Sch]). In the following lemma we define the infinitesimal action of the Lie algebra of the Picard group on the *isospectral sets*. We show that this map is the dual map of $\Omega_{V,W}$ with respect to S erre duality and the symplectic form. For this purpose we represent the elements of the Lie algebra of the Picard group by those data, which enter in the first Cousin Problem (see e. g. [Ah, Chapter 5 2.1.] and [G-P-R, Chapter III.  3.3.]). These data are the singular parts of local meromorphic functions, sometimes called Mittag-Leffler distributions. If we identify those data with zero, which are the singular parts of global meromorphic functions, the resulting space is canonically isomorphic with the Lie algebra of the Picard group [Fo,  18.2].

We shall use the notations introduced in the context of the *Fundamental domain* 2.1, but the operators under consideration act on the Hilbert bundle corresponding to the *Trivialization* 2.2.

Lemma 3.2. *Let f be a meromorphic function with finitely many poles on some open neighbourhood $\mathcal{U}$ of $\mathcal{F}(V, W)/\Lambda^*$. Then there exists a meromorphic function $\tilde{\mathbf{A}}_f(\hat{p})$ from the complex plane $\hat{p} \in \mathbb{C}$ into the bounded operators from the Banach spaces $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ into $L^{p'}(\mathbb{R}^2/\Lambda) \times L^{p'}(\mathbb{R}^2/\Lambda)$ for all $1 < p' < p < \infty$ with the following properties:*

(i) *For all $(n_1, n_2) \in \mathbb{Z}^2$ and all $\hat{p} \in \mathbb{C}$ the following identity is valid:*

$$\tilde{\mathbf{A}}_f(\hat{p})\psi_{n_1\hat{\kappa}+n_2\tilde{\kappa}} = \psi_{n_1\hat{\kappa}+n_2\tilde{\kappa}}\tilde{\mathbf{A}}_f(\hat{p} + n_1).$$

(ii) *The commutator $[\tilde{\mathbf{A}}_f(\hat{p}), \tilde{\mathbf{D}}_{\hat{\gamma}, \tilde{\gamma}}(V, W, \hat{p})]$ does not depend on $\hat{p}$ and is equal to an operator*

$$\text{of the form } \delta\tilde{\mathbf{D}}_{\hat{\gamma}, \tilde{\gamma}}(V, W, \cdot) = \begin{pmatrix} 0 & \frac{\delta W_f}{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1} \\ \frac{\delta V_f}{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1} & 0 \end{pmatrix}, \text{ with some variations } \delta V_f, \delta W_f \in$$

$\bigcap_{q < \infty} L^q(\mathbb{R}^2/\Lambda)$. Moreover, this variation does not change the complex Fermi curve.

Equivalently, the form $\Omega_{V, W}(\delta V_f, \delta W_f)$ is identically zero.

(iii) *For all elements $\tilde{\chi}$ and $\tilde{\xi}$ of $\bigcup_{r \geq 2} L^r(\mathbb{R}^2/\Lambda) \times L^r(\mathbb{R}^2/\Lambda)$ and all $\varepsilon, \delta > 0$ there exists a $\delta' > 0$, such that for all $\hat{p}$ in*

$$\{\hat{p} \in \mathbb{C} \mid |\hat{p} - \hat{p}'| > \delta \pmod{\mathbb{Z}} \text{ for all values } \hat{p}' \text{ of the function } \hat{p} \text{ at the poles of } f\}$$

with $|\hat{p}| > \delta'$ the function $|\langle \tilde{\chi}, \tilde{\mathbf{A}}(\hat{p})\tilde{\xi} \rangle|$ is bounded by ε .

(iv) *For all variations $\delta V, \delta W \in L^2(\mathbb{R}^2/\Lambda)$ the total residue of the form $f \cdot \Omega_{V, W}(\delta V, \delta W)$ on $\mathcal{U}$ is equal to the value of a symplectic form at $(\delta V, \delta W)$ and $(\delta V_f, \delta W_f)$:*

$$\text{Res}_{\mathcal{U}}(f \cdot \Omega_{V, W}(\delta V, \delta W)) = \frac{1}{2\pi^2\sqrt{-1}} \int_{\mathbb{R}^2/\Lambda} (\delta V \delta W_f - \delta W \delta V_f) d^2x.$$

Proof. If f is a linear combination of two functions with two associated functions fulfilling conditions (i)–(iv), then the linear combination $\tilde{\mathbf{A}}_f$ of the two associated functions has the required properties. Hence we may assume that f has a pole at one point $[k'] \in \mathcal{F}(V, W)/\Lambda^*$. Let $\tilde{\mathbf{F}}(\cdot)$ denote the local sum of $f \cdot \tilde{\mathbf{P}}_{\hat{\gamma}}$ over all sheets of $\mathcal{F}(V, W)/\Lambda^*$ considered as a covering space over $\hat{p} \in \mathbb{C}$, which contain the element $[k']$ (compare with Remark 2.19). By definition this is a meromorphic function from some neighbourhood of $\hat{p}'$ into the finite rank operators on $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$. Therefore, the singular part $\tilde{\mathbf{F}}_{\text{singular}}(\cdot)$ is a meromorphic function on the whole plane $\hat{p} \in \mathbb{C}$. Now we claim that the infinite sum

$$\tilde{\mathbf{A}}_f(\hat{p}) = \sum_{(n_1, n_2) \in \mathbb{Z}^2} \psi_{n_1\hat{\kappa}+n_2\tilde{\kappa}} \tilde{\mathbf{F}}_{\text{singular}}(\hat{p} + n_1) \psi_{-n_1\hat{\kappa}-n_2\tilde{\kappa}}$$

converges in the strong operator topology. Due to Theorem 2.3 the operator-valued function $\tilde{\mathbf{F}}_{\text{singular}}(\hat{p})$ is a finite sum of operators of the form $\frac{|\tilde{\psi}\rangle\langle\tilde{\phi}|}{(\hat{p}-\hat{p}')^l} : \tilde{\chi} \mapsto \langle\langle\tilde{\phi}, \tilde{\chi}\rangle\rangle(\hat{p} - \hat{p}')^{-l}\tilde{\psi}$ with

elements $\tilde{\phi}$ and $\tilde{\psi}$ of the Banach spaces $\bigcap_{q < \infty} L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$. For all $l \in \mathbb{N}$ and $\hat{q} \in [0, 1]$ the sum $d_l(\hat{p}, \hat{q}) = \sum_{n \in \mathbb{Z}} (\hat{p} - \hat{p}' + n)^{-l} \exp(2n\pi\sqrt{-1}\hat{q})$ converges. Let us calculate these limits. Fourier decomposition gives the relation

$$d_1(\hat{p}, \hat{q}) = \frac{2\pi\sqrt{-1} \exp(2\pi\sqrt{-1}(\hat{p}' - \hat{p})\hat{q})}{1 - \exp(2\pi\sqrt{-1}(\hat{p}' - \hat{p}))} \quad \forall \hat{q} \in [0, 1].$$

Moreover, they satisfy the recursion equation

$$2\pi\sqrt{-1}(\hat{p} - \hat{p}')d_{l+1}(\hat{p}, \hat{q}) + \frac{\partial d_{l+1}(\hat{p}, \hat{q})}{\partial \hat{q}} = 2\pi\sqrt{-1}d_l(\hat{p}, \hat{q}) \quad \forall \hat{q} \in \mathbb{R}/\mathbb{Z}.$$

Due to the Riemann–Lebesgue Lemma [R-S-II, Theorem IX.7] for $l > 1$ the functions $d_l(\hat{p}, \hat{q})$ are continuous and periodic with period 1 with respect to $\hat{q}$. This implies the following recursion relation for the functions $\tilde{d}_{l+1}(\hat{p}, \hat{q}) = \exp(2\pi\sqrt{-1}(\hat{p} - \hat{p}')\hat{q}) d_l(\hat{p}, \hat{q})$:

$$\tilde{d}_l(\hat{p}, \hat{q}) = 2\pi\sqrt{-1} \left(\int_0^{\hat{q}} \tilde{d}_l(\hat{p}, \hat{q}') d\hat{q}' + \frac{1}{(\exp(2\pi\sqrt{-1}(\hat{p} - \hat{p}')) - 1)} \int_0^1 \tilde{d}_l(\hat{p}, \hat{q}') d\hat{q}' \right).$$

Since $\tilde{d}_1(\hat{p}, \hat{q})$ is equal to $(2\pi\sqrt{-1}) (1 - \exp(2\pi\sqrt{-1}(\hat{p}' - \hat{p})))^{-1}$, these functions $\tilde{d}_l(\hat{p}, \hat{q})$ are polynomials with respect to $\hat{q}$ and $(\exp(2\pi\sqrt{-1}(\hat{p}' - \hat{p})) - 1)^{-1}$ times $\tilde{d}_1(\hat{p}, \hat{q})$. The unique solution of this recursion is given by the generating function

$$\sum_{l \in \mathbb{N}} t^{l-1} \tilde{d}_l(\hat{p}, \hat{q}) = \frac{(2\pi\sqrt{-1}) (1 - \exp(2\pi\sqrt{-1}(\hat{p}' - \hat{p})))^{-1} \exp(2\pi\sqrt{-1}\hat{q}t)}{1 - (\exp(2\pi\sqrt{-1}(\hat{p}' - \hat{p})) - 1)^{-1} (\exp(2\pi\sqrt{-1}t) - 1)}.$$

In fact, a transformation of the recursion relation

$$f_{l+1}(x) = a \left(\int_0^x f_l(x') dx' + b \int_0^1 f_l(x') dx' \right) \quad \text{with } f_1(x) = c$$

into an integral equation of the corresponding generating function yields the unique solution

$$\sum_{l \in \mathbb{N}} t^{l-1} f_l(x) = \frac{c \exp(atx)}{1 - b(\exp(at) - 1)}.$$

Hence the functions $d_l(\hat{p}, \hat{q})$ may be considered as meromorphic functions with respect to $\hat{p} \in \mathbb{C}$, whose values are bounded functions of $\hat{q} \in [0, 1]$. Finally, using the identity $\sum_{n \in \mathbb{Z}} \exp(2n\pi\sqrt{-1}(\tilde{q} - \tilde{q}')) = \delta(\tilde{q} - \tilde{q}')$ we conclude that the following infinite sum of finite

rank operators converges in the strong operator topology to an operator, which is given by an integral kernel with respect to $\hat{q} \in [0, 1]$ depending on $\tilde{q} \in \mathbb{R}/\mathbb{Z}$:

$$\left(\sum_{(n_1, n_2) \in \mathbb{Z}^2} \psi_{n_1 \hat{\kappa} + n_2 \tilde{\kappa}} \frac{|\tilde{\psi}\rangle\langle\tilde{\phi}|}{(\hat{p} - \hat{p}' + n_1)^i} \psi_{-n_1 \hat{\kappa} - n_2 \tilde{\kappa}} \tilde{\chi} \right) (\hat{q}, \tilde{q}) =$$

$$\text{vol}(\mathbb{R}^2/\Lambda) \int_{\hat{q}-1}^{\hat{q}} d_l(\hat{p}, \hat{q} - \hat{q}') \begin{pmatrix} \tilde{\psi}_1(\hat{q}, \tilde{q}) \tilde{\phi}_1(\hat{q}', \tilde{q}) & \tilde{\psi}_1(\hat{q}, \tilde{q}) \tilde{\phi}_2(\hat{q}', \tilde{q}) \\ \tilde{\psi}_2(\hat{q}, \tilde{q}) \tilde{\phi}_1(\hat{q}', \tilde{q}) & \tilde{\psi}_2(\hat{q}, \tilde{q}) \tilde{\phi}_2(\hat{q}', \tilde{q}) \end{pmatrix} \begin{pmatrix} \tilde{\chi}_1(\hat{q}', \tilde{q}) \\ \tilde{\chi}_2(\hat{q}', \tilde{q}) \end{pmatrix} d\hat{q}'.$$

This proves the claim. Thus, due to Hölder's inequality [R-S-I, Theorem III.1 (c)] the function $\tilde{A}_f(\cdot)$ is a meromorphic function from the complex plane $\hat{p} \in \mathbb{C}$ into the bounded operators from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ to $L^{p'}(\mathbb{R}^2/\Lambda) \times L^{p'}(\mathbb{R}^2/\Lambda)$ for all $1 < p' < p < \infty$. By definition this function $\tilde{A}_f(\cdot)$ satisfies condition (i).

In order to show the existence of the commutator of this function with the unbounded-operator valued function $\tilde{D}_{\hat{\gamma}, \tilde{\gamma}}(V, W, \cdot)$ in (ii) we have to ensure that the corresponding operators maps the domain of $\tilde{D}_{\hat{\gamma}, \tilde{\gamma}}(V, W, \cdot)$ (which does not depend on $\hat{p}$) into this domain. With the help of Theorem 2.3 it is quite straightforward to chose appropriate Banach spaces and to show this property. For all $(n_1, n_2) \in \mathbb{Z}^2$ and all $\hat{p} \in \mathbb{C}$ we have

$$\tilde{D}_{\hat{\gamma}, \tilde{\gamma}}(V, W, \hat{p}) \psi_{n_1 \hat{\kappa} + n_2 \tilde{\kappa}} = \psi_{n_1 \hat{\kappa} + n_2 \tilde{\kappa}} \left(\tilde{D}_{\hat{\gamma}, \tilde{\gamma}}(V, W, \hat{p} + n_1) + n_2 \pi \mathbf{1} \right).$$

Hence for all $\hat{p} \in \mathbb{C}$ the commutator is equal to

$$\begin{aligned} [\tilde{A}_f(\hat{p}), \tilde{D}_{\hat{\gamma}, \tilde{\gamma}}(\hat{p})] &= \sum_{(n_1, n_2) \in \mathbb{Z}^2} \left[\psi_{n_1 \hat{\kappa} + n_2 \tilde{\kappa}} \tilde{F}_{\text{singular}}(\hat{p} + n_1) \psi_{-n_1 \hat{\kappa} - n_2 \tilde{\kappa}}, \tilde{D}_{\hat{\gamma}, \tilde{\gamma}}(\hat{p}) \right] \\ &= \sum_{(n_1, n_2) \in \mathbb{Z}^2} \psi_{n_1 \hat{\kappa} + n_2 \tilde{\kappa}} \left[\tilde{F}_{\text{singular}}(\hat{p} + n_1), \tilde{D}_{\hat{\gamma}, \tilde{\gamma}}(\hat{p} + n_1) \right] \psi_{-n_1 \hat{\kappa} - n_2 \tilde{\kappa}}. \end{aligned}$$

The operator-valued functions $\tilde{F}(\cdot)$ and $\tilde{D}_{\hat{\gamma}, \tilde{\gamma}}(\cdot)$ commute pointwise, because the projection $\tilde{P}_{\hat{\gamma}}$ is the spectral projection of $\tilde{D}_{\hat{\gamma}, \tilde{\gamma}}(\cdot)$. If we insert the explicit form of the operator-valued function $\tilde{D}_{\hat{\gamma}, \tilde{\gamma}}(\hat{p})$, then we obtain

$$[\tilde{A}_f(\hat{p}), \tilde{D}_{\hat{\gamma}, \tilde{\gamma}}(\hat{p})] = \pi \sum_{\kappa \in \Lambda^*} \psi_{\kappa} \left[\tilde{F}_{-1}, \begin{pmatrix} \frac{\hat{\kappa}_2 - \sqrt{-1} \hat{\kappa}_1}{\hat{\kappa}_2 - \sqrt{-1} \tilde{\kappa}_1} & 0 \\ 0 & \frac{\hat{\kappa}_2 + \sqrt{-1} \hat{\kappa}_1}{\hat{\kappa}_2 + \sqrt{-1} \tilde{\kappa}_1} \end{pmatrix} \right] \psi_{-\kappa},$$

where $\tilde{F}_{-1}$ is the residue of the operator-valued form $\tilde{F}(\hat{p}) d\hat{p}$ at the pole $\hat{p} = \hat{p}'$. This shows that the commutator does not depend on $\hat{p}$ and that the diagonal elements with respect to the decomposition $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ vanishes. Moreover, since $\sum_{\kappa \in \Lambda^*} \psi_{\kappa}(x)$ is

equal to $\text{vol}(\mathbb{R}^2/\Lambda)$ times Dirac's δ -function $\delta(x)$, for all finite rank operators of the form $\tilde{\chi} \mapsto \langle\langle \tilde{\phi}, \tilde{\chi} \rangle\rangle \tilde{\psi}$ with elements $\tilde{\phi}$ and $\tilde{\psi}$ of $\bigcap_{q < \infty} L^q(\mathbb{R}^2/\Lambda)$, the infinite sum

$$\tilde{\chi} \mapsto \sum_{\kappa \in \Lambda^*} \langle\langle \tilde{\phi}, \psi_{-\kappa} \tilde{\chi} \rangle\rangle \psi_{\kappa} \tilde{\psi}$$

converges in the strong operator topology to the operator $\text{vol}(\mathbb{R}^2/\Lambda)$ times $\begin{pmatrix} \tilde{\psi}_1\tilde{\phi}_1 & \tilde{\psi}_1\tilde{\phi}_2 \\ \tilde{\psi}_2\tilde{\phi}_1 & \tilde{\psi}_2\tilde{\phi}_2 \end{pmatrix}$, where the functions $\tilde{\psi}_1, \tilde{\psi}_2, \tilde{\phi}_1$ and $\tilde{\phi}_2$ are considered as the operators of multiplication with these functions. Hence the commutator is equal to a variation $\delta\tilde{D}_{\hat{\gamma},\tilde{\gamma}}$ associated to unique variations δV_f and δW_f of the potentials. Due to Lemma 3.1 the form $\Omega_{V,W}(\delta V_f, \delta W_f)$ is equal to

$$\Omega_{V,W}(\delta V_f, \delta W_f) = \text{tr} \left(\tilde{P}_{\hat{\gamma}} \delta\tilde{D}_{\hat{\gamma},\tilde{\gamma}} \right) \frac{d\hat{p}}{\pi} = \text{tr} \left(\tilde{P}_{\hat{\gamma}} \left[\tilde{A}_f(\hat{p}), \tilde{D}_{\hat{\gamma},\tilde{\gamma}}(\hat{p}) \right] \right) \frac{d\hat{p}}{\pi}$$

for all $\hat{p} \neq \hat{p}'$. If we choose $\hat{p}$ to be equal to the corresponding value of the point, where $\tilde{P}_{\hat{\gamma}}$ is evaluated, then the projection $\tilde{P}_{\hat{\gamma}}$ commutes with the operator $\tilde{D}_{\hat{\gamma},\tilde{\gamma}}(\hat{p})$ and the trace vanishes. Thus the form vanishes on an open dense subset of the *complex Fermi curve* and therefore on the whole *complex Fermi curve*. This proves (ii).

For all elements $\tilde{\chi} \in L^1(\mathbb{R}/\mathbb{Z})$ the Riemann–Lebesgue Lemma [R-S-II, Theorem IX.7] shows that the estimate

$$\left| \int_0^1 \chi(\hat{q}') \exp(2\pi\sqrt{-1}(\hat{p}' - \hat{p})\hat{q}') d\hat{q}' \right| \leq \exp(2\pi\Im(\hat{p} - \hat{p}')) \cdot |\check{f}(\Re(\hat{p}' - \hat{p}))|$$

holds with some continuous function $\check{f}$ on $\mathbb{R}$, which vanishes at infinity. Due to the form of the functions $d_l(\hat{p}, \hat{q})$, this implies the estimate (iii).

The total residue of the form $f \cdot \Omega_{V,W}$ on $\mathcal{U}$ is equal to

$$\begin{aligned} \text{Res}_{\mathcal{U}}(f \cdot \Omega_{V,W}) &= \text{Res}_{\mathcal{U}} \left(\text{tr} \left(f \cdot \tilde{P}_{\hat{\gamma}} \begin{pmatrix} 0 & \frac{\delta W}{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1} \\ \frac{\delta V}{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1} & 0 \end{pmatrix} \right) \frac{d\hat{p}}{\pi} \right) \\ &= \text{tr} \left(\text{Res}_{\hat{p}=\hat{p}'} \left(\tilde{F}(\hat{p}) \begin{pmatrix} 0 & \frac{\delta W}{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1} \\ \frac{\delta V}{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1} & 0 \end{pmatrix} \frac{d\hat{p}}{\pi} \right) \right) \\ &= \frac{1}{\pi} \text{tr} \left(\tilde{F}_{-1} \begin{pmatrix} 0 & \frac{\delta W}{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1} \\ \frac{\delta V}{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1} & 0 \end{pmatrix} \right). \end{aligned}$$

Since $\tilde{F}_{-1}$ is a finite rank operator, it is an integral operator. The trace of the composition of an integral operator with the operator of multiplication with some function is equal to the integral over the product of the diagonal part of the integral kernel and the function. On the other hand the infinite sum $\sum_{\kappa \in \Lambda^*} \psi_{\kappa} \tilde{F}_{-1} \psi_{-\kappa}$ converges in the strong operator topology to the operator of $\text{vol}(\mathbb{R}^2/\Lambda)$ times the multiplication with the diagonal part of the integral kernel of $\tilde{F}_{-1}$. By definition of δV_f and δW_f we have

$$\left[\sum_{\kappa \in \Lambda^*} \psi_{\kappa} \tilde{F}_{-1} \psi_{-\kappa}, \begin{pmatrix} \frac{\hat{\kappa}_2 - \sqrt{-1}\hat{\kappa}_1}{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1} & 0 \\ 0 & \frac{\hat{\kappa}_2 + \sqrt{-1}\hat{\kappa}_1}{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1} \end{pmatrix} \right] = \frac{1}{\pi} \begin{pmatrix} 0 & \frac{\delta W_f}{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1} \\ \frac{\delta V_f}{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1} & 0 \end{pmatrix}.$$

$$\begin{aligned}
\text{This implies } & \frac{1}{\pi \operatorname{vol}(\mathbb{R}^2/\Lambda)} \sum_{\kappa \in \Lambda^*} \psi_{\kappa} \tilde{F}_{-1} \psi_{-\kappa} = \\
& = \frac{1}{\pi^2 \operatorname{vol}(\mathbb{R}^2/\Lambda)} \left(\begin{array}{c} \cdot \\ \left(\frac{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1}{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1} - \frac{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1}{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1} \right)^{-1} \frac{\delta V_f}{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1} \\ \cdot \end{array} \begin{array}{c} \left(\frac{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1}{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1} - \frac{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1}{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1} \right)^{-1} \frac{\delta W_f}{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1} \\ \cdot \end{array} \right) \\
& = \frac{g(\tilde{\kappa}, \tilde{\kappa})}{2\pi^2 \sqrt{-1}} \left(\begin{array}{c} \cdot \\ \frac{\delta W_f}{\tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1} \\ \frac{-\delta V_f}{\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1} \\ \cdot \end{array} \right),
\end{aligned}$$

and the formula (iv) follows.

q.e.d.

The integral on the left hand side of the formula (iv) is equal to the value of some unique symplectic form on the Hilbert space of all potentials evaluated at the variations $(\delta V, \delta W)$ and $(\delta V_f, \delta W_f)$. Due to (ii) the kernel of the map $f \mapsto (\delta V_f, \delta W_f)$ contains data f entering in first Cousin Problem, whose corresponding operator-valued function $\tilde{A}_f(\cdot)$ commute with the unbounded-operator function $\tilde{D}_{\tilde{\gamma}, \tilde{\gamma}}(V, W, \cdot)$. The eigenvalue of this function $\tilde{A}_f(\cdot)$ yields a global meromorphic function on the *complex Fermi curve*. Due to the definition of the function $\tilde{A}_f(\cdot)$, the singular part of this meromorphic function coincides with the singular part of f on $\mathcal{U}$, and has no other poles. Therefore, this function is a solution of the corresponding first Cousin Problem. More precisely, it can be shown that on the complement of the singular part described in the initial data f this function is holomorphic with respect to the structure sheaf of all eigenvalues of holomorphic functions commuting with the function $\tilde{D}_{\tilde{\gamma}, \tilde{\gamma}}(V, W, \cdot)$. Furthermore, the points $\infty^{\pm}$ are identified to an ordinary double point (compare with [Sch]). In fact property (iii) implies that these functions vanish at $\infty^{\pm}$. If we identify all data f entering in the first Cousin problem with zero, which are the singular parts of global meromorphic functions, then the quotient space is naturally isomorphic to the Lie algebra of the Picard group [Fo, §18.2]. We conclude that the map $f \mapsto (\delta V_f, \delta W_f)$ induces a map from the Lie algebra of the Picard group into the tangent space of the phase space. The pairing on the left hand side of the formula (iv) between the space of regular one-forms and the Lie algebra of the Picard group is given by S erre Duality [Fo, §17]. As a remarkable consequence of the formula (iv) the image of the map $\Omega_{V,W}$ described in Lemma 3.1 is surjective, if the *complex Fermi curve* has finite genus (compare with Proposition 3.7). We expect that a similar statement holds also in the general case (compare with [Sch, Chapter 7]).

Remark 3.3. *The reduction to the fixed points of the involution $(V, W) \mapsto (W, V)$ (compare with Section 2.3) destroys this structure. More precisely, for geometric algebraic solutions (corresponding to complex Fermi curves of finite geometric genus) the number of independent integrals is equal to the geometric genus of the quotient of the complex Fermi curve modulo the holomorphic involution σ . On the other hand the number of degrees of freedom of the isospectral sets (i. e. the set of all real potentials U , whose complex Fermi curves are equal to some given complex Fermi curve) is equal to the dimension of the Prym varieties associated to the complex Fermi curves together with the holomorphic involution σ (see e. g.*

[L-B, Chapter 12]). Due to the Riemann–Hurwitz formula the difference of these numbers is proportional to the number of fixed points of the holomorphic involution. Any bound on this difference would yield a bound on the geometric genus of spectral curves corresponding to minimal tori in S^3 .

Let us now investigate the compatibility of the involutions and the corresponding relations with the map $f \mapsto (\delta V_f, \delta W_f)$. The operator $\begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix}$ is self-adjoint and satisfies the relation

$$J \begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix} + \begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix}^t J = 0.$$

Hence the operator $\tilde{D}_{\hat{\gamma}, \hat{\gamma}}(\hat{p})$ satisfies the relations

$$\begin{aligned} -\tilde{D}_{\hat{\gamma}, \hat{\gamma}}^t(-\hat{p}) &= - \begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix}^{-1} J^{-1} \tilde{D}_{\hat{\gamma}, \hat{\gamma}}(\hat{p}) \begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix}^t J \\ &= \begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix}^{-1} J^{-1} \tilde{D}_{\hat{\gamma}, \hat{\gamma}}(\hat{p}) J \begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix} \\ &= \begin{pmatrix} -\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 & 0 \\ 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \end{pmatrix}^{-1} \tilde{D}_{\hat{\gamma}, \hat{\gamma}}(\hat{p}) \begin{pmatrix} -\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 & 0 \\ 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \end{pmatrix}, \\ \tilde{D}_{\hat{\gamma}, \hat{\gamma}}^*(\bar{\hat{p}}) &= \begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix}^{-1} \tilde{D}_{\hat{\gamma}, \hat{\gamma}}(\hat{p}) \begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix}, \\ -\tilde{\bar{D}}_{\hat{\gamma}, \hat{\gamma}}(-\bar{\hat{p}}) &= J^{-1} \tilde{D}_{\hat{\gamma}, \hat{\gamma}}(\hat{p}) J. \end{aligned}$$

Therefore, the projection $\tilde{P}_{\hat{\gamma}}$ satisfies the relations

$$\begin{aligned} \tilde{P}_{\hat{\gamma}} &= \begin{pmatrix} -\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 & 0 \\ 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \end{pmatrix} \sigma^* \left(\tilde{P}_{\hat{\gamma}}^t \right) \begin{pmatrix} -\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 & 0 \\ 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \end{pmatrix}^{-1} \\ &= \begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix} \rho^* \left(\tilde{P}_{\hat{\gamma}}^* \right) \begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix}^{-1} \\ &= J \eta^* \left(\tilde{\bar{P}}_{\hat{\gamma}} \right) J^{-1}. \end{aligned}$$

The following lemma is verified by some easy calculations.

Lemma 3.4. *The map $f \mapsto \tilde{A}_f$ has the following properties:*

- (i) *If the pair of potentials is of the form $(U, \bar{U})$ and f is a meromorphic function with finitely many poles on some open subset $\mathcal{U}$ of the complex Fermi curve, then $\sigma^* f$ is a meromorphic function with finitely many poles on $\sigma(\mathcal{U})$. The associated operator-valued functions satisfy for all $\hat{p} \in \mathbb{C}$:*

$$\begin{pmatrix} -\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 & 0 \\ 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \end{pmatrix} \tilde{A}_{\sigma^* f}^t(-\hat{p}) \begin{pmatrix} -\tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 & 0 \\ 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \end{pmatrix}^{-1} = \tilde{A}_f(\hat{p}),$$

and the associated variation $(\delta V_{\sigma^ f}, \delta W_{\sigma^* f})$ is equal to $(-\delta W_f, -\delta V_f)$.*

- (ii) *If the potentials are real and f is a meromorphic function with finitely many poles on some open subset $\mathcal{U}$ of the complex Fermi curve, then $\rho^* \bar{f}$ is a meromorphic function*

with finitely many poles on $\rho(\mathcal{U})$. For all $\hat{p} \in \mathbb{C}$ the associated operator-valued functions satisfy

$$\begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix} \tilde{\mathbf{A}}_{\rho^* \bar{f}}^*(\hat{p}) \begin{pmatrix} 0 & \tilde{\kappa}_2 + \sqrt{-1}\tilde{\kappa}_1 \\ \tilde{\kappa}_2 - \sqrt{-1}\tilde{\kappa}_1 & 0 \end{pmatrix}^{-1} = \tilde{\mathbf{A}}_f(\hat{p}).$$

The associated variation $(\delta V_{\rho^* \bar{f}}, \delta W_{\rho^* \bar{f}})$ is equal to $(-\delta \bar{V}_f, -\delta \bar{W}_f)$.

- (iii) If the pair of potentials is of the form $(U, \bar{U})$ and f is a meromorphic function with finitely many poles on some open subset $\mathcal{U}$ of the complex Fermi curve, then $\eta^* \bar{f}$ is a meromorphic function with finitely many poles on $\eta(\mathcal{U})$. The associated operator-valued functions satisfy for all $\hat{p} \in \mathbb{C}$:

$$\mathbf{J} \tilde{\mathbf{A}}_{\eta^* \bar{f}}^*(-\bar{\hat{p}}) \mathbf{J}^{-1} = \tilde{\mathbf{A}}_f(\hat{p}),$$

and the associated variation $(\delta V_{\eta^* \bar{f}}, \delta W_{\eta^* \bar{f}})$ is equal to $(\delta \bar{W}_f, \delta \bar{V}_f)$

- (iv) The constant operator $\tilde{\mathbf{A}}_f(\hat{p}) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ fulfills condition (i)–(ii) of Lemma 3.2. The corresponding variation is equal to $\delta V_f = -2V$ and $\delta W_f = 2W$. **q.e.d.**

The corresponding function f in (iv) would be a function, which is in some neighbourhood of ∞^- equal to 1 and on some neighbourhood of ∞^+ equal to -1 . Since the two infinities should be considered as one double point, this function is not holomorphic (compare with [Sch]). Moreover, the forms $\Omega_{V,W}(\delta V, \delta W)$ have in general two simple poles at the two infinities, and the residue is proportional to the variation of the functional $\int VW d^2x$.

The foregoing Lemmata have an application, which will be an essential tool in the classification of relative minimizers. By the residue formula $\sum_{\text{poles of } f\omega} \text{Res}(f\omega) = 0$ on a compact Riemann surface every meromorphic function f induces a relation of the values of a holomorphic (meromorphic) form ω at the poles of f (and the poles of ω). Due to S erre duality [Fo, §17.] all relations on the values of all holomorphic forms are of this form. We shall see now that the same is true for general complex Fermi curves.

Proposition 3.5. *Let $\mathcal{F}(V, W)$ be a complex Fermi curve with the property that for all variations δV and δW , whose regular forms $\Omega_{V,W}(\delta V, \delta W)$ vanish at finitely many points $y_1, \dots, y_L \in \mathcal{F}(V, W)/\Lambda^*$, the corresponding variation $4 \int_{\mathbb{R}^2/\Lambda} (V(x)\delta W(x) + \delta V(x)W(x)) d^2x$ of the first integral vanishes. Then this complex Fermi curve $\mathcal{F}(V, W)/\Lambda^*$ can be compactified to a compact Riemann surface by adding the points $\infty^\pm$ (compare with condition Finite type potentials (i)). Moreover, this compactified complex Fermi curve has a global meromorphic function whose poles are contained in $\{y_1, \dots, y_L\}$ and which takes the values ± 1 at $\infty^\pm$.*

Proof. The form $\Omega_{V,W}(\delta V, \delta W)$ vanishes at y_l , if and only if the residue of this form multiplied with a meromorphic function f_l on a neighbourhood $\mathcal{U}$ of y_l vanishes, where f has a first order pole at y_l and no other poles. Therefore, for all variations $(\delta V, \delta W)$, whose residues

$$\text{Res}_{y_1}(f_1 \Omega_{V,W}(\delta V, \delta W)), \dots, \text{Res}_{y_L}(f_L \Omega_{V,W}(\delta V, \delta W))$$

vanish, the variation of the *first integral* has to vanish. Due to Lemma 3.2 (iv) the former residues vanish, if and only if the symplectic form of $(\delta V, \delta W)$ and the variations

$$(\delta V_{f_1}, \delta W_{f_1}), \dots, (\delta V_{f_L}, \delta W_{f_L})$$

vanish. Since the variation of the *first integral* is equal to four times the symplectic form of $(V, -W)$ and $(\delta V, \delta W)$, we conclude that $(-V, W)$ is a complex linear combination of $(V_{f_1}, W_{f_1}), \dots, (V_{f_L}, W_{f_L})$. Due to Lemma 3.2 (ii) this implies that the sum of the operator mentioned in Lemma 3.4 (iv) and a complex linear combination of the operators

$$\tilde{A}_{f_1}(\cdot), \dots, \tilde{A}_{f_L}(\cdot)$$

commutes with $\tilde{D}_{\tilde{\gamma}, \tilde{\gamma}}(\cdot)$. The eigenvalues of this family of operators commuting with $\tilde{D}_{\tilde{\gamma}, \tilde{\gamma}}(\cdot)$ defines a meromorphic function on $\mathcal{F}(V, W)$. Due to Lemma 3.2 (i) this function induces a meromorphic function on $\mathcal{F}(V, W)/\Lambda^*$. By construction this function solves the first Cousin Problem (see e. g. [Ah, Chapter 5 2.1.] and [G-P-R, Chapter III. §3.3.]) of the data given by the corresponding complex linear combination of $f_1, \dots, f_L$. Moreover, due to Theorem 2.12 Lemma 3.2 (iii) and Lemma 3.4 (iv) this function takes values in $\{z \in \mathbb{C} \mid |z + 1| < \varepsilon\}$ on any open set of the form $\mathcal{V}_{\varepsilon, \delta}^-$ (ε arbitrary) and values in $\{z \in \mathbb{C} \mid |z - 1| < \varepsilon\}$ on any set of the form $\mathcal{V}_{\varepsilon, \delta}^+$ (ε arbitrary). If $\varepsilon < 1$, then the normalizations of the corresponding handles described in Theorem 2.12 have two connected components. Therefore, the addition of the points $\infty^\pm$ compactifies the *complex Fermi curve* $\mathcal{F}(V, W)/\Lambda^*$ to a compact Riemann surface of finite genus. The solution of the first Cousin problem yields a meromorphic function on this compact Riemann surface whose poles are contained in $\{y_1, \dots, y_l\}$ and which takes at $\infty^\pm$ the values ± 1 . **q.e.d.**

Remark 3.6. *The proof carries over to more general situations, where the vanishing of $\Omega_{V,W}(\delta V, \delta W)$ at $y_1, \dots, y_L$ with orders $m_1 - 1, \dots, m_L - 1$ implies that the corresponding variation of the first integral vanishes. Furthermore, due to the Separating hyperplane theorem [R-S-I, Theorem V.4], the conclusion of the proposition holds as well, if for variations $(\delta V, \delta W)$, whose forms $\Omega_{V,W}(\delta V, \delta W)$ takes values in a convex sub-cone of $\mathbb{C}$ at the points $y_1, \dots, y_L$, the corresponding variation of the first integral takes values in the complement of that convex sub-cone. Finally, by construction of the variations $(\delta V_f, \delta W_f)$ in the proof of Lemma 3.2, the conclusion of the proof of this proposition $(-V, W)$ being a linear combination of the variations $(\delta V_{f_1}, \delta W_{f_1}), \dots, (\delta V_{f_L}, \delta W_{f_L})$ implies that these potentials are smooth. In fact, due to Remark 2.7 the resolvent of the free Dirac operator is a bounded operator from the Sobolev spaces $W^{m-1,p}(\mathbb{R}^2/\Lambda) \times W^{m-1,p}(\mathbb{R}^2/\Lambda)$ onto $W^{m,p}(\mathbb{R}^2/\Lambda) \times W^{m,p}(\mathbb{R}^2/\Lambda)$ for all $m \in \mathbb{N}$ and $1 < p < \infty$. Hence, due to the proof of Theorem 2.3, for all $m \in \mathbb{N}$ the eigenfunctions of potentials in the Sobolev spaces $\bigcap_{q < \infty} W^{m-1,q}(\mathbb{R}^2/\Lambda)$ belong to the Sobolev spaces $\bigcap_{q < \infty} W^{m,q}(\mathbb{R}^2/\Lambda)$. We conclude inductively that the potential of a critical point is indeed smooth.*

If the pair of potentials is of the form $(U, \bar{U})$, then due to Lemma 2.26 U has to be of finite type. Consequently, due to the obvious generalization of Lemma 2.27 to compact one-dimensional complex spaces Y (instead of compact smooth Riemann surfaces) U is actually analytic.

We shall see that this proposition applies to critical points of the Willmore functional, and yields a function theoretic characterization of these critical points. The corresponding global meromorphic function induces a holomorphic mapping from the spectral curve onto $\mathbb{P}^1$. It is well known [D-K-N] that if the preimage of one element of $\mathbb{P}^1$ is contained in $\{\infty^-, \infty^+\}$, then the corresponding potentials solve a differential equation, and belong to a reduction of the integrable system. The known descriptions of Willmore surfaces [Ba, B-B, Bo] and of conformal minimal immersion into S^3 [Hi, P-S] are exactly of this form.

3.2 Deformations of *complex Fermi curves*

In this section we shall prove the smoothness of some subsets of the moduli space. This result will become the essential tool in the investigation of relative minimizers of the transformed variational problem of the Willmore functional. Due to condition *Quasi-momenta* (i) the complex space $Y \setminus \{\infty^-, \infty^+\}$ is a one-sheeted covering of $\mathcal{F}(Y, \infty^-, \infty^+, k)$. The elements of these subsets are locally isomorphic to $\mathcal{F}(Y, \infty^-, \infty^+, k)$ and therefore local complete intersections of $\mathbb{C}^2$ [G-P-R, Chapter VI. §2.1.] with fixed arithmetic genus. To make this precise we introduce another condition on the data $(Y, \infty^-, \infty^+, k)$.

Quasi-momenta (v) There exists an open covering of $Y \setminus \{\infty^-, \infty^+\}$ with the property that the restriction of any branch of k to any neighbourhood of this covering is an isomorphism of this neighbourhood onto some one-dimensional complex subspace of $\mathbb{C}^2$.

Let $\mathcal{M}_{g,\Lambda}$ be the set of all data fulfilling conditions *Quasi-momenta* (i)–(ii) and *Quasi-momenta* (v) with the property, that the arithmetic genus of Y is equal to g . For all data $(Y', \infty^-, \infty^+, k)$, where Y' are smooth compact Riemann surfaces, there exist some data $(Y, \infty^-, \infty^+, k)$ in some $\mathcal{M}_{g,\Lambda}$, where the normalization of Y is equal to Y' and the pullback of the function k on Y under the normalization map is equal to the corresponding function of k on Y' . Hence the union $\bigcup_{g \in \mathbb{N}_0} \mathcal{M}_{g,\Lambda}$ contains all *complex Fermi curves* of finite *geometric genus*.

Proposition 3.7. *The set $\mathcal{M}_{g,\Lambda}$ is a $g+1$ -dimensional complex manifold. For all elements $(Y, \infty^-, \infty^+, k)$ the tangent space of this manifold at this point is isomorphic to the space of all regular forms of the complex space $\tilde{Y}$, which is obtained from Y by identifying ∞^- and ∞^+ to an ordinary double point. More precisely, for any such regular form ω there exists a one-dimensional complex family of data, such that $\omega = \check{p}d\hat{p}$.*

Remark 3.8. *Here we assume that Y is connected. More precisely, we assume that if Y has a disconnected normalization, then $\tilde{Y}$ has a singularity containing elements of both components. Otherwise the regular forms of $\tilde{Y}$ coincide with the regular forms of Y , and $\mathcal{M}_{g,\Lambda}$ is a g -dimensional manifold. Moreover, in this case the first integral is locally constant on this manifold.*

Here we have used the notation introduced in *Fundamental domain* 2.1. Furthermore, we make the convention that the function $\hat{p}$ does not depend on the deformation.

Proof. For given data $(Y, \infty^-, \infty^+, k) \in \mathcal{M}_{g,\Lambda}$ we shall first construct an open neighbourhood in some complex manifold, whose elements correspond to deformations of Y . More precisely, we consider all deformations of compact one-dimensional complex spaces of arithmetic genus g , on which the multi-valued function $\hat{p} = g(\hat{\gamma}, k)$ exists with the analogous properties *Quasi-momenta* (i)–(ii), and which is a local complete intersection in $\mathbb{C}^2$ [G-P-R, Chapter VI. §2.1.]. Here we carry over the methods of [Ko] to the case of complex spaces instead of complex manifolds. The tools are essentially developed by Grauert (see e. g. [Gr, Ba-St, Gr-Re, G-K, G-P-R]). Afterwards we shall extend any regular form of $\tilde{Y}$ to a holomorphic family of regular forms on the corresponding family of complex spaces. Then we shall show that each of these holomorphic families of regular forms defines some flow on the manifold, which preserves the subset corresponding to elements of $\mathcal{M}_{g,\Lambda}$. Finally, we show that the regular forms on the family of deformed complex spaces may be considered as a holomorphic sub-bundle of the tangent bundle of the manifold of deformations constructed before, and that this sub-bundle is the tangent bundle along the submanifold corresponding to elements of $\mathcal{M}_{g,\Lambda}$. Along these lines the proof is achieved in ten steps.

1. Since the arithmetic genus is finite, the union of the singular points of Y and the zeroes of $d\hat{p}$ is a finite subset of Y . Now we choose a finite covering $\mathcal{U}_0, \dots, \mathcal{U}_L$ of Y , such that $\mathcal{U}_1, \dots, \mathcal{U}_L$ are disjoint open neighbourhoods, each of which contains either one singular point or one zero of $d\hat{p}$. Without loss of generality we may assume that the function $\hat{p}$ maps all these open subsets onto Weierstraß coverings over open subsets $\mathcal{V}_1, \dots, \mathcal{V}_L$ of $\hat{p} \in \mathbb{C}$. We in addition assume that the subset $\mathcal{U}_0$ is contained in the complement of some neighbourhood of the union of all singular points and all zeroes of $d\hat{p}$. Finally, the two marked points ∞^- and ∞^+ are assumed to be elements of the complement of the closure of $\mathcal{U}_1 \cup \dots \cup \mathcal{U}_L$ in $\mathcal{U}_0$. We conclude that the restriction of the function $\hat{p}$ to $\mathcal{U}_0$ is locally biholomorphic, and therefore $\mathcal{U}_0$ may be identified with some union of open subsets of $\hat{p} \in \mathbb{P}^1$. Due to condition *Quasi-momenta* (v) any $\mathcal{U}_l$ of the subsets $\mathcal{U}_1, \dots, \mathcal{U}_L$ is isomorphic to the complex spaces defined by some equation $R(\hat{p}, \check{p}) = 0$ over some open subset $\mathcal{V}_l$ of $\mathbb{C}^2$. Moreover, we may choose this function $R(\hat{p}, \check{p})$ to be a polynomial with respect to $\check{p}$, whose degree d is equal to the number of sheets of this Weierstraß covering with respect to the function $\hat{p}$, and whose highest coefficient is 1. It would be more accurate to decorate the functions R and their degrees d with an index $l = 1, \dots, L$. We omit these indices, and hope that from the context it will be clear, which objects are associated to which member of the covering.

2. Let us simplify these local descriptions to polynomial equations $Q(\hat{p}, \check{p}) = 0$ together with an equation $\check{p} = P(\hat{p}, \check{p})$, which is polynomial with respect to $\check{p}$ of degree $d - 1$, and whose coefficients are holomorphic functions depending on $\hat{p} \in \mathcal{V}_l$. If we choose the neighbourhoods $\mathcal{U}_1, \dots, \mathcal{U}_L$ small enough, then there always exists such a simplification, where the coefficients of $Q(\hat{p}, \check{p})$ are equal to the corresponding lowest non-vanishing Taylor coefficients of $R(\hat{p}, \check{p})$. More precisely, if the preimage of the corresponding zero of $d\hat{p}$ in the normalization is a single point, then $Q(\hat{p}, \check{p})$ can be chosen to be of the form $\check{p}^d - \hat{p}^m$, where $c\hat{p}^m$ is the lowest non-vanishing coefficient of R of this form. In this case d and m have to be co-prime, and we define the degree of $\hat{p}$ as d and the degree of $\check{p}$ (and $\check{p}$) as m . In general the *Local contribution to the arithmetic genus* 4.3 of the equation $R(\hat{p}, \check{p}) = 0$ at the corresponding zero of $d\hat{p}$ has to be the same as the corresponding *Local contribution to the arithmetic genus* 4.3 of the

equation $Q(\hat{p}, \check{p}) = 0$.

3. If $\dot{Q}(\hat{p}, \check{p})$ is any polynomial with respect to $\hat{p}$ and $\check{p}$, whose degree with respect to $\check{p}$ is smaller than d , the equation $Q(\hat{p}, \check{p}, t) = Q(\hat{p}, \check{p}) + t\dot{Q}(\hat{p}, \check{p})$ defines a deformation of the complex space $\mathcal{U}_l$. Also for small $|t|$ the function $\hat{p}$ of this deformation defines a Weierstraß covering over $\mathcal{V}_l$, whose restriction to the corresponding deformed sets $\mathcal{U}_l \cap \mathcal{U}_0$ is biholomorphic to the undeformed set. Hence this deformation defines a deformation of the entire compact complex space. If we consider $\check{p}$ as a function depending on $\hat{p}$ and t , the derivative $\partial\check{p}/\partial t$ is equal to $-\dot{Q}(\hat{p}, \check{p})/(\partial Q(\hat{p}, \check{p}, t)/\partial\check{p})$. If for small $|t|$ this quotient is a holomorphic function on the deformed space, then we conclude that at least for small values of $|t|$ the deformed function $\check{p}$ may be expressed by the undeformed functions $\hat{p}$ and $\check{p}$ and vice versa. Therefore, the family of complex spaces are all isomorphic. Consequently we choose finitely many monomials $Q_1, \dots, Q_r$ with respect to $\hat{p}$ and $\check{p}$, whose degree with respect to $\check{p}$ is smaller than d , and which induce a basis of the space of global sections of the sheaf $\mathcal{O}/\mathcal{O}(\partial Q(\hat{p}, \check{p})/\partial\check{p})$ on the subset $\mathcal{U}_l$. Let $\tilde{\mathcal{M}}_l \subset \mathbb{C}^r$ be some small open neighbourhood of 0, and let Z_l be the complex subspace of $(\hat{p}, \check{p}, t_1, \dots, t_r) \in \mathcal{V}_l \times \mathbb{C} \times \tilde{\mathcal{M}}_l$ defined by the equation

$$Q(\hat{p}, \check{p}, t_1, \dots, t_r) = Q(\hat{p}, \check{p}) + t_1 Q_1(\hat{p}, \check{p}) + \dots + t_r Q_r(\hat{p}, \check{p}) = 0.$$

There exists a natural map $f_l : Z_l \rightarrow \tilde{\mathcal{M}}_l$, which is induced by the projection $\mathcal{V}_l \times \mathbb{C} \times \tilde{\mathcal{M}}_l \rightarrow \tilde{\mathcal{M}}_l$. For all elements $t_1, \dots, t_r \in \tilde{\mathcal{M}}_l$ the complex subspaces defined by the pullback under f_l under the unique maximal ideal corresponding to $t_1, \dots, t_r$ is obviously the deformation of $\mathcal{U}_l$ defined by the equation $Q(\hat{p}, \check{p}, t_1, \dots, t_r) = 0$. Hence Z_l is some “universal family” of deformations of $\mathcal{U}_l$. Now we claim that the map f_l is flat. In fact, f_l is the composition of some Weierstraß map $Z_l \rightarrow \mathcal{V}_l \times \tilde{\mathcal{M}}_l$ and the projection $\mathcal{V}_l \times \tilde{\mathcal{M}}_l \rightarrow \tilde{\mathcal{M}}_l$. The first map is flat due to [G-P-R, Chapter II Proposition 2.10] and the Weierstraß Isomorphism [Gr-Re, Chapter 2. §4.2.]. The second map is flat due to [G-P-R, Chapter II Corollary 2.7]. Hence [G-P-R, Chapter II Proposition 2.6] shows the claim. If $\tilde{\mathcal{M}}_l$ is chosen small enough, the deformations of the subsets $\mathcal{U}_l \cap \mathcal{U}_0$ do not contain any zero of the function $\partial Q(\hat{p}, \check{p}, t_1, \dots, t_r)/\partial\check{p}$. In this case we may glue Z_l with the complex space $\mathcal{U}_0 \times \tilde{\mathcal{M}}_l$ along the deformations of the subset $\mathcal{U}_l \cap \mathcal{U}_0$. If we repeat this for all $l = 1, \dots, L$ we obtain a universal family $f : Z \rightarrow \tilde{\mathcal{M}} = \tilde{\mathcal{M}}_1 \times \dots \times \tilde{\mathcal{M}}_L$, of deformations of the compact complex space Y . We conclude that the map f is proper and flat.

4. Now we shall prove that for all $l = 1, \dots, L$ the corresponding polynomials $Q_1, \dots, Q_r$ form a basis of all sections of the sheaf $\mathcal{O}/\mathcal{O}(\partial Q(\hat{p}, \check{p}, t_1, \dots, t_r)/\partial\check{p})$ on Z_l . The support of this sheaf is the complex subspace $\tilde{Z}_l$ of Z_l defined by the equation $\partial Q(\hat{p}, \check{p}, t_1, \dots, t_r)/\partial\check{p} = 0$. The union of these subspaces is a subspace $\tilde{Z}$ of Z , and the map f induces a map $\tilde{f} : \tilde{Z} \rightarrow \tilde{\mathcal{M}}$. This map is again proper and flat. In fact, it is a finite map, and the dimension of the stalk of the preimage of any point of $\tilde{\mathcal{M}}$ considered as a $\mathbb{C}$ -vector space is equal to the number of zeroes of the form $d\hat{p}$ on the corresponding deformation of Y . Due to the Semi-Continuity theorem [G-P-R, Chapter III Theorem 4.7] the arithmetic genus of all these deformations is equal to g . Moreover, since all deformations are local complete intersections in $\mathbb{C}^2$ [G-P-R, Chapter VI. §2.1.], Lemma 2.20 shows that the sheaf of regular forms are locally free. Therefore, the degree of the canonical divisor is equal to $2g - 2$ [Se, Chapter IV. §3.11]. Since $d\hat{p}$ has two poles at the non-singular points ∞^- and ∞^+ , the number of zeroes of $d\hat{p}$ is equal to $2g + 2$

and therefore constant. Again [G-P-R, Chapter II Proposition 2.10] shows that $\check{f}$ is flat. Now the Semi-Continuity theorem [G-P-R, Chapter III Theorem 4.7] implies that the direct image sheaf of the structure sheaf of $\check{Z}$ under $\check{f}$ defines some holomorphic vector bundles on $\tilde{\mathcal{M}}$ of rank $2g + 2$. Hence for any choice of sections, which form a basis in the fibre over $0 \in \tilde{\mathcal{M}}$, these sections form a basis of all fibres in some neighbourhood. If $\tilde{\mathcal{M}}_l$ is chosen small enough, then the corresponding elements $Q_1, \dots, Q_r$ form a basis of all global sections of the sheaf $\mathcal{O}/\mathcal{O}(\partial Q(\hat{p}, \check{p}, t_1, \dots, t_r)/\partial \check{p})$ on Z_l .

5. Due to Lemma 2.20 the sheaf of regular forms on the family of deformations of Y is locally free. Therefore, [G-P-R, Chapter II. Proposition 2.6] implies that this sheaf is also f -flat, and the Semi-Continuity theorem [G-P-R, Chapter III Theorem 4.7] shows that the direct image of the sheaf of sections of the regular forms of the deformations of Y under f defines some vector bundle on $\tilde{\mathcal{M}}$ of rank g . The same argumentation carries over to the direct image of the sheaf of regular forms of the corresponding deformations of $\check{Y}$, which is defined by identifying the non-singular points ∞^- and ∞^+ to an ordinary double point. If Y is connected (i. e. either the normalization of Y is connected or the two connected components of the normalization of Y are connected by some singularity), then this yields a holomorphic vector bundle of rank $g + 1$ on $\tilde{\mathcal{M}}$, otherwise a holomorphic vector bundle of rank g (compare with Remark 3.8). We conclude that any basis of regular forms of $\check{Y}$, extends to a holomorphic family of regular forms on the preimage of some open neighbourhood of $0 \in \tilde{\mathcal{M}}$.

6. If for the deformations of the sets $\mathcal{U}_l$ we make the ansatz $\check{p} = P(\hat{p}, \check{p})$ and conceive the functions $\hat{p}$ and $\check{p}$ as functions depending on $\hat{p}$ and the moduli, then the differential equation $\dot{\hat{p}}d\hat{p} = \omega$, where ω is a section of the vector bundle on $\tilde{M}$ constructed above, transforms into the differential equation

$$\begin{aligned} \frac{\omega}{\partial P(\hat{p}, \check{p})/\partial \check{p}} &= \frac{\dot{\hat{p}}d\hat{p}}{\partial P(\hat{p}, \check{p})/\partial \check{p}} = \left(\frac{\dot{P}(\hat{p}, \check{p})}{\partial P(\hat{p}, \check{p})/\partial \check{p}} + \dot{\check{p}} \right) d\hat{p} \\ &= \left(\frac{\dot{P}(\hat{p}, \check{p})}{\partial P(\hat{p}, \check{p})/\partial \check{p}} - \frac{\dot{Q}(\hat{p}, \check{p})}{\partial Q(\hat{p}, \check{p})/\partial \check{p}} \right) d\hat{p}. \end{aligned}$$

The partial derivative $\partial P(\hat{p}, \check{p})/\partial \check{p}$ of the initial $P(\hat{p}, \check{p})$ has a holomorphic inverse on $\mathcal{U}_l$. By reduction of $\tilde{\mathcal{M}}$ we may achieve that the same is true for all deformed $P(\hat{p}, \check{p})$. Due to Lemma 2.20 any regular form on some deformed Y of the family constructed above may be written on each $\mathcal{U}_l$ of the deformed subsets $\mathcal{U}_1, \dots, \mathcal{U}_L$ as the quotient of some holomorphic function $f(\hat{p}, \check{p})$ divided by $\partial Q(\hat{p}, \check{p}, t_1, \dots, t_r)/\partial \check{p}$. Therefore, the same is true for $\frac{\omega}{\partial P(\hat{p}, \check{p})/\partial \check{p}}$. Since the polynomials $Q_1, \dots, Q_r$ forms a basis of the sheaf $\mathcal{O}/\mathcal{O}(\partial Q(\hat{p}, \check{p}, t_1, \dots, t_r)/\partial \check{p})$, any such regular form may be written locally as

$$\frac{\omega}{\partial P(\hat{p}, \check{p})/\partial \check{p}} = \frac{c_1 Q_1(\hat{p}, \check{p}) + \dots + c_r Q_r(\hat{p}, \check{p})}{\partial Q(\hat{p}, \check{p}, t_1, \dots, t_r)/\partial \check{p}} d\hat{p} + g(\hat{p}, \check{p}) d\hat{p},$$

where $c_1, \dots, c_r$ are uniquely defined complex numbers (depending holomorphically on the moduli in $\tilde{\mathcal{M}}$) and $g(\hat{p}, \check{p})$ is a unique holomorphic function on $\mathcal{U}_l$ (also depending holomor-

phically on the moduli). Hence we obtain the differential equation

$$\dot{t}_1 = -c_1, \dots, \dot{t}_r = -c_r \text{ and } \dot{P}(\hat{p}, \check{p}) = g(\hat{p}, \check{p}) \frac{\partial P(\hat{p}, \check{p})}{\partial \check{p}}.$$

7. With the help of the Weierstraß Isomorphism [Gr-Re, Chapter 2. §4.2.] the holomorphic functions $P(\hat{p}, \check{p})$, $f(\hat{p}, \check{p})$ and $g(\hat{p}, \check{p})$ may be uniquely represented as polynomials with respect to $\check{p}$ of degree $d-1$ (where d denotes the number of sheets of the Weierstraß covering $\mathcal{U}_l \rightarrow \mathcal{V}_l$), whose coefficients are holomorphic functions depending on $\hat{p}$. It is quite easy to see that with this choice the holomorphic function $\frac{f(\hat{p}, \check{p})}{\partial P(\hat{p}, \check{p})/\partial \check{p}}$ is also of this form, and the coefficients of this polynomial of degree $d-1$ with respect to $\check{p}$ are rational functions of the corresponding coefficient of $P(\hat{p}, \check{p})$ and $f(\hat{p}, \check{p})$. We conclude that the numbers $c_1, \dots, c_r$ and the coefficient of the polynomial $g(\hat{p}, \check{p})$ with respect to $\check{p}$ depend holomorphically on the coefficients of the polynomials $P(\hat{p}, \check{p})$ and $f(\hat{p}, \check{p})$ with respect to $\check{p}$. On $\mathcal{U}_0$ we have the ordinary differential equation $\dot{P}(\hat{p}) = \check{p} = \omega/d\hat{p}$. We conclude that the section ω defines a holomorphic vector field on the product of $\tilde{M}$ with the space of polynomials $P(\hat{p}, \check{p})$ on $\mathcal{U}_0 \times \dots \times \mathcal{U}_L$ of the corresponding degrees $d-1$ with respect to $\check{p}$, whose coefficients depend holomorphically on $\hat{p} \in \mathcal{V}_0 \times \dots \times \mathcal{V}_L$. Since these vector fields are holomorphic they may be integrated to some holomorphic flow at least in some neighbourhood of $0 \in \tilde{\mathcal{M}}$ [La, Chapter IV §1.], which have to be chosen small enough in order to guarantee the assumption that $\partial P(\hat{p}, \check{p})/\partial \check{p}$ has a holomorphic inverse on $\mathcal{U}_l$ for $l = 1, \dots, L$.

8. This implies that in some neighbourhood of $0 \in \tilde{\mathcal{M}}$ on the whole family of the integrated flow two functions $\hat{p}$ and $\check{p} + P(\hat{p}, \check{p})$ are defined, which may be combined to a function k with $\hat{p} = g(\hat{\gamma}, k)$ and $\check{p} = g(\check{\gamma}, k)$. This function k fulfills conditions *Quasi-momenta* (i)–(ii). Moreover, due to the assumption that $\partial P(\hat{p}, \check{p})/\partial \check{p}$ has a holomorphic inverse on $\mathcal{U}_l$, the map $(\hat{p}, \check{p}) \mapsto (\hat{p}, \check{p})$ is locally a biholomorphic map. Therefore, also condition *Quasi-momenta* (v) is fulfilled for small moduli. This shows that any holomorphic family of regular forms on the family of deformations of Y constructed above define a vector field on the manifold of moduli, whose flow preserves the subspace of deformations contained in $\mathcal{M}_{g,\Lambda}$.

9. Now we shall prove that the subset of $\tilde{\mathcal{M}}$, on whose elements besides the function $\hat{p}$ there exists another function $\check{p}$ such that the resulting function k fulfills conditions *Quasi-momenta* (i)–(ii), is a complex subspace of this complex manifold. It is part of the Krichever construction [Kr-1] that the space translations of these soliton equations are induced by the action of some linear subgroup of the Picard group. From this point of view the existence of the function k with the properties *Quasi-momenta* (i)–(ii) is equivalent to the existence of two periods of a natural two-dimensional subgroup of the Picard group associated to the choice of the two marked points ∞^- and ∞^+ . This periodicity is equivalent to the assumption that the two line bundles corresponding to the translation by $\hat{\gamma}$ and $\check{\gamma}$ are trivial. Since the function $\hat{p}$ exists on the whole universal family $\tilde{Z}$, the line bundle corresponding to the translation by $\hat{\gamma}$ is trivial for the whole family. Now we construct the line bundle corresponding to the shift by $\check{\gamma}$. The corresponding global sections are of the form $c \exp(2\pi\sqrt{-1}\check{p})$, where c is some complex number and $\check{p} = g(\check{\gamma}, k)$ together with $\hat{p} = g(\hat{\gamma}, k)$ defines a holomorphic function k on Y fulfilling conditions *Quasi-momenta* (i)–(ii). Let $\mathcal{U}^+$ and $\mathcal{U}^-$ denote small open disjoint neighbourhoods of ∞^- and ∞^+ , which are contained in the complement of $\mathcal{U}_1, \dots, \mathcal{U}_L$. Due

to condition *Quasi-momenta* (i) some branch of the function $\check{p}$ is proportional to $\frac{\check{\gamma}_1 - \sqrt{-1}\check{\gamma}_2}{\check{\gamma}_1 + \sqrt{-1}\check{\gamma}_2}$ times $\hat{p}$ at ∞^- and proportional to $\frac{\check{\gamma}_1 + \sqrt{-1}\check{\gamma}_2}{\check{\gamma}_1 - \sqrt{-1}\check{\gamma}_2}$ times $\hat{p}$ at ∞^+ up to order $\mathbf{O}(1/\hat{p})$. Hence the line bundle, which describes the lifting of the translation by $\check{\gamma}$, may be defined by the trivial bundle on $Y \setminus \{\infty^-, \infty^+\}$, $\mathcal{U}^+$ and $\mathcal{U}^-$, which are glued by the function $\exp\left(2\pi\sqrt{-1}\hat{p}\frac{\check{\gamma}_1 + \sqrt{-1}\check{\gamma}_2}{\check{\gamma}_1 - \sqrt{-1}\check{\gamma}_2}\right)$ on $(Y \setminus \{\infty^+\}) \cap \mathcal{U}^+$ and by the function $\exp\left(2\pi\sqrt{-1}\hat{p}\frac{\check{\gamma}_1 - \sqrt{-1}\check{\gamma}_2}{\check{\gamma}_1 + \sqrt{-1}\check{\gamma}_2}\right)$ on $(Y \setminus \{\infty^+\}) \cap \mathcal{U}^-$. Due to the Direct Image theorem [Gr-Re, Chapter 10. §4.6.] the direct image of the sheaf of sections of this line bundle under $\tilde{f}$ is a coherent sheaf on $\tilde{\mathcal{M}}$. The subset of this manifold, on which another function $\check{p}$ is defined and whose combination k fulfills conditions *Quasi-momenta* (i)–(ii) is exactly the support of this direct image sheaf. In fact, any non-zero germ of this direct image sheaf defines a global holomorphic function on $Y \setminus \{\infty^-, \infty^+\}$ with essential singularity at the two marked points. Since the degree of the family of line bundles on the family of complex spaces is zero, this global holomorphic function does not vanish, and the logarithm of this function divided by some uniquely defined complex number yields a function $\check{p}$ with the desired properties. Since the support of any coherent sheaf is an analytic set [Gr-Re, Annex. §4.5.] this establishes the claim.

10. This last result implies that the subset of $\tilde{\mathcal{M}}$ corresponding to elements of $\mathcal{M}_{g,\Lambda}$ contains an open dense subset of nonsingular points, and this open dense subset forms a submanifold of $\tilde{\mathcal{M}}$. Moreover, the derivative of $\check{p}$ with respect to any vector field along this submanifold defines some regular form on the corresponding family of complex subspaces. We conclude that the dimension of this complex space is equal to $g+1$, and that the tangent sheaf extends to a locally free sheaf of rank $g+1$. Now the same arguments as in the proof of [G-P-R, Chapter II. §1.4. Theorem 1.12.] (i. e. [G-P-R, Chapter II. §1.4. Fact 1.13.] combined with [G-P-R, Chapter I. §10.5. (10.20)]) imply that this subset is smooth. **q.e.d.**

For the proof of the existence of the minimizer we shall only consider *complex Fermi curves* of bounded genus. Therefore, we should investigate the subspace of this moduli space containing only *complex Fermi curves* of finite *geometric genus*. Obviously, a given *complex Fermi curve* of *geometric genus* g corresponds to some element in $\mathcal{M}_{g,\Lambda}$ if and only if the normalization does not contain zeroes of both differentials $d\hat{p}$ and $d\check{p}$. In this case Y may be chosen to be the normalization. Otherwise there exists a unique representative element $(Y, \infty^-, \infty^+, k) \in \mathcal{M}_{g',\Lambda}$ of minimal arithmetic genus g' . In this case Y is locally isomorphic to the images of the normalization under k , and the arithmetic genus is equal to the genus of the normalization plus the *Local contribution to the arithmetic genus* 4.3 of all these common zeroes of $d\hat{p}$ and $d\check{p}$, which are called *cusps*. Unfortunately, the subspace of $\mathcal{M}_{g',\Lambda}$ containing all *complex Fermi curves* of *geometric genus* g , has itself *cusps* at *complex Fermi curves*, whose normalization contains *cusps*.

Proposition 3.9. *For any complex Fermi curve $\mathcal{F}(Y, \infty^-, \infty^+, k) \in \mathcal{M}_{g',\Lambda}$, whose normalization contains a cusp and has genus $g < g'$, and all meromorphic forms ω on the normalization of Y with at most first-order poles at $\infty^\pm$ and no other poles, there exists a one-dimensional complex subspace of some neighbourhood of $\mathcal{F}$ in $\mathcal{M}_{g',\Lambda}$, which contains $\mathcal{F}$, and which is contained in the set of the complex Fermi curves of geometric genus g .*

Moreover, this one-dimensional subspace contains a cusp at $\mathcal{F}$ in tangent direction of the line corresponding to ω . More precisely, the preimage of $\mathcal{F}$ in the normalization of this one-dimensional subspace contains a single point. Furthermore, with respect to any local parameter of the normalization at this point the lowest non-vanishing derivative is proportional to ω , considered as an element of the tangent space of $\mathcal{M}_{g',\Lambda}$ over $\mathcal{F}$.

Proof. This proposition is proven in nine steps.

1. First we remark that due to Proposition 3.7 the flow of a vector field on some neighbourhood of some *complex Fermi curve* without *cusps*, which takes values in the subspace of the tangent space corresponding to meromorphic forms on the normalization of the corresponding *complex Fermi curve* with at most first-order poles at $\infty^\pm$ and no other poles, preserves the genus of the normalization of the corresponding *complex Fermi curves*.

2. We shall see that again we may transform a *cusp* of the *complex Fermi curve* into *multiple points* (i. e. singularities, whose preimage in the normalization does not contain zeroes of both differentials $d\hat{p}$ and $d\check{p}$) by blowing up the moduli space. We choose the unique representative $(Y, \infty^-, \infty^+, k)$ of minimal arithmetic genus described above. Without loss of generality we may choose the local descriptions of the elements of $\tilde{M}$ to be locally zero sets of equations $\check{p}^d - \hat{p}^m = 0$ with two co-prime natural numbers d and m . Since we are interested in deformations, which preserve the *geometric genus*, we change the description of $\tilde{M}_l$ near the *cusps*. All these deformations correspond locally to those deformations in $\tilde{M}_l$, whose equations $Q(\hat{p}, \check{p}, t_1, \dots, t_r) = 0$ describe algebraic curves birational isomorphic to $\mathbb{P}^1$. In particular, the functions $\hat{p}$ and $\check{p}$ may be written as polynomials

$$\hat{p} = z^d + \hat{t}_1 z^{d-1} + \hat{t}_2^2 z^{d-2} + \dots + \hat{t}_d^d \text{ and } \check{p} = z^m + \check{t}_1 z^{m-1} + \check{t}_2^2 z^{m-2} + \dots + \check{t}_m^m.$$

The coefficients are written in terms of powers of the moduli $(\hat{t}_1, \dots, \hat{t}_d, \check{t}_1, \dots, \check{t}_m) \in \check{\mathcal{M}}_l$ in such a way, that the re-scaling $z \mapsto \lambda z$ acts projectively on the moduli. For all these moduli there exists an unique polynomial with

$$Q(\hat{p}, \check{p}, \hat{t}_1, \dots, \hat{t}_d, \check{t}_1, \dots, \check{t}_m) = 0.$$

If we define the degree of $\hat{p}$ as d and the degree of $\check{p}$ as m this polynomial has common degree dm , and the highest coefficient is equal to $\check{p}^d - \hat{p}^m$.

3. We should remark that the translations $z \mapsto z + z_0$ do not change Q , but change the moduli. Furthermore, a finite groups acts on the moduli without changing the coefficients of the polynomials $\hat{p}$ and $\check{p}$ with respect to z . For this reason the moduli space should be defined as the quotient by the action of the translations and the finite group on the moduli. The orbits of this group action define one-dimensional curves in the parameter space parameterized by z_0 . At some points these curves might have singularities (i. e. the derivatives of all parameters with respect to z_0 vanish), but obviously for all orbits this happens only at finitely many values of z_0 . Therefore, we may choose some non-singular representative and use the implicit function theorem [R-S-I, Supplementary material V.5 Theorem S.11] in order to obtain locally smooth parameters of the quotient by this group action. In the sequel we will ignore this ambiguity of the moduli $(\hat{t}_1, \dots, \hat{t}_d, \check{t}_1, \dots, \check{t}_m)$ and assume that the real moduli space is locally the quotient by this group action.

4. If we glue the local normalizations of the deformed spaces, parameterized again by z on $\mathcal{U}_0$ in the natural way, then the arguments in the proof of Proposition 3.7 carry over to the modified moduli space $\check{\mathcal{M}}$. They show that the space of meromorphic forms on the normalizations, which have at most first-order poles at $\infty^\pm$ and no other poles, define a holomorphic vector bundle on $\check{\mathcal{M}}$ of rank $g+1$ or g (compare with Remark 3.8). Therefore, all these forms on the normalization of the initial *complex Fermi curve* extend to holomorphic sections of this vector bundle. Now we shall construct the deformations corresponding to the sections of this vector bundle by blowing up the moduli space $\check{\mathcal{M}}$ at the origin corresponding to the initial *complex Fermi curve*. The blowing up re-parameterizes locally the moduli as elements of the tautological bundle over the corresponding projective space [Har, pp. 28-31]. Locally this blowing up is the product of the projective space with a one-dimensional space parameterizing the fibre of the tautological bundle. We should choose an element $(\hat{t}_1, \dots, \hat{t}_d, \check{t}_1, \dots, \check{t}_m) \in \mathbb{P}^{d+m-1}$ of the projective space, such that the deformation changes in lowest-order the parameter of the fibre. All coefficients of the polynomial $Q(\hat{p}, \check{p})$ corresponding to some moduli are homogeneous polynomials of the moduli, whose degree is equal to dm minus the common degree of the corresponding monomial of $\hat{p}$ and $\check{p}$. Due to Lemma 2.20 the regular form ω may be locally uniquely written as

$$\omega = \frac{\dot{Q}(\hat{p}, \check{p})}{\partial Q(\hat{p}, \check{p})/\partial \check{p}} d\hat{p},$$

where $\dot{Q}$ is a polynomial with respect to $\check{p}$ of degree $d-1$, whose coefficients are holomorphic functions depending on $\hat{p}$. We conclude that besides the coefficient of degree zero (corresponding to $\check{p}^d - \hat{p}^m$) those coefficients of $Q(\hat{p}, \check{p})$ corresponding to the projective element $(\hat{t}_1, \dots, \hat{t}_d, \check{t}_1, \dots, \check{t}_m) \in \mathbb{P}^{d+m-1}$ we start with should be zero, which are of lower degree than the degree of the unique coefficient, whose monomial corresponds to the lowest non-vanishing Taylor coefficient of ω . Furthermore, the coefficient of $Q(\hat{p}, \check{p})$ corresponding to the projective element we start with, whose monomial corresponds to the lowest non-vanishing Taylor coefficient, should not vanish.

5. Now we shall determine those projective moduli $(\hat{t}_1, \dots, \hat{t}_d, \check{t}_1, \dots, \check{t}_m)$ we have to start with, if ω does not vanish at the corresponding *cusp*. The common degree of the unique monomial $\dot{Q}(\hat{p}, \check{p})$, corresponding locally to a holomorphic non-vanishing form, is equal to $(d-1)(m-1)$. Therefore, besides the coefficient of degree zero all coefficients of degree lower than $dm - (d-1)(m-1) = d+m-1$ of the polynomial $Q(\hat{p}, \check{p})$, which corresponds to those projective moduli we start with, should vanish. There are exactly $d+m-2$ such coefficients. Hence we have to determine the zero set of $d+m-2$ homogeneous polynomials on $\mathbb{P}^{d+m-1}$. Due to [Har, Chapter I Theorem 7.2] this set is at least one-dimensional. In order to investigate these solutions let us first determine the derivative of the coefficients with respect to some infinitesimal change $(\dot{\hat{p}}, \dot{\check{p}})$ of the polynomials $\hat{p}$ and $\check{p}$. The equations

$$\frac{dQ(\hat{p}, \check{p})}{dt} = \dot{Q}(\hat{p}, \check{p}) + \frac{\partial Q(\hat{p}, \check{p})}{\partial \hat{p}} \dot{\hat{p}} + \frac{\partial Q(\hat{p}, \check{p})}{\partial \check{p}} \dot{\check{p}} = 0 \quad \frac{\partial Q(\hat{p}, \check{p})}{\partial \hat{p}} d\hat{p} + \frac{\partial Q(\hat{p}, \check{p})}{\partial \check{p}} d\check{p} = 0$$

imply the identity

$$\dot{Q}(\hat{p}, \check{p}) \frac{d\hat{p}}{\partial Q(\hat{p}, \check{p})/\partial \check{p}} = -\dot{Q}(\hat{p}, \check{p}) \frac{d\check{p}}{\partial Q(\hat{p}, \check{p})/\partial \hat{p}} = \dot{\hat{p}} d\check{p} - \dot{\check{p}} d\hat{p}.$$

For the initial curve defined by $\check{p}^d - \hat{p}^m = 0$ the right hand side divided by $\dot{Q}(\hat{p}, \check{p})$ is equal to $z^{-(d-1)(m-1)} dz$. If we consider the projective deformation from the initial curve (defined by $\check{p}^d - \hat{p}^m = 0$) into one of these solutions, then the derivative with respect to the lowest power of the deformation parameter (the parameter of the fibre of the tautological bundle), which does not vanish, yields a meromorphic one-form with poles only at $z = \infty$. We conclude that the corresponding $\dot{Q}(\hat{p}, \check{p})$ has to be a polynomial with respect to z , whose degree is larger than $(d-1)(m-1)$. In particular, the coefficients of degree $d+m-1$ of the polynomials $Q(\hat{p}, \check{p})$, which correspond to these solutions, cannot vanish. Therefore, for these deformations the derivative with respect to the $(d+m-1)$ -th power of the deformation parameter at the initial curve corresponds to a one-form proportional to dz .

6. Now we claim that the normalizations of these curves do not have a *cusp*. In fact, otherwise we shift the *cusp* into the origin ($z = 0$). In this case, due to the foregoing formula, the derivative with respect to the lowest non-vanishing (i. e. the $(d+m-1)$ -th) power of the deformation parameter, corresponds to a one-form, which vanishes at $z = 0$. This contradicts the considerations in step 5.

7. Now we show that, if we choose some non-singular representative $(\hat{t}_1, \dots, \hat{t}_d, \check{t}_1, \dots, \check{t}_m)$ in the orbit under the group described above, then the derivative of the map, which maps the moduli onto the coefficients of the corresponding polynomial $Q(\hat{p}, \check{p})$, has rank $d+m-1$. This statement is equivalent to the statement, that the derivative of the map, which maps the coefficients of the polynomials $\hat{p}$ and $\check{p}$ with respect to z onto the coefficients of the corresponding polynomial $Q(\hat{p}, \check{p})$, has rank $d+m-1$. The foregoing formula shows that a tangent vector $(\dot{\hat{p}}, \dot{\check{p}})$ in the space of polynomials $\hat{p}$ and $\check{p}$ with respect to z does not change Q , if and only if the right hand side vanishes. Since the curve defined by the equation $Q(\hat{p}, \check{p}) = 0$ does not have *cusps*, $\dot{\hat{p}}$ has to be zero at the zeroes of $d\hat{p}$, and $\dot{\check{p}}$ has to be zero at the zeroes of $d\check{p}$. We conclude that $(\dot{\hat{p}}, \dot{\check{p}})$ has to be proportional to $(d\hat{p}/dz, d\check{p}/dz)$, which corresponds exactly to translations.

8. With the help of the implicit function theorem [R-S-I, Supplementary material V.5 Theorem S.11] we conclude from the last step that in some arbitrary small neighbourhood of those non-singular moduli, we start with if ω does not vanish at the *cusp*, there also exist moduli, whose $Q(\hat{p}, \check{p})$ has besides the highest term $\check{p}^d - \hat{p}^m$ vanishing coefficients of degree larger than those corresponding to the lowest non-vanishing coefficient of ω . Therefore, we may choose these moduli in such a way, that the corresponding curves $Q(\hat{p}, \check{p}) = 0$ also have no *cusps*, and that the derivative of the map, which maps the moduli onto the coefficients of $Q(\hat{p}, \check{p})$, has rank $d+m-1$. This in general defines the projective moduli, we start with.

9. Finally, we multiply the pullback of the section of the vector bundle of meromorphic forms on the normalizations with poles of at most first-order at $\infty^\pm$ and no other poles, to the blowing up of the moduli space $\check{\mathcal{M}}$, with the parameter of the fibre of the tautological bundle to the power, which is equal to the order of the zero of ω plus $d+m-2$, and which is also equal to the degree of the corresponding coefficient of Q minus one. With the help of the construction of Proposition 3.7 the flow defined by $\check{p}d\hat{p} = \omega$ can be locally integrated along the vector field on the blowing up of the moduli space $\check{\mathcal{M}}$. The image of the corresponding curve in $\check{\mathcal{M}}$ is contained in $\mathcal{M}$ and yields the desired curve. **q.e.d.**

This proposition can be generalized to deformations of one-sheeted coverings of the *complex Fermi curves*, which are locally not complete intersection in $\mathbb{C}^2$.

If we consider deformations, which preserve an anti-holomorphic involution with fixed points, then the real regular forms describe the corresponding tangent space. In case of a real cusp of the *complex Fermi curve*, the construction of Proposition 3.9 has to be done over the real numbers. In particular, the moduli we start with, should be real. More precisely, it suffices to find two polynomials

$$\hat{p}(z) = z^d + a_1 z^{d-1} + \dots + a_d \text{ and } \check{p}(z) = z^m + b_1 z^{m-1} + \dots + b_m$$

with real coefficients $a_1, \dots, a_d$ and $b_1, \dots, b_m$, whose coefficients of the corresponding polynomial $Q(\hat{p}, \check{p})$ of degree $1, \dots, d+m-2$ vanish. Due to the considerations in part 5. of the forgoing proof this is equivalent to the relation

$$\hat{p}'(z/t)t^d \frac{d}{dt} \check{p}(z/t)t^m - \check{p}'(z/t)t^m \frac{d}{dt} \hat{p}(z/t)t^d = t^{d+m-2} \mathbf{O}(1) \text{ with respect to } z.$$

This relation in turn is equivalent to the relation

$$m\hat{p}'(z)\check{p}(z) - d\check{p}'(z)\hat{p}(z) = \mathbf{O}(1) \text{ with respect to } z.$$

First we remark that due to the invariance under the translations $z \mapsto z + z_0$, we may assume that $a_1 = 0$. Moreover, due to the structure of this relation, the corresponding equations of order $z^{m+d-1}, \dots, z^{d-1}$ are linear in $b_1, \dots, b_m$. Therefore, we may use these equations to obtain $b_1, \dots, b_m$ as polynomials with respect to $a_2, \dots, a_d$. The corresponding polynomial $\check{p}(z)$ is determined by the m lowest Taylor coefficients of $(z^{-d}\hat{p}(z))^{m/d} = (1 + a_1 z^{-1} + \dots + a_d z^{-d})^{m/d}$ at infinity:

$$\check{p}(z) = z^m (z^{-d}\hat{p}(z))^{m/d} + \mathbf{O}(z^{-1}).$$

It remains to find zeroes of $d-2$ homogeneous polynomials with respect to $a_2, \dots, a_d$ of common degree $m+1, \dots, m+d-2$, where the degree of a_l is equal to l for all $l = 2, \dots, d$. We conclude that for $d = 2$ there always exists one unique real solution.

If $d = 3$ and m is even, then there also exists a real solution. In fact in this case $a_3 = 0$ is always a solution. If m is odd and $m+1$ is not divisible by 12 then there again exists a real solution, since a polynomial, whose degree is even but not divisible by 12 with respect to a_2 and a_3 is either zero for $a_2 = 0$ or $a_3 = 0$ or is an odd polynomial either with respect to a_2 or a_3 . A direct calculation shows that for $m = 11$ there also exists a real solution.

If $d = 4$, the choice $a_3 = 0$ reduces the number of equations by one. Therefore, we have to find a real zero of a polynomial of degree $m+1$ with respect to a_2 and a_4 . If $m+1$ is not divisible by 8 such a polynomial has always a real solution.

We conjecture that there always exists a real solution of these equations. But if $d+m-1$ is even (and therefore not both degrees are odd), then the real part of the one-dimensional subspace of the moduli space, which is constructed in Proposition 3.9 with the help of a real polynomials constructed above, covers only one half of the real line in the tangent space of the moduli space corresponding to any real regular form, which does not vanish at the

cusps. Since in the simplest case $d = 2$ and $m = 3$ there exists only one real solution, the subspace of the moduli space containing *complex Fermi curves*, which are invariant under the anti-holomorphic involution, and which have the same *geometric genus* as the original *complex Fermi curve* with the real cusp, covers only a half-plane of the real tangent space.

3.3 The compactified moduli spaces with bounded genus

In Section 4.1 we shall find some necessary condition for a *complex Fermi curve* to correspond to some immersion. Since the Willmore functional is equal to the *first integral* of the corresponding *Weierstraß curve* 1.4, we may transform the variational problem of the Willmore functional into a variational problem of the *first integral* on the space of all *complex Fermi curves*, which obey the *Weak Singularity condition* 4.2. For this purpose we should endow the set of all *complex Fermi curves* with some topology. By definition all *complex Fermi curves* are closed subsets of $\mathbb{C}^2$. Consequently we shall use some topology on the set of closed subsets of $\mathbb{C}^2$. The set of closed subsets of some compact Hausdorff space has a natural topology in contrast to the space of closed subsets of a non-compact Hausdorff space [Mi]. Obviously the closures of all *complex Fermi curves* considered as subsets of the one-point-compactification $\overline{\mathbb{C}^2}$ of $\mathbb{C}^2$ are given by the union of the *complex Fermi curves* (contained in $\mathbb{C}^2$) with the one point set $\{\infty\}$ containing the single point at infinity. In the sequel we shall identify the closed subsets of $\mathbb{C}^2$ with the corresponding closures in $\overline{\mathbb{C}^2}$.

Let us therefore introduce a topology on the space of all closed subsets of some topological space ([Ho, Chapter 5.4] and [Mi]). In [Mi] several topologies are given, and we will use the so called *finite topology*. For all open sets $\mathcal{O}$ the subsets of the set of closed subsets of the form $\{\mathcal{A} \mid \mathcal{A} \cap \mathcal{O} \neq \emptyset\}$ and $\{\mathcal{A} \mid \mathcal{A} \subset \mathcal{O}\}$ are open and generate the set of open subsets of the *finite topology*. We recall some of the properties of this topology proven in [Mi]. If the underlying space is a compact Hausdorff space, the space of closed subsets endowed with this *finite topology* is again a compact Hausdorff space. Moreover, if the underlying space is in addition metrizable the space of all closed subsets is also metrizable and coincides with the topology of the Hausdorff metric. Finally, the subspace of all finite subsets is dense, therefore the space of all closed subsets is separable if the underlying space is separable. The following lemma is a special case of results proven in [Mi].

Lemma 3.10. *The space of all closed subsets of $\overline{\mathbb{C}^2}$ is a separable compact metrizable space. q.e.d.*

Due to Theorem 2.3, for any pair of potentials (V, W) the mapping $k \mapsto R(V, W, k, 0)$ is a meromorphic mapping from $\mathbb{C}^2$ into the compact operators on $L^2(\Delta) \times L^2(\Delta)$. Moreover, this mapping uniquely determines the pair of potentials. The topology of uniform convergence on compact subsets of these mappings [Co-I, Chapter VII. §1.] yields a topology on $(V, W) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$, which we call the *compact open topology* of the resolvents $R(V, W, k, 0)$. Obviously this topology is equivalent to the *compact open topology* of the resolvents of $\tilde{D}_{\hat{\gamma}, \hat{\gamma}}(V, W, \hat{p})$ (compare with Section 2.4).

We claim that the mapping $(V, W) \mapsto \bar{\mathcal{F}}(V, W)$ is continuous with respect to this topology and the *finite topology*. In fact, since the two families of subsets $\{\bar{\mathcal{F}} \subset \overline{\mathbb{C}^2} \mid \bar{\mathcal{F}} \cap \mathcal{O} \neq \emptyset\}$ and

$\{\bar{\mathcal{F}} \subset \overline{\mathbb{C}^2} \mid \bar{\mathcal{F}} \subset \mathcal{O}\}$ indexed by the open subsets $\mathcal{O}$ of $\overline{\mathbb{C}^2}$ form a basis of the open subsets of the *finite topology*, it suffices to show that the corresponding preimages are open. If the closure $\bar{\mathcal{F}}$ of a *complex Fermi curve* is contained in an open subset of the former family, then there exists a finite $k \in \mathcal{F} \cap \mathcal{O}$. Furthermore, due to the application of the arguments in the proof of Theorem 2.3 concerning the spectral projections to the corresponding spectral projections of $\tilde{D}_{\hat{\gamma}, \hat{\gamma}}(V, W, \hat{p})$, the same is true for all pairs of potentials in a neighbourhood with respect to the *compact open topology*. If the closure $\bar{\mathcal{F}}$ is contained in an open subset of the latter family, then $\mathcal{O}$ has to contain ∞ . Furthermore, all *complex Fermi curves* of pairs of potentials in a neighbourhood (with respect to the *compact open topology*) are also disjoint with the complement of $\mathcal{O}$, since this complement is a compact subset of $\mathbb{C}^2$. This proves

Lemma 3.11. *The mapping $(V, W) \mapsto \bar{\mathcal{F}}(V, W)$ is a continuous mapping from $(V, W) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ with the compact open topology of $R(V, W, k, 0)$ into the closed subsets of $\overline{\mathbb{C}^2}$ with the finite topology. q.e.d.*

Let $\mathcal{M}_{\Lambda, \eta}$ and $\mathcal{M}_{\Lambda, \eta, \mathcal{W}}$ denote the moduli spaces

$$\begin{aligned} \mathcal{M}_{\Lambda, \eta} &= \{ \mathcal{F}(U, \bar{U}) \mid U \in L^2(\mathbb{R}^2/\Lambda) \} \\ \mathcal{M}_{\Lambda, \eta, \mathcal{W}} &= \{ \mathcal{F}(U, \bar{U}) \mid U \in L^2(\mathbb{R}^2/\Lambda) \text{ and } 4\|U\|^2 \leq \mathcal{W} \}. \end{aligned}$$

Both sets are subsets of the space of closed subsets of $\overline{\mathbb{C}^2}$. The subset of these moduli spaces, which contain the corresponding *complex Fermi curves* of real potentials are denoted by $\mathcal{M}_{\Lambda, \eta, \sigma}$ and $\mathcal{M}_{\Lambda, \eta, \sigma, \mathcal{W}}$, respectively.

We will see that neither $\mathcal{M}_{\Lambda, \eta}$ nor $\mathcal{M}_{\Lambda, \eta, \mathcal{W}}$ is compact, if $\mathcal{W}$ is not smaller than 4π (compare with Remark 2.5 and Remark 2.8). In fact, in Section 2.5 we have introduced conditions on compact one-dimensional complex spaces, which for the dense subclass of *connected* complex spaces implies, that they are *complex Fermi curves*. In Lemma 4.16 we will investigate the subclass of complex spaces with two connected components. It will turn out that the simplest example is not a *complex Fermi curve* of any potential, and that the corresponding *first integral* is equal to 4π . But they are the *complex Fermi curves* of perturbations of Dirac operators acting on direct sums of line bundles of degree ± 1 . In the next section we shall prove that the set of all *complex Fermi curves* of higher-degree perturbations of Dirac operators, whose *first integral* is bounded by any $\mathcal{W} > 0$ is compact. Higher-degree perturbations of Dirac operators also occur in ‘quaternionic function theory’ [P-P].

In this section we restrict ourself to the moduli spaces of *complex Fermi curves* of bounded *geometric genus*: Let $\mathcal{M}_{g, \Lambda, \eta}$ and $\mathcal{M}_{g, \Lambda, \eta, \mathcal{W}}$ denote the moduli spaces

$$\begin{aligned} \mathcal{M}_{g, \Lambda, \eta} &= \{ \mathcal{F} \in \mathcal{M}_{\Lambda, \eta} \mid \text{geometric genus of } \mathcal{F} \text{ is not larger than } g \} \\ \mathcal{M}_{g, \Lambda, \eta, \mathcal{W}} &= \mathcal{M}_{\Lambda, \eta, \mathcal{W}} \cap \mathcal{M}_{g, \Lambda, \eta}. \end{aligned}$$

The corresponding subsets of these moduli spaces, which contain only *complex Fermi curves* of real potentials are denoted by $\mathcal{M}_{g, \Lambda, \eta, \sigma}$ and $\mathcal{M}_{g, \Lambda, \eta, \sigma, \mathcal{W}}$, respectively.

In Section 2.5 we characterized these *complex Fermi curves* of finite genus by the conditions *Quasi-momenta* (i)–(iii) of Section 2.5. There exists an explicit formula of the *first*

integral, whose restriction to *Weierstraß curves* 1.4 is equal to the Willmore functional, in terms of the data $(Y, \infty^-, \infty^+, k)$.

Lemma 3.12. *If the complex Fermi curve $\mathcal{F}(Y, \infty^-, \infty^+, k)$ of some data $(Y, \infty^-, \infty^+, k)$, which fulfills conditions *Quasi-momenta* (i)–(ii) of Section 2.5, is equal to the complex Fermi curve of some pair of potentials (V, W) , then the corresponding first integral is equal to the residue*

$$4 \int_{\mathbb{R}^2/\Lambda} V(x)W(x)d^2x = \mathcal{W}(Y, \infty^-, \infty^+, k) = 8\pi^2\sqrt{-1} \operatorname{vol}(\mathbb{R}^2/\Lambda) \operatorname{Res}_{\infty^+}(k_1 dk_2).$$

Obviously the residue on the right hand side does not depend on the choice of the branch of k . Below we will show, that whenever the data fulfill condition *Quasi-momenta* (iii) of Section 2.5 these numbers are non-negative.

Proof. The complex Fermi curve corresponding to the zero potentials is given by the union of the solutions of the equations $g(k+\kappa, k+\kappa) = 0$ for $\kappa \in \Lambda^*$. An easy calculation shows that in this case the formula is correct. If we apply formula Lemma 3.2 (iv) to the variation described in Lemma 3.4 (iv) the right hand side is equal to $1/(\pi^2\sqrt{-1})$ times the variation of the *first integral*. Since this variation corresponds to the function, which in some neighbourhood of ∞^+ is equal to 1 and in some neighbourhood of ∞^- equal to -1 , the left hand side is equal to the residue of $\Omega_{V,W}(\cdot, \cdot)$ at ∞^+ minus the residue of this form at ∞^- . If we consider the function $\tilde{p}$ as a function depending on $\hat{p}$ and the deformation parameters, we conclude from the definition of Ω that the left hand side is equal to the residue of the form $\delta\tilde{p}d\hat{p}$ at ∞^+ . This implies that the *first integral* is equal to the residue at ∞^+ of the form $2\pi^2\sqrt{-1}\tilde{p}d\hat{p}$. The right hand side of the formula of the lemma is invariant under linear conformal transformations of the lattice Λ . If we choose $\gamma = (1, 0)$ the formula follows. **q.e.d.**

Consequently, define $\bar{\mathcal{M}}_{g,\Lambda,\eta}$ as the set of all complex Fermi curves $\mathcal{F}(Y, \infty^-, \infty^+, k)$, whose data fulfill conditions *Quasi-momenta* (i)–(iii) of Section 2.5 and whose *geometric genus* is not larger than g . Analogously we denote the subset of all varieties $\mathcal{F}(Y, \infty^-, \infty^+, k)$ of data, whose *first integral* $\mathcal{W}(Y, \infty^-, \infty^+, k)$ is not larger than $\mathcal{W}$ by $\bar{\mathcal{M}}_{g,\Lambda,\eta,\mathcal{W}}$. The subsets of these moduli spaces containing the corresponding complex Fermi curves, which are invariant under the involution σ , are denoted by $\bar{\mathcal{M}}_{g,\Lambda,\eta,\sigma}$ and $\bar{\mathcal{M}}_{g,\Lambda,\eta,\sigma,\mathcal{W}}$, respectively. Now we can state the main theorem of this section.

Theorem 3.13. *The spaces $\bar{\mathcal{M}}_{g,\Lambda,\eta,\mathcal{W}}$ are compact and the completions of $\mathcal{M}_{g,\Lambda,\eta,\mathcal{W}}$.*

For the proof of this theorem we need some preparation. Let $\bar{\mathcal{M}}_{g,\eta}$ denote the set of all varieties $\mathcal{F}(Y, \infty^-, \infty^+, k)$, whose data obey conditions *Quasi-momenta* (i) and (iii) of Section 2.5, but instead of condition *Quasi-momenta* (ii) we assume condition

Quasi-momenta (ii') The difference of two arbitrary branches of the function k is some element of $\mathbb{R}^2$.

In this case $d\hat{p}$ and $d\check{p}$ are meromorphic forms, and the multi-valued function k is determined by this form up to the addition of some complex numbers. So let us investigate for a given data $(Y, \infty^-, \infty^+, k)$ the space of such forms $d\hat{p}$ and $d\check{p}$.

Lemma 3.14. *Let Y be a compact smooth Riemann surface of genus g with two marked points ∞^- and ∞^+ with the property that each connected component of Y contains at least one point. Moreover, let η be an anti-holomorphic involution, which interchanges the two marked points. Then for any local parameter z at ∞^+ , which takes the value zero at ∞^+ , there exists a unique meromorphic form dp with the following properties:*

Form (i) $dp - d(1/z)$ extends to a holomorphic form on some neighbourhood of ∞^+ .

Form (ii) The integral of dp along any cycle in $H_1(Y, \mathbb{Z})$ is real.

Form (iii) dp transforms under η as $\eta^*dp = -\bar{d}p$.

Form (iv) The integral of dp along any real cycle (i. e. the cycle is homolog to the image under η) vanishes.

Proof. Obviously conditions *Form* (i)–(iii) are equivalent to conditions *Form* (i) together with the conditions *Form* (iii)–(iv). Hence we may assume this last triple of conditions. The Riemann–Roch theorem [Fo, Theorem 16.9 and Theorem 17.16] implies that the space of meromorphic forms, which have poles of at most second order without residues at the two marked points and no other poles is a $(g+2)$ -dimensional complex space. Since the involution η acts on this space as an anti-linear involution, the eigenspace of this action corresponding to the eigenvalue -1 is a $(g+2)$ -dimensional real space. Hence there exists some dp obeying conditions *Form* (i) and (ii). This form is determined up to the addition of some holomorphic form ω with the property that $\eta^*\omega = -\bar{\omega}$. Since the real part of $H_1(Y, \mathbb{Z})$ forms a Lagrangian subgroup with respect to the intersection form (i. e. a subgroup, which coincides with the group of all elements having zero intersection form with all elements of this subgroup), any set of generators of this real part gives rise to a dual base of holomorphic differential forms ω with the property that $\eta^*\omega = -\bar{\omega}$ [F-K, III.3.3. Proposition]. This proves the Lemma. **q.e.d.**

We conclude that for all data $(Y, \infty^-, \infty^+, \eta, k) \in \bar{\mathcal{M}}_{g,\eta}$ the components $\hat{p}$ completely determine the other components, and more generally, any real component of k determines the function k . So in the sequel we will associate to the elements of $\bar{\mathcal{M}}_{g,\Lambda,\eta}$ data of the form $(Y, \infty^-, \infty^+, p)$ where p is a multi-valued in $\mathbb{C}$ instead of $\mathbb{C}^2$ (either equal to $\hat{p}$ or any other real component of k). In this case we assume that these data obey the conditions analogous to *Quasi-momenta* (i), (ii') and (iii). Obviously, $(Y, \infty^-, \infty^+, \hat{p})$ and $(Y, \infty^-, \infty^+, \check{p})$ are data of this form. Also the formula of Lemma 3.12 for the *first integral* extends to elements of $\bar{\mathcal{M}}_{g,\eta}$. On these larger moduli spaces we may *renormalize* the functions k by multiplication with some positive number. The *first integral* changes under these transformations by multiplication with the square of this positive number. It is quite easy to see that the quotient of the moduli spaces $\bar{\mathcal{M}}_{g,\eta}$ modulo these transformations and modulo shifts of k by some additive constant, is the moduli space of all compact Riemann surfaces of genus g with two marked points $\infty^\pm$, at most two connected components, each containing at least one of

the two marked points, and the anti-holomorphic involution η without fixed points interchanging $\infty^\pm$. Therefore, the description of these Riemann surfaces in Lemma 3.16 yields a parameterization of this moduli space. A simple count of degrees of freedom shows that $\bar{\mathcal{M}}_{g,\eta}$ has real dimension $3g + 2$. This coincides with the dimension of the moduli space of the corresponding Riemann surfaces with two marked points $(3g - 1)$ plus the degree of freedom of renormalization of k and the two additive degrees of freedom given by the complex additive constant just mentioned.

Let us now show that for all these data $(Y, \infty^-, \infty^+, p)$ the Riemann surface Y may be realized as two copies of $\mathbb{P}^1$, which are glued along several cuts. In the sequel we will use this construction for different choices of the function p , therefore we shall decorate the corresponding symbols with an index p . First we choose a branch of the function p in some small neighbourhood of ∞^+ . Obviously there exists an open neighbourhood $\mathcal{U}_p$ of ∞^+ obeying the following conditions:

Neighbourhood (i) The fixed branch of p near ∞^+ maps $\mathcal{U}_p$ biholomorphically onto the complement in $\mathbb{P}^1$ of finitely many disjoint compact convex sets contained in $\mathbb{C} \subset \mathbb{P}^1$. In particular, $\mathcal{U}_p$ contains no zero of dp .

Neighbourhood (ii) Each connected component of the complement of $\mathcal{U}_p \cup \eta(\mathcal{U}_p)$ contains at most one connected component of the boundary of the closures of $\mathcal{U}_p$.



Due to condition *Neighbourhood* (i) the connected components of the boundary of the closure of $\mathcal{U}_p$ are in one to one correspondence with the compact convex subsets mentioned there.

Lemma 3.15. *There exists a unique maximal open neighbourhood $\mathcal{U}_p^+$ of ∞^+ obeying conditions Neighbourhood (i)–(ii). Moreover, the set $\mathcal{U}_p^- = \eta(\mathcal{U}_p^+)$ is the unique maximal open neighbourhood of ∞^- obeying the corresponding conditions for the branch $-\eta^*p$ near ∞^- . Both sets $\mathcal{U}_p^+$ and $\mathcal{U}_p^-$ are disjoint.*

Proof. Let us first prove that the union of two open neighbourhoods $\mathcal{U}_p$ and $\mathcal{U}'_p$ of ∞^+ obeying these conditions obeys also these conditions. Obviously the union of two complements of finitely many disjoint compact convex sets is again a complement of finitely many disjoint compact convex sets. Due to condition *Quasi-momenta* (ii') the subset of all elements in the intersection of the images of $\mathcal{U}_p$ and $\mathcal{U}'_p$ under the corresponding branches of p , whose preimages are the same element of $\mathcal{U}_p \cap \mathcal{U}'_p$, is open and closed. Hence it is a connected component of $\mathcal{U}_p \cap \mathcal{U}'_p$. But the image of $\mathcal{U}_p \cap \mathcal{U}'_p$ under the corresponding branches of p is the complement of finitely many not necessary disjoint compact convex subsets of $\mathbb{P}^1$ and therefore connected. Hence the branches of p corresponding to $\mathcal{U}_p$ and $\mathcal{U}'_p$, coincide, and $\mathcal{U}_p \cup \mathcal{U}'_p$ obeys condition *Neighbourhood* (i). Since the complement of $\mathcal{U}_p \cup \mathcal{U}'_p \cup \eta(\mathcal{U}_p \cup \mathcal{U}'_p)$ is equal to the intersection of the complements of $\mathcal{U}_p \cup \eta(\mathcal{U}_p)$ and $\mathcal{U}'_p \cup \eta(\mathcal{U}'_p)$, the union $\mathcal{U}_p \cup \mathcal{U}'_p$ obeys also condition *Neighbourhood* (ii). This argumentation extends to the union of an arbitrary set of neighbourhoods obeying these conditions. Therefore, the union of all neighbourhoods obeying these conditions is the unique maximal $\mathcal{U}_p^+$. Obviously, the set $\mathcal{U}_p^- = \eta(\mathcal{U}_p^+)$ has the corresponding properties (with ∞^- interchanged by ∞^+). Arguments similar to those used above show that $\mathcal{U}_p^+ \cap \mathcal{U}_p^-$ has to be open and closed. Therefore, it has to be empty, since it does not contain $\infty^\pm$. **q.e.d.**



In the sequel we want to extend the open sets $\mathcal{U}_p^+$ and $\mathcal{U}_p^-$ to open disjoint sets, which are mapped by the corresponding branches of p biholomorphically onto the complement of finitely many finite parts of straight lines in $\mathbb{C} \subset \mathbb{P}^1$. The copy of $\mathbb{P}^1$ containing the image of $\mathcal{U}_p^+$ is denoted by $\mathbb{P}_p^+$ and the copy containing the image of $\mathcal{U}_p^-$ is denoted by $\mathbb{P}_p^-$. It will turn out that the Riemann surface Y may be realized by gluing $\mathbb{P}_p^+$ and $\mathbb{P}_p^-$ along several cuts, and the gluing map is always of the form $p \mapsto p$ plus some element of $\mathbb{R}$. We shall describe such surgeries by figures. On these figures small letters will denote the values of the functions p at the zeroes of dp . If a sheet of $\mathbb{P}_p^+$ is glued with some sheets of $\mathbb{P}_p^-$ these two sheets will be denoted by a capital letter with exponent $\pm$. If two sheets of $\mathbb{P}_p^+$ or $\mathbb{P}_p^-$ are these two sheets are denoted by a capital letter with indices 1 and 2 and exponent $\pm$. Let us first give three typical examples:

A horizontal cut. In this case dp has two simple zeroes at the values $p = a$ and $p = b$ with respect to the branch of p near ∞^+ and the values $p = -\bar{b}$ and $p = -\bar{a}$ with respect to the branch of p near ∞^- , respectively, where a and b are complex numbers with $\Im(a) = \Im(b)$ and $\Re(a) < \Re(b)$. The difference between the branch near ∞^+ minus the branch near ∞^- is therefore equal to $a + \bar{b} = b + \bar{a}$. Finally, the Riemann surface Y may be realized by gluing the slit along the straight line connecting a and b of $\mathbb{P}_p^+$ with the slit along the straight line connecting $-\bar{b}$ and $-\bar{a}$ of $\mathbb{P}_p^-$. More precisely, for the gluing we use the map $p \mapsto p + a + \bar{b}$, which transforms the branch near of $\mathbb{P}_p^+$ into the branch of $\mathbb{P}_p^-$.



figure of $\mathbb{P}_p^+$ figure of $\mathbb{P}_p^-$

A pair of parallel cuts. In this case dp has four simple zeroes at the values $p = a_1, a_2, b_1, b_2$ on $\mathbb{P}_p^+$, where $\Im(a_1) = \Im(a_2)$, $\Im(b_1) = \Im(b_2)$, $\Im(a_1) > \Im(b_1)$, $\Re(a_1) < \Re(a_2)$ and $\Re(b_1) - \Re(b_2) = \Re(a_1) - \Re(a_2)$. Also there are four other zeroes at the values $p = -\bar{a}_2, -\bar{a}_1, -\bar{b}_2, -\bar{b}_1$. In this case the slit along the straight lines of $\mathbb{P}_p^+$ connecting a_1 and b_1 is glued with the slit along straight lines connecting a_2 and b_2 , and the analog slits of $\mathbb{P}_p^-$ are also glued. Hence on $\mathbb{P}_p^+$ and on $\mathbb{P}_p^-$ two pairs of zeroes of dp are identified, and all together we have four zeroes of dp . Besides this two pairs of parallel slits we have two horizontal slits. One connects the two slits of $\mathbb{P}_p^+$ and the other connects the two slits of $\mathbb{P}_p^-$. Both are part of a line $\Im(p) = c$ with some arbitrary constant in the interval $\Im(a_1) \geq c \geq \Im(b_1)$. Since the two slits at the end of each of these horizontal slits are glued, these horizontal slits are glued to form circles. Finally, these two horizontal slits of $\mathbb{P}_p^+$ and $\mathbb{P}_p^-$ are glued by some map $p \mapsto p$ plus some real constant. But if we add to this constant the period $\Re(a_2) - \Re(a_1)$ the identification of the two circles does not change. Obviously we may shift these two slits (this means that we change c in the interval $\Im(a_1) \geq c \geq \Im(b_1)$) and simultaneously shift the constant of the gluing map in

some specific way, without changing the data $(Y, \infty^-, \infty^+, \eta, p)$ and the branch of p at ∞^+ .

figure of $\mathbb{P}_p^+$ figure of $\mathbb{P}_p^-$

A modified pair of parallel cuts. This is a variation of the *pairs of parallel cuts*, where the two sheets at the horizontal cut of $\mathbb{P}_p^+$ are glued by some map $p \mapsto p$ plus some real constant, and similarly for the two sheets at the horizontal cut of $\mathbb{P}_p^-$. In this case the Riemann surface Y has two connected components:

figure of $\mathbb{P}_p^+$ figure of $\mathbb{P}_p^-$

In the special case that the constant of the gluing map is zero, we could remove the two horizontal cuts. However, we will not do so, because otherwise k will not extend to a single-valued function on $\mathbb{P}_p^+$.

Lemma 3.16. *Let the data $(Y, \infty^-, \infty^+, p)$ be given by any data $(Y, \infty^-, \infty^+, k) \in \bar{\mathcal{M}}_{g,\eta}$ with $p = k_1$ (or any real component of k). Then for all branches of p in some neighbourhood of ∞^+ the Riemann surface Y may be realized by two copies of $\mathbb{P}^1$, which are glued along a combination of finitely many horizontal cuts and finitely many pairs of parallel cuts. The involution η is given by the map $p \mapsto -\bar{p}$ interchanging $\mathbb{P}_p^+$ and $\mathbb{P}_p^-$. Moreover, the first integral of the data $(Y, \infty^-, \infty^+, k)$ is equal to the Dirichlet integral of the positive form*

$$2\pi\sqrt{-1} \operatorname{vol}(\mathbb{R}^2/\Lambda) (dk_1 + \sqrt{-1}k_2) \wedge \overline{(dk_1 + \sqrt{-1}k_2)}$$

over $\mathbb{P}_p^+$. In particular, the first integral is always some non-negative real number. The first integral vanishes, if and only if the data $(Y, \infty^-, \infty^+, k)$ are isomorphic to the following data: Y is biholomorphic to the disjoint union of $\mathbb{P}_{k_1}^+ \cup \mathbb{P}_{k_1}^-$ with the natural marked points ∞^- and ∞^+ . The component k_1 is the natural meromorphic function on both copies of $\mathbb{P}_{k_1}$. The component k_2 is on $\mathbb{P}_{k_1}^+$ equal to $\sqrt{-1}k_1$ and on $\mathbb{P}_{k_1}^-$ equal to $-\sqrt{-1}k_1$. Finally, η is the involution induced by the map $k \mapsto -\bar{k}$, which interchanges the components $\mathbb{P}_{k_1}^+$ and $\mathbb{P}_{k_1}^-$.

Proof. **1.** First we claim that the convex compact sets excluded from the image of $\mathcal{U}_p^+$, mentioned in *Neighbourhood* (i), are the convex hulls of the values of p at finitely many zeroes of dp . For all $\mathcal{U}_p$ obeying conditions *Neighbourhood* (i)–(ii), the connected components of the boundary of $\mathcal{U}_p$ are mapped by p on convex Jordan curves around the excluded domains in $\mathbb{C} \subset \mathbb{P}_p^+$. Since p is locally biholomorphic if and only if dp has no zero, we may enlarge $\mathcal{U}_p$ and therefore shrink these Jordan curves, until they form the boundary of the convex hull of finitely many values of p at zeroes of dp . In case where the image of $\mathcal{U}_p^+$ is the complement of finitely many disjoint *horizontal cuts*, the Riemann surface Y is biholomorphic to the union of $\mathbb{P}_p^+$ and $\mathbb{P}_p^-$ glued in the prescribed way.

2. Now we claim that no straight line with constant imaginary part of p , which does not contain any zero of dp , can connect $\mathcal{U}_p^+$ and $\mathcal{U}_p^-$. Let $s_0 \in \mathbb{R}$ be the imaginary part of p at one zero of dp , and consider the family of all those straight lines, whose constant imaginary part belongs to a small neighbourhood of s_0 , and whose open ends, where the real part of p goes to $-\infty$, belong either to $\mathcal{U}_p^+$ or $\mathcal{U}_p^-$. Some of the zeroes of dp have the property, that if we move around them and return to the part with very small real parts of p we may connect $\mathcal{U}_p^+$ with $\mathcal{U}_p^-$. Now all zeroes of dp with this property, and which belong to one connected component of the complement of $\mathcal{U}_p^+ \cup \mathcal{U}_p^-$, have to be permuted by the involution η . Hence the number of such zeroes has to be even, and therefore these straight lines cannot connect $\mathcal{U}_p^+$ and $\mathcal{U}_p^-$.



3. Let us call a straight line with constant imaginary part of p periodic, if there exists some period $\Pi \in \mathbb{R}$, such that the preimages of $p + \Pi$ coincide with the preimages of p . If the imaginary part of p varies in some interval, such that the corresponding periodic lines contain no zero of dp , then the corresponding periods are constant, since dp is closed. Obviously, all straight lines with constant imaginary part of p are either periodic, or have open ends intersecting one of the two neighbourhoods $\mathcal{U}_p^+$ or $\mathcal{U}_p^-$ of the two poles of dp at $\infty^\pm$. Hence for generic real s (i. e. for s in the complement of the finite set of values of the imaginary part of p at all zeroes of dp) there exist exactly two non-periodic straight lines with constant imaginary part of p equals s , one having non-empty intersection with $\mathcal{U}_p^+$ and another one having non-empty intersection with $\mathcal{U}_p^-$. In particular, the Riemann surfaces decomposes into three types of regions. On one hand *regions of periodic lines* and on the other hand two regions of non-periodic lines intersecting either $\mathcal{U}^+$ or $\mathcal{U}^-$. Moreover, the common boundaries of different regions with periodic lines, are periodic lines, which contain a zero of dp . Furthermore, the common boundaries of regions with periodic lines with one of the two latter regions are of the same form. The common boundaries of the latter two regions, however, are *horizontal cuts*.

4. Now we claim that for all real s , the part of the complement of $\mathcal{U}_p^+ \cup \mathcal{U}_p^-$, on which the imaginary part of p is equal to s is a union of straight lines, whose sum of lengths is equal



to the sum of those segments of the straight lines with imaginary part equals s , which are excluded from $\mathcal{U}_p^+$ and $\mathcal{U}_p^-$. Obviously this is true for very large (and very small) values of s . On every component of a *region of periodic lines* the imaginary part of p takes the maximum and the minimum either on a common boundary with $\mathcal{U}^+$ or with $\mathcal{U}^-$. Hence the claim follows from the description of the boundaries of the three types of regions described in part 3. Moreover, the same arguments show that the complement of all *regions of periodic lines* is the disjoint union $\mathcal{U}^+ \cup \mathcal{U}^-$.

5. We have decomposed the Riemann surface Y into three regions: the neighbourhoods $\mathcal{U}_p^+$ and $\mathcal{U}_p^-$ (or their closures) and the *regions of periodic lines*. We have already seen that for any value of the imaginary part of p , the corresponding lines together form two copies of lines of $\mathbb{P}^1$, with constant imaginary part. Consequently we will now see that these parts may be glued in such a way, that they form the two copies $\mathbb{P}_p^+$ and $\mathbb{P}_p^-$. In doing so we have some freedom and no canonical choice. But, as we already have seen all connected components of the boundary of the closures of $\mathcal{U}_p^+$ and $\mathcal{U}_p^-$, which do not form a *horizontal cut*, are glued with some *regions of periodic lines*, and for each single *region of periodic lines* (with only one period Π) the corresponding gluing rule is of the form $p \mapsto p + \text{some real constant}$. This together with the condition that for all imaginary values of p the sum of the lengths of the corresponding periodic lines is exactly equal to the excluded parts of those lines with open ends intersecting $\mathcal{U}_p^+$ or $\mathcal{U}_p^-$ is the essential content of the statement that Y may be realized by a combination of several *horizontal cuts* and several *pair of parallel cuts*. In order to make this more precise, let us describe these combinations in some more detail.



6. If we try to realize Y in general by a combination of several *horizontal cuts* and several *pairs of parallel cuts* we have to envisage the possibility that different *pairs of parallel cuts* penetrate each other. In the following two figures we show two equivalent such possibilities, where all pairs of parallel slits, which differ by some real period, are glued:

figure of $\mathbb{P}_p^+$ figure of $\mathbb{P}_p^+$

In such cases we introduce additional horizontal cuts as shown in the following figures:

figure of $\mathbb{P}_p^+$ figure of $\mathbb{P}_p^-$

Furthermore, we permit the situation, where that the parallel cuts of *pair of parallel cuts* are two lines with some corners, and where one is a shift of the other by some real period. A *horizontal cut* and a *pair of parallel cuts* may also penetrate each other: Either the horizontal cut belongs to the *regions of periodic lines*, or to the part $\mathcal{U}_p^+ \cup \mathcal{U}_p^-$ where these lines are not periodic. In the following figures, we show two *pairs of parallel cuts* with corners and two *horizontal cuts*, one inside of the part of periodic lines and one outside.

figure of $\mathbb{P}_p^+$ figure of $\mathbb{P}_p^-$

As a result of the preceding considerations all Riemann surfaces Y may be realized as a combination of several *horizontal cuts* with several *pairs of parallel cuts* and some additional horizontal cuts connecting pairs of parallel cuts.

7. Let us assume now that we have given some data $(Y, \infty^-, \infty^+, k) \in \bar{\mathcal{M}}_{g,\eta}$, and consider the corresponding realization of Y as two planes $\mathbb{P}_p^\pm$ glued in the way described above. The positive two-form

$$2\pi\sqrt{-1} \operatorname{vol}(\mathbb{R}^2/\Lambda) (dk_1 + \sqrt{-1}dk_2) \wedge \overline{(dk_1 + \sqrt{-1}dk_2)}$$

over the part $\mathbb{P}_p^+$ is equal to

$$4\pi \operatorname{vol}(\mathbb{R}^2/\Lambda) (d\Re(k_1) \wedge d\Im(k_1) + d\Re(k_2) \wedge d\Im(k_2) + d\Re(k_1) \wedge d\Re(k_2) + d\Im(k_1) \wedge d\Im(k_2)).$$

On $\mathbb{P}_p^+$ the function k is single-valued, hence this integral is equal to the integral of the

one-form

$$4\pi \operatorname{vol}(\mathbb{R}^2/\Lambda) (\Re(k_1)d\Re(k_2) - \Im(k_2)d\Im(k_1) - \Im(k_1)d\Re(k_2) - \Im(k_2)d\Re(k_1))$$

over the boundary of $\mathbb{P}_p^+$. On the other hand, due to Lemma 3.12 the *first integral* is equal to the integral of the one-form

$$4\pi \operatorname{vol}(\mathbb{R}^2/\Lambda) (\Re(k_1)d\Re(k_2) + \Im(k_2)d\Im(k_1) + \sqrt{-1}\Im(k_1)d\Re(k_2) - \sqrt{-1}\Im(k_2)d\Re(k_1))$$

over the boundary of $\mathbb{P}_p^+$. Hence it would suffice to prove that the integral of the four one-forms $\Im(k_1)d\Re(k_1)$, $\Im(k_2)d\Re(k_2)$, $\Im(k_2)d\Im(k_1)$, $\Im(k_1)d\Re(k_2)$ and $\Im(k_2)d\Re(k_1)$ over the boundary of $\mathbb{P}_p^+$ all vanish. Let us first prove this statement for the two basic cases of a *horizontal cut* or a *pair of parallel cuts*. In the case of a *horizontal cut* the boundary of $\mathbb{P}_p^+$ is a cycle of Y , which is invariant under the involution η . The transformation of the function k under this involution implies that in this case the integrals of the one-forms $\Im(k_1)d\Re(k_1)$, $\Im(k_2)d\Re(k_2)$, $\Im(k_1)d\Re(k_2)$ and $\Im(k_2)d\Re(k_1)$ over the boundary of $\mathbb{P}_p^+$ vanish. Since on a horizontal cut the form $d\Im(k_1)$ vanishes, the integral $\Im(k_2)d\Im(k_1)$ over the boundary of $\mathbb{P}_p^+$ also vanishes. If we consider a *pair of parallel cuts* we shall first remark that all forms $\Im(k_1)d\Re(k_1)$, $\Im(k_2)d\Re(k_2)$, $\Im(k_2)d\Im(k_1)$, $\Im(k_1)d\Re(k_2)$ and $\Im(k_2)d\Re(k_1)$ are invariant under a translation of (k_1, k_2) by any element of $\mathbb{R}^2$. This implies that all the integrals along the parallel cuts vanish. Moreover, this argument also extends to a horizontal cut connecting two parallel cuts, which cuts two parts of $\mathbb{P}_p^+$ into different parts. If the horizontal cut, which connects both parallel cuts, then it is a boundary of $\mathbb{P}_p^+$ and $\mathbb{P}_p^-$, it is the sum of two circles of Y , which are interchanged by η . The same arguments as in the case of a *horizontal cut* show that also in this case all the integrals of the forms $\Im(k_1)d\Re(k_1)$, $\Im(k_2)d\Re(k_2)$, $\Im(k_2)d\Im(k_1)$, $\Im(k_1)d\Re(k_2)$ and $\Im(k_2)d\Re(k_1)$ along the horizontal cut connecting the two parallel cuts vanish. These arguments obviously extend to the combination of several *horizontal cuts* with several *pair of parallel cuts* of the general case. Moreover, these arguments also show, that the *first integral* is equal to the integral of the positive form $2\pi\sqrt{-1}\operatorname{vol}(\mathbb{R}^2/\Lambda)(dk_1 + \sqrt{-1}dk_2) \wedge (dk_1 + \sqrt{-1}dk_2)$ over $\mathcal{U}_p^+$ plus the integral of the positive form $2\pi\sqrt{-1}\operatorname{vol}(\mathbb{R}^2/\Lambda)(dk_1 \wedge d\bar{k}_1 + dk_2 \wedge d\bar{k}_2)$ over the *subregion of periodic lines* of $\mathbb{P}_p^+$. The second summand is larger or equal to $4\pi \operatorname{vol}(\mathbb{R}^2/\Lambda)$ times the volume of this *subregion of periodic lines* of $\mathbb{P}_p^+$. In particular, the *first integral* is larger or equal to the volume of this part.

8. We conclude that the *first integral* is a non-negative real number. It is zero, if and only if dk_2/dk_1 is identically equal to $\sqrt{-1}$ on $\mathbb{P}_{k_1}^+$. In this case Y has two connected components, on one dk_2/dk_1 is equal to $\sqrt{-1}$ and on the other equal to $-\sqrt{-1}$. Since all periods of dk_1 and dk_2 are real, this implies that they have to be zero in this case. Hence k_1 and k_2 may be chosen to be single-valued meromorphic functions with one pole on each connected component of Y . Therefore, both connected components have to be biholomorphic to $\mathbb{P}^1$. This proves the Lemma. **q.e.d.**

For all data $(Y, \infty^-, \infty^+, \eta, p)$ let $D(Y, \infty^-, \infty^+, \eta, p)$ denote the maximal diameters of the compact convex subsets mentioned in *Neighbourhood* (i), which form the complement of the image of $\mathcal{U}_p^+$. Since the diameter is invariant under translations, it depends only on the form dp .

Lemma 3.17. *For all $g \in \mathbb{N}_0$ and $\mathcal{W} > 0$ there exists some $C > 0$, such that for all $(Y, \infty^-, \infty^+, \eta, k) \in \bar{\mathcal{M}}_{g, \Lambda, \eta, \mathcal{W}}$ the diameters $D(Y, \infty^-, \infty^+, \eta, \hat{p})$ and $D(Y, \infty^-, \infty^+, \eta, \check{p})$ are not larger than C .*

Proof. Let us assume on the contrary that there exists some sequence of data in $\bar{\mathcal{M}}_{g, \Lambda, \eta, \mathcal{W}}$, such that either the diameter of the corresponding data $(Y, \infty^-, \infty^+, \eta, \hat{p})$ or the diameter of the data $(Y, \infty^-, \infty^+, \eta, \check{p})$ tends to infinity. By passing to some subsequence we may assume that for all elements of the subsequence the second diameter is larger than the former diameter and that the second sequence of diameters is monotone increasing. We *renormalize* the functions k and obtain some sequence of data in $\bar{\mathcal{M}}_{g, \eta}$, whose corresponding data $(Y, \infty^-, \infty^+, \eta, \check{p})$ have constant diameter equal to 1, and whose *first integral* converges to zero. Some of the excluded domains of $\mathcal{U}_{\check{p}}^+$ corresponding to the sequence of data $(Y, \infty^-, \infty^+, \eta, \check{p})$ tend to infinity. Now we use the invariance under the shifts described in Lemma 2.11 to transform the sequence into another sequence in $\bar{\mathcal{M}}_{g, \Lambda, \eta, \mathcal{W}}$ of data, such that the *renormalized* data have excluded domains of $\mathcal{U}_{\check{p}}^+$ corresponding to the *renormalized* $(Y, \infty^-, \infty^+, \eta, \check{p})$ inside of some finite subset of $\mathbb{C}$ with diameter equal to 1.

In the proof of Lemma 3.16 we showed that the *first integral* is larger or equal to $4\pi \text{vol}(\mathbb{R}^2/\Lambda)$ times the volume of the *regions of periodic lines* of $\mathbb{P}_{\hat{p}}^+$. Since the *first integral* is invariant under conformal linear transformations of Λ , in general the *first integral* is larger or equal to $4\pi \text{vol}(\mathbb{R}^2/\Lambda)/g(\gamma, \gamma)$ times the volume of the *regions of periodic lines* of $\mathbb{P}_{\check{p}}^+$ with respect to the component $p = g(\gamma, k)$. Since the periods of $\hat{p}$ and $\check{p}$ are integers, we conclude that the *regions of periodic lines* of $\mathbb{P}_{\hat{p}}^+$ and $\mathbb{P}_{\check{p}}^+$ of the *renormalized* sequence converge to finitely many horizontal lines, respectively. Now we claim that on all compact subsets in the complement of these finitely many horizontal lines of $\mathbb{P}_{\hat{p}}$ the function $k_1 + \sqrt{-1}k_2$ converges uniformly to zero. In fact, due to Cauchy's formula [Co-I, Chapter IV. §6.1] the value of this function at any element $\hat{p}'$ in the image of $\mathcal{U}_{\check{p}}^+$ under the function $\hat{p}$ is equal to

$$\frac{1}{2\pi\sqrt{-1}} \int_{\partial\mathcal{U}_{\check{p}}^+} (k_1 + \sqrt{-1}k_2) \frac{d\hat{p}}{\hat{p} - \hat{p}'} = -\frac{1}{2\pi\sqrt{-1}} \int_{\partial\mathcal{U}_{\check{p}}^+} \ln(1 - \hat{p}/\hat{p}') (dk_1 + \sqrt{-1}dk_2).$$

We remark that this formula is only valid, if $\hat{p}'$ is not contained inside of any of the cycles corresponding to the component of the boundary of $\mathcal{U}_{\check{p}}^+$ (in other words the formula is valid, if there exists a path not crossing the boundary of $\mathcal{U}_{\check{p}}^+$, which connects $\hat{p}'$ with ∞^+). This formula yields the branch of the function $k_1 + \sqrt{-1}k_2$, which vanishes at ∞^+ . In particular, if we use the left hand side to calculate the contribution of some component of the boundary of $\mathcal{U}_{\check{p}}^+$, this contribution does not depend on some constant, which is added to the function $k_1 + \sqrt{-1}k_2$ nearby this boundary component. In fact, due to Lemma 3.16 the function $k_1 + \sqrt{-1}k_2$ extends to some single-valued function on $\mathcal{U}_{\check{p}}^+$. On the other hand we may estimate $|(dk_1 + \sqrt{-1}dk_2)/d\hat{p}|$ at every point $\hat{p}'$ in the image of $\mathcal{U}_{\check{p}}^+$ under $\hat{p}$, whose distance to the boundary of $\mathcal{U}_{\check{p}}^+$ is larger than some $\varepsilon > 0$ by some constant (depending only on the lattice) times $\mathcal{W}/\varepsilon$. We conclude that the function $k_1 + \sqrt{-1}k_2$ converges to zero uniformly on compact subsets of the complement of the horizontal straight lines.

We conclude that all accumulation points of zeroes of $d\check{p}$ cannot belong to the interior of $\mathbb{P}_{\check{p}}^+$ corresponding to the limit. Hence also the maximal difference of the real parts of the domains excluded from $U_{\check{p}}^+$ corresponding to the sequences $(Y, \infty^-, \infty^+, \eta, \check{p})$ converges to zero, which contradicts the assumptions. Therefore, the diameters of the elements of $\bar{\mathcal{M}}_{g,\Lambda,\eta,\mathcal{W}}$ are bounded. **q.e.d.**

Lemma 3.18. *For all fixed $g \in \mathbb{N}_0$, $C > 0$ and $\mathcal{W} > 0$ and any sequence of elements $(Y, \infty^-, \infty^+, k) \in \bar{\mathcal{M}}_{g,\Lambda,\eta,\mathcal{W}}$ whose diameters of the corresponding sequences of members $D(Y, \infty^-, \infty^+, \eta, \hat{p})$ and $D(Y, \infty^-, \infty^+, \eta, \check{p})$ are not larger than C , there exists some subsequence and some choice of the corresponding cuts of $\mathbb{P}_{\hat{p}}^+ \setminus \{\infty^+\}$ described in Lemma 3.16, which converges in the sense of [Co-II, §15.4 Definition 4.1] to some choice of $\mathbb{P}_{\check{p}}^+ \setminus \{\infty^+\}$ of some data $(Y, \infty^-, \infty^+, \eta, \hat{p})$. Moreover, the function $\check{p}(\hat{p})$ of this subsequence converges uniformly on compact subsets to the corresponding functions of the limit. Finally, the first integral of the limit is not larger than the $\liminf$ of the subsequence of first integrals.*

Proof. Let us choose some constant $C > 0$ and consider all data $(Y, \infty^-, \infty^+, \eta, k) \in \bar{\mathcal{M}}_{g,\Lambda,\eta,\mathcal{W}}$ with the property $D(Y, \infty^-, \infty^+, \eta, \hat{p}) \leq C$. Let us first prove that any sequence of such data has a subsequence and some choice of cuts described in Lemma 3.16 with the property, that the corresponding domains $\mathbb{P}_{\hat{p}}^+ \setminus \{\infty^+\}$ converge in the sense of [Co-II, §15.4 Definition 4.1] to some choice of cuts corresponding to some data in this space. Obviously we may choose some subsequence such that the images of the sets $\mathcal{U}_{\hat{p}}^+ \setminus \{\infty^+\}$ under some choice of the branches of the function $\hat{p}$ converge to some open domain in $\mathbb{C}$. Moreover, by passing to some subsequence we may achieve that the imaginary values of $\hat{p}$ at all zeroes of $d\hat{p}$ converge and moreover that if s is not the limit of some sequence of imaginary parts of zeroes of $d\hat{p}$, that all periods of periodic straight lines with imaginary part equal to s converge to some finite set of periods. By passing again to some subsequence we may achieve in addition that the corresponding sequence of images of $\mathcal{U}_{\hat{p}}^+$ under $\hat{p}$ converge to the corresponding image of some data in this set. From the considerations in the proof of Lemma 3.16 it follows that we may chose the cuts of these subsequence in such a way, that the corresponding domains $\mathbb{P}_{\hat{p}}^+ \setminus \{\infty^+\}$ converge in the sense of [Co-II, §15.4 Definition 4.1] to some choice of cuts corresponding to some data in this space.

Now we claim that there exists some constant $C' > 0$ such that the mapping $\hat{p} \mapsto \check{p}$ defines some biholomorphic mapping from the subset $\{\hat{p} \in \mathbb{C} \mid B(\hat{p}, C') \subset \hat{p}[\mathcal{U}_{\hat{p}}^+]\}$ into some domain of $\mathbb{C}$. This follows from the formula for the function $k_1 + \sqrt{-1}k_2$ used in the proof of Lemma 3.17. We conclude that this biholomorphic mapping $\hat{p} \mapsto \check{p}$ maps some complement in $\mathbb{C}$ of finitely many compact domains into a complement of finitely many compact domains. More generally, the *complex Fermi curve* is therefore the union of finitely many compact pieces and two open ends, which are mapped by $\hat{p}$ and $\check{p}$ onto the complement in $\mathbb{C}$ of finitely many disjoint compact subsets. If we consider the *complex Fermi curve* as a finite-sheeted covering with respect to the mapping $\check{p}$, then we may also apply Lemma 3.16 to this covering. We conclude that the four functions $\exp(2\pi\sqrt{-1}\hat{p})$, $\exp(-2\pi\sqrt{-1}\hat{p})$, $\exp(2\pi\sqrt{-1}\check{p})$ and $\exp(-2\pi\sqrt{-1}\check{p})$ are bounded by some constant. Moreover, this argument carries over to some subsequence of any sequence in $\bar{\mathcal{M}}_{g,\Lambda,\eta,\mathcal{W}}$ with the properties described in the Lemma.

With the help of Montel's theorem [Co-I, Chapter VII §2.9] we conclude that there exists some subsequence, such that the functions $\check{p}(\hat{p})$ converge uniformly on compact subsets of $\mathbb{P}_{\hat{p}}^+ \setminus \{\infty^+\}$ to some function, which has to be equal to the corresponding function of the limit of the data.

The single-valued function dk_2/dk_1 has no poles on the interior of $\mathbb{P}_{\hat{p}}^+$ and is equal to $\sqrt{-1}$ at ∞^+ . We conclude that for all data in $\bar{\mathcal{M}}_{g,\eta}$ the function $(dk_2/dk_1 - \sqrt{-1})$ converges to zero, if the absolute value of $\hat{p} \in \mathbb{P}_{\hat{p}}^+$ converges to infinity. Since $dk_1 + \sqrt{-1}dk_2 = \sqrt{-1}((dk_2/dk_1 - \sqrt{-1})dk_1)$ we conclude that for all data in $\bar{\mathcal{M}}_{g,\eta}$ and all $\varepsilon > 0$ there exists some compact subset K of some choice of $\mathbb{P}_{\hat{p}}^+ \setminus \{\infty^+\}$ corresponding to this data, such that the integral of the form

$$2\pi\sqrt{-1} \operatorname{vol}(\mathbb{R}^2/\Lambda)(dk_1 + \sqrt{-1}dk_2) \wedge \overline{(dk_1 + \sqrt{-1}dk_2)}$$

over K is an element of the interval $[\mathcal{W} - \varepsilon, \mathcal{W}]$, where $\mathcal{W}$ denotes the *first integral* of the data. Since $\check{p}(\hat{p})$ converges uniformly on compact subsets of $\mathbb{P}_{\hat{p}}^+$, we conclude that the *first integral* of the limit is smaller than the sum of any accumulation point of the sequence of values of the *first integral* plus ε . Since this is true for all ε , the *first integral* of the limit has to be smaller or equal to the $\liminf$ of the sequence of values of the *first integral*. **q.e.d.**

It is possible to prove a sharper statement: For a given sequence of data some of the excluded domains of corresponding images of $\mathcal{U}_p^+$ under p , will tend to infinity. To all these regions we may associate a part of the *first integral*, by integrating the corresponding one-form around these excluded domains. In the limit the excluded domains, which tend to infinity are removed, and therefore the *first integral* of the limit is the limit of those parts of the *first integrals*, which correspond to excluded domains inside of bounded domains. In general, at least if the excluded domains have enough distance from each other, all parts of the *first integral* are positive. Therefore, the limit of the *first integral* is smaller than all accumulation points of the sequence of values of the *first integrals* of any convergent series of data.

Proof of Theorem 3.13. Due to Lemma 3.10 the space of all closed subsets of $\mathbb{C}^2$ is a separable metrizable space. Hence $\bar{\mathcal{M}}_{g,\Lambda,\mathcal{W}}$ is compact if and only if any sequence has a convergent subsequence. Due to the Lemma 3.18 and Lemma 3.17 any sequence of $\bar{\mathcal{M}}_{g,\Lambda,\mathcal{W}}$ has a subsequence, which converges in sense described above. It is easy to see that that this implies that the corresponding subsets $\mathcal{F}(Y, \infty^-, \infty^+, k) \subset \mathbb{C}^2$ converges. Moreover, due to Lemma 3.18 the *first integral* of the limit is not larger than $\mathcal{W}$. **q.e.d.**

3.4 Limits of complex Fermi curves

In this section we shall investigate the limits of the sequences of resolvents corresponding to weakly convergent sequences of complex potentials $(U_n)_{n \in \mathbb{N}}$ in $L^2(\mathbb{R}^2/\Lambda)$ with bounded L^2 -norm.

Due to the Banach–Alaoglu theorem [R-S-I, Theorem IV.21] and the Riesz Representation theorem [Ro-2, Chapter 13 Section 5] any sequence $(U_n)_{n \in \mathbb{N}}$ in $L^2(\mathbb{R}^2/\Lambda)$ with bounded L^2 -norm has a subsequence, with the property that the corresponding sequence of measures $U_n \bar{U}_n d^2x$ converge weakly to a measure on $\mathbb{R}^2/\Lambda$. With the help of Lemma 2.6 we conclude

that the corresponding sequence of resolvents converge, if the limit of the measure does not contain point measures of mass larger or equal to the constant S_p^{-2} . In fact, if the weak limit of the measures does not contain point measures of mass larger or equal to S_p^{-2} , we may cover $\mathbb{R}^2/\Lambda$ by open sets, whose measures with respect to the limit of the measures is smaller than S_p^{-2} . Due to the compactness of $\mathbb{R}^2/\Lambda$ this open covering has a finite subcovering. For any finite open covering, the function on $\mathbb{R}^2/\Lambda$, which associates to each x the radius of the maximal open disc around x , which is entirely contained in one member of the covering, is continuous. We exclude the trivial case, where one member of the covering contains the whole torus and therefore all discs. So this function is the maximum of the distances of the corresponding point to all complements of the members of the covering. Therefore, there exists a small $\varepsilon > 0$, such that all discs with radius 2ε are contained in one member of the finite subcovering. Obviously, for any member of the subcovering there exists a continuous $[0, 1]$ -valued function, whose support is contained in this member of the subcovering, and which is equal to 1 on those discs $B(x, \varepsilon)$, whose extensions $B(x, 2\varepsilon)$ are contained in this member of the subcovering. Since the sequence of measures converges weakly, the integrals of these functions with respect to the measures $U_n(x)\bar{U}_n(x)d^2x$ corresponding to the sequence are also smaller than S_p^{-2} , with the exception of finitely many elements of the sequence. This shows that with the exception of finitely many elements of the sequence, the norms $\|U_n\|_{2,\varepsilon}$ are smaller than some $C_p < S_p^{-1}$.

Lemma 3.19. *If a weak limit of the sequence of finite Baire measures $U_n\bar{U}_nd^2x$ on $\mathbb{R}/\Lambda$ does not contain point measures with mass larger or equal to S_p^{-2} , then there exists a $C_p < S_p^{-1}$, an $\varepsilon > 0$ and a subsequence of the bounded sequence U_n in $L^2(\mathbb{R}^2/\Lambda)$ whose norms $\|\cdot\|_{2,\varepsilon}$ are smaller than C_p .* **q.e.d.**

Since the L^2 -norm of the sequence is bounded, there exist only finitely many points $z_1, \dots, z_L$ in $\mathbb{C}/\Lambda_{\mathbb{C}}$, whose mass with respect to the weak limit of the measures $U_n(x)\bar{U}_n(x)d^2x$ is larger or equal to S_p^{-2} . Here we used the complex coordinates z instead of the real coordinates x . Moreover, since we consider only local potentials (i. e. multiplication by functions) the limits of the corresponding sequence of resolvents should yield perturbations of the resolvents of the Dirac operator corresponding to the weak limit U . Furthermore, we expect that the support of these perturbations is contained in this finite set $\{z_1, \dots, z_L\}$. In the first subsection we characterize the natural candidates for these perturbations. In the last subsection we will show that this is indeed the case. The proof is intricate and will proceed through several steps.

3.4.1 Finite rank perturbations

It will turn out that in general the limits of the resolvents $R(U_n, \bar{U}_n, k, \lambda)$ depend non-continuously on λ . Since we are interested in the limits of the *complex Fermi curves* we shall concentrate on the limits of $R(U_n, \bar{U}_n, k, 0)$. As a preparation we first derive an explicit formula for the integral kernels of $R(0, 0, k, 0)$.

These integral kernels may be expressed in terms of a Theta function corresponding to the elliptic curve $\mathbb{C}/\Lambda_{\mathbb{C}}$ together with a canonical homology bases. The choice of a *Fundamental*

domain 2.1 determines these data, therefore we decorate the corresponding Theta function with an index Δ . Let θ_Δ denote the following Theta function:

$$\theta_\Delta(z) = \frac{\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2}{\theta'_1(0, \frac{\check{\gamma}_1 + \sqrt{-1}\check{\gamma}_2}{\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2})} \theta_1 \left(\frac{z}{\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2}, \frac{\check{\gamma}_1 + \sqrt{-1}\check{\gamma}_2}{\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2} \right).$$

Here $\theta_1(z, \tau)$ is the first one-dimensional Theta-function with modulus τ [Bat, 13.19]. Since the orientation of the basis $\hat{\gamma}$ and $\check{\gamma}$ is positive, the imaginary part of the modulus $\frac{\check{\gamma}_1 + \sqrt{-1}\check{\gamma}_2}{\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2}$ is positive and so the one-dimensional Theta-functions are well defined. The properties of these functions implies that the functions θ_Δ are uniquely determined by the following properties:

Quasi-periodicity: Under the shifts of the elements of Λ considered as element in $\mathbb{C}$ this function transform as follows:

$$\begin{aligned} \theta_\Delta(z + \hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2) &= -\theta_\Delta(z) \\ \theta_\Delta(z + \check{\gamma}_1 + \sqrt{-1}\check{\gamma}_2) &= -\exp \left(-\pi\sqrt{-1} \frac{2z + \check{\gamma}_1 + \sqrt{-1}\check{\gamma}_2}{\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2} \right) \theta_\Delta(z). \end{aligned}$$

Holomorphicity: The function $z \mapsto \theta_\Delta(z)$ is a holomorphic entire function.

Anti-symmetry: Under the map $z \mapsto -z$ this function is anti-symmetric.

Zeroes: The zeroes of this function are exactly the elements of $\Lambda_{\mathbb{C}}$. More precisely, at the origin it has an expansion of the form $\theta_\Delta(z) = z + O(z^3)$.

Now Dolbeault's Lemma [Gu-Ro, Chapter I Section D 2. Lemma] implies that the inverse of the operators $\bar{\partial}_{[k]}$ and $\partial_{[k]}$ have the following integral kernels:

$$\begin{aligned} \mathcal{K}_1(z - z', k) \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}} &= \exp \left(2\pi\sqrt{-1}g(\hat{\gamma}, k) \frac{z - z'}{\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2} \right) \frac{\theta_\Delta(z - z' + z^+(k))}{\theta_\Delta(z - z')\theta_\Delta(z^+(k))} \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}} \\ \mathcal{K}_2(z - z', k) \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}} &= \exp \left(2\pi\sqrt{-1}g(\hat{\gamma}, k) \frac{\bar{z} - \bar{z}'}{\hat{\gamma}_1 - \sqrt{-1}\hat{\gamma}_2} \right) \frac{\bar{\theta}_\Delta(z - z' + \bar{z}^-(k))}{\bar{\theta}_\Delta(z - z')\bar{\theta}_\Delta(\bar{z}^-(k))} \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}}, \end{aligned}$$

where $z^+(k) = \text{vol}(\mathbb{R}^2/\Lambda)(\sqrt{-1}k_1 - k_2)$ and $z^-(k) = \text{vol}(\mathbb{R}^2/\Lambda)(\sqrt{-1}k_1 + k_2)$. In fact, this choice implies the relations

$$\begin{aligned} \exp \left(2\pi\sqrt{-1}g(\hat{\gamma}, k) \frac{\check{\gamma}_1 + \sqrt{-1}\check{\gamma}_2}{\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2} \right) \exp \left(-\pi\sqrt{-1} \frac{2z^+(k)}{\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2} \right) &= \exp(2\pi\sqrt{-1}g(\check{\gamma}, k)) \\ \exp \left(2\pi\sqrt{-1}g(\hat{\gamma}, k) \frac{\check{\gamma}_1 - \sqrt{-1}\check{\gamma}_2}{\hat{\gamma}_1 - \sqrt{-1}\hat{\gamma}_2} \right) \exp \left(\pi\sqrt{-1} \frac{2z^-(k)}{\hat{\gamma}_1 - \sqrt{-1}\hat{\gamma}_2} \right) &= \exp(2\pi\sqrt{-1}g(\check{\gamma}, k)). \end{aligned}$$

Hence both integral kernels define with respect to z sections of the line bundle corresponding to $[k]$ and with respect to z' sections of the line bundle corresponding to $[-k]$. The first integral kernel is well defined if and only $z^+(k)$ is not an element of $\Lambda_{\mathbb{C}}$ and the second

integral kernel is well defined if and only if $\bar{z}^-(k)$ is not an element of $\Lambda_{\mathbb{C}}$. The generators of the dual lattice are equal to $\hat{\kappa} = J\tilde{\gamma}/\text{vol}(\mathbb{R}^2/\Lambda)$ and $\check{\kappa} = J\hat{\gamma}/\text{vol}(\mathbb{R}^2/\Lambda)$. Therefore, the functions $z^{\pm}(k)$ obey the relations

$$\begin{aligned} z^+(k + \hat{\kappa}) &= z^+(k) + \tilde{\gamma}_1 + \sqrt{-1}\tilde{\gamma}_2 & z^+(k + \check{\kappa}) &= z^+(k) + \hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2 \\ z^-(k + \hat{\kappa}) &= z^-(k) - \overline{\tilde{\gamma}_1 + \sqrt{-1}\tilde{\gamma}_2} & z^-(k + \check{\kappa}) &= z^-(k) - \overline{\hat{\gamma}_1 + \sqrt{-1}\hat{\gamma}_2}. \end{aligned}$$

Hence the first condition is equivalent to the condition that the line bundle corresponding to $[k]$ has a nontrivial holomorphic section and $\bar{\partial}_{[k]}$ has a kernel and the second condition is equivalent to the condition that this line bundle has a anti-holomorphic section and $\partial_{[k]}$ has a kernel. More generally, these integrals kernels considered as functions of k are invariant under shifts of the dual lattice, and therefore depend only on $[k] \in \mathbb{C}^2/\Lambda^*$. We collect these calculations in the following

Lemma 3.20. *The resolvent $R(0, 0, k, 0)$ has the following integral kernel:*

$$\begin{pmatrix} 0 & \mathcal{K}_1(z - z', k) \\ -\mathcal{K}_2(z - z', k) & 0 \end{pmatrix} \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}} = \begin{pmatrix} 0 & \mathcal{K}_1(z - z', k) \\ -\mathcal{K}_2(z - z', k) & 0 \end{pmatrix} \frac{d^2x'}{\pi}.$$

q.e.d.

The following properties characterize the natural candidates of those perturbations, which occur as limits of sequences of resolvents.

Finite rank Perturbation (i) The resolvents fit together to a meromorphic family of operators depending on $k \in \mathbb{C}^2$, which differs from the family of resolvents $k \mapsto R(V, W, k, 0)$ by a family of finite rank operators.

Finite rank Perturbation (ii) The application of the Dirac operators $D(V, W, k)$ on all elements of the range of the finite rank operators in (i) are tempered distributions, whose support is contained in a finite set $\{z_1, \dots, z_L\}$. The same holds for the corresponding transposed operators. The support of these distributions is called the support of the *Finite rank Perturbation*.

Finite rank Perturbation (iii) The corresponding perturbations of the Dirac operators $D(V, W, k)$ does not depend on k .

Since any tempered distribution, which is located in finitely many points, is a sum of derivatives of δ -functions [R-S-I, Theorem V.11] condition *Finite rank Perturbation* (ii) implies that the difference of the resolvents minus $R(V, W, k, 0)$ is an operator with integral kernel

$$\sum_{m,n} \psi_m(z, k) S(k)_{m,n} \phi_n(z', k) d^2x',$$

where $S(k)$ is a $L \times L$ -matrix-valued function depending on $[k] \in \mathbb{C}^2/\Lambda^*$, and $\psi_1, \dots, \psi_K$ and $\phi_1, \dots, \phi_K$ have the property that the applications of $D(V, W, k)$ and $D^t(V, W, k)$ yields

derivatives of δ -functions, respectively. Therefore, these functions are values of the derivatives of the integral kernel at the support of the *Finite rank Perturbation*. In the following ten steps we will make this more precise. The first three steps can be considered as an application of Krein's formula [A-K]. However, these perturbations violate the general rule, that the domain of the perturbation should contain the domain of the unperturbed operator ([Ka, Chapter four] and [A-K, Chapter 1]).

1. Let us first characterize all *Finite rank Perturbations* of $R(0, 0, k, 0)$, whose support is just one point z_l . All these perturbations have integral kernels of the form

$$\Psi(z, k)S(k)\Phi(z', k)d^2x',$$

where $S(k)$ is a $M \times M$ matrix depending on $k \in \mathbb{C}^2$, whose entries are 2×2 -matrices, and $\Psi(z, k)$ and $\Phi(z, k)$ are the following row-vectors and column-vectors with M entries indexed by $m = 0, \dots, M-1$, respectively:

$$\Psi_m(z, k) = \frac{(-1)^m}{\pi m!} \begin{pmatrix} 0 & \mathcal{K}_1^{(m)}(z-z_l, k) \\ -\mathcal{K}_2^{(m)}(z-z_l, k) & 0 \end{pmatrix} \quad \Phi_m(z, k) = \frac{1}{\pi m!} \begin{pmatrix} 0 & \mathcal{K}_1^{(m)}(z_l-z, k) \\ -\mathcal{K}_2^{(m)}(z_l-z, k) & 0 \end{pmatrix},$$

where $\mathcal{K}_1^{(m)}(z, k)$ and $\mathcal{K}_2^{(m)}(z, k)$ denotes the m -th holomorphic and anti-holomorphic derivative with respect to z of the functions $\mathcal{K}_1$ and $\mathcal{K}_2$, respectively. This ansatz should be understood as the 2×2 matrix, which is obtained by the matrix multiplication of the transposed vector Ψ with L entries times the $L \times L$ matrix $S(k)$ times the vector Φ with again L entries, all whose entries are 2×2 matrices. Formally it would be also possible to consider anti-holomorphic derivatives of $\mathcal{K}_1$ or holomorphic derivatives of $\mathcal{K}_2$. But these perturbations have integral kernels, which are supported only at the points $z_1, \dots, z_L$. We shall not consider such highly singular perturbations, since we are only interested in perturbations of the resolvents, which can be obtained as the limit $\varepsilon \downarrow 0$ of perturbations, whose integral kernel are supported in the complement of $B(z_1, \varepsilon) \cup \dots \cup B(z_L, \varepsilon)$.

Formally the corresponding perturbations P of $D(0, 0, k, 0)$ should have integral kernels of the form

$$\check{\Psi}(z)P\check{\Phi}(z')d^2x' \quad \text{with} \quad \check{\Psi}_m = \frac{(-1)^m}{m!} \begin{pmatrix} \bar{\partial}^m \delta(z-z_l) & 0 \\ 0 & \partial^m \delta(z-z_l) \end{pmatrix} \quad \check{\Phi}_m = \frac{(-1)^m}{m!} \begin{pmatrix} \partial^m \delta(z-z_l) & 0 \\ 0 & \bar{\partial}^m \delta(z-z_l) \end{pmatrix}.$$

Again $\check{\Psi}$ and $\check{\Phi}$ are row-vectors and column-vectors, respectively, with the given entries indexed by $m = 0, \dots, M-1$. They are related to Ψ and Φ :

$$\Psi(z, k) = R(0, 0, k, 0)\check{\Psi}(z) \quad \Phi^t(z, k) = R^t(0, 0, k, 0)\check{\Phi}^t(z).$$

Since the resolvent of the perturbed operator is equal to

$$R(0, 0, k, 0)(\mathbf{1} - PR(0, 0, k, 0))^{-1},$$

we should calculate the inverse operator on the right hand side.

Lemma 3.21. *Let F be a Banach subspace of the Banach space E and A a linear operator on E , whose range is contained in F . Then the operator $\mathbf{1}_E - A$ on E has an inverse if and only if the operator $\mathbf{1}_F - A_F$ on F has an inverse, where A_F denotes the restriction of A to F considered as an operator on F . Finally, if the inverse of the latter exists, then the inverse of the former is equal to*

$$(\mathbf{1}_E - A)^{-1} = \mathbf{1}_E + (\mathbf{1}_F - A_F)^{-1} A.$$

In order to simplify notation we do not distinguish between operators with the same range, but which are mappings to different Banach spaces. Therefore, A denotes either an operator on E or an operator from E to F . Furthermore, the operators $\mathbf{1}_F$ and A_F denote either operators on F or operators from F to E .

Proof. Since the range of A is contained in F , the operator $\mathbf{1}_E - A$ leaves the subspace F invariant. If the inverse of $\mathbf{1}_E - A$ exists, then it also leaves invariant the subspace F , and the restriction to the subspace F of the inverse is an inverse of the operator $\mathbf{1}_F - A_F$. Conversely, if the inverse of the latter operator exists, then the operator $\mathbf{1}_E + (\mathbf{1}_F - A_F)^{-1} A$ is a right inverse of $\mathbf{1}_E - A$:

$$(\mathbf{1}_E - A) (\mathbf{1}_E + (\mathbf{1}_F - A_F)^{-1} A) = \mathbf{1}_E - A + (\mathbf{1}_F - A_F) (\mathbf{1}_F - A_F)^{-1} A = \mathbf{1}_E.$$

Moreover, since the range of A is contained in F , the operator $A(\mathbf{1}_E - A)$ on E is equal to the operator $(\mathbf{1}_F - A_F) A$. Therefore, the operator $\mathbf{1}_E + (\mathbf{1}_F - A_F)^{-1} A$ is also a left inverse of $\mathbf{1}_E - A$:

$$(\mathbf{1}_E + (\mathbf{1}_F - A_F)^{-1} A) (\mathbf{1}_E - A) = \mathbf{1}_E - A + (\mathbf{1}_F - A_F)^{-1} (\mathbf{1}_F - A_F) A = \mathbf{1}_E.$$

q.e.d.

In order to apply this Lemma the integrals $\int \partial^n \delta(z - z_l) \mathcal{K}_1^{(m)}(z - z_l, k) d^2x$ and $\int \bar{\partial}^n \delta(z - z_l) \mathcal{K}_2^{(m)}(z - z_l, k) d^2x$ have to be evaluated. In doing so we use in some neighbourhood of z_l polar coordinates (r, φ) with $z - z_l = r \exp(2\pi\sqrt{-1}\varphi)$ and make a partial Fourier transformation with respect to φ : any function f may be decomposed into an infinite sum

$$f(r, \varphi) = \sum_{n \in \mathbb{N}} \hat{f}(n, r) \exp(2n\pi\sqrt{-1}\varphi).$$

The integrals are regularized by the rule

$$\int f(z) \delta(z - z_l) d^2x = \hat{f}(0, 0)$$

and the corresponding compatible rules for the holomorphic and anti-holomorphic derivatives of the δ -function. With these rules we obtain the formulas

$$\int_{\mathbb{R}^2/\Lambda} \Phi(z, k) \check{\Psi}(z) d^2x = \int_{\mathbb{R}^2/\Lambda} \check{\Phi}(z) \Psi(z, k) d^2x = R(k).$$

Here $\mathbf{R}(k)$ denotes the $M \times M$ matrices, whose entries indexed by $m, n = 0, \dots, M-1$ are equal to

$$\mathbf{R}(k)_{m,n} = \binom{m+n}{m} \frac{(-1)^n}{\pi} \begin{pmatrix} 0 & \mathcal{K}_{1,m+n}(k) \\ -\mathcal{K}_{2,m+n}(k) & 0 \end{pmatrix}.$$

Here $\mathcal{K}_{1,n}$ and $\mathcal{K}_{2,n}$ denote the Laurent series at $z = 0$ of the two functions $\mathcal{K}_1$ and $\mathcal{K}_2$ in the integral kernels of $\mathbf{R}(0, 0, k, 0)$ (compare with Lemma 3.20):

$$\mathcal{K}_1(z, k) = z^{-1} + \sum_{n=0}^{\infty} \mathcal{K}_{1,n}(k) z^n \quad \mathcal{K}_2(z, k) = \bar{z}^{-1} + \sum_{n=0}^{\infty} \mathcal{K}_{2,n}(k) \bar{z}^n.$$

Hence, due to Lemma 3.21, the inverse of the operator $\mathbf{1} - \mathbf{PR}(0, 0, k, 0)$ is equal to $\mathbf{1}$ plus the operator with integral kernel

$$\check{\Psi}(z) \mathbf{S}(k) \Phi(z', k) d^2 x',$$

where $\mathbf{S}(k)$ is the composition of $\mathbf{P}$ with the inverse of the restriction of $\mathbf{1} - \mathbf{PR}(k)$ to the range of $\mathbf{P}$. If $\mathbf{E}$ and $\mathbf{F}$ denote the kernel and the range of $\mathbf{P}$, respectively, then let $\tilde{\mathbf{P}}$ and $\tilde{\mathbf{R}}(k)$ denote the operators

$$\tilde{\mathbf{P}} : \mathbb{C}^2 \otimes \mathbb{C}^M / \mathbf{E} \xrightarrow{\mathbf{P}} \mathbf{F}, \quad \tilde{\mathbf{R}}(k) : \mathbf{F} \hookrightarrow \mathbb{C}^2 \otimes \mathbb{C}^M \xrightarrow{\mathbf{R}(k)} \mathbb{C}^2 \otimes \mathbb{C}^M \twoheadrightarrow \mathbb{C}^2 \otimes \mathbb{C}^M / \mathbf{E}.$$

Thus $\mathbf{S}(k)$ is given by

$$\mathbf{S}(k) : \mathbb{C}^2 \otimes \mathbb{C}^M \twoheadrightarrow \mathbb{C}^2 \otimes \mathbb{C}^M / \mathbf{E} \xrightarrow{(\tilde{\mathbf{P}}^{-1} - \tilde{\mathbf{R}}(k))^{-1}} \mathbf{F} \hookrightarrow \mathbb{C}^2 \otimes \mathbb{C}^M.$$

We remark that by definition of $\mathbf{E}$ and $\mathbf{F}$ the operator $\tilde{\mathbf{P}}$ is invertible. It is quite easy to see that all these functions $\mathbf{S}(k)$ obey for all $k, k' \in \mathbb{C}^2$ the relation

$$\mathbf{S}(k) - \mathbf{S}(k') = \mathbf{S}(k) (\mathbf{R}(k) - \mathbf{R}(k')) \mathbf{S}(k').$$

Conversely, any solution of these equations is of the form

$$\mathbf{S}(k) : \mathbb{C}^2 \otimes \mathbb{C}^M \twoheadrightarrow \mathbb{C}^2 \otimes \mathbb{C}^M / \mathbf{E} \xrightarrow{(\tilde{\mathbf{Q}} - \tilde{\mathbf{R}}(k))^{-1}} \mathbf{F} \hookrightarrow \mathbb{C}^2 \otimes \mathbb{C}^M,$$

where $\mathbf{F}$ and $\mathbf{E}$ are subspaces of $\mathbb{C}^2 \otimes \mathbb{C}^M$ such that the co-dimension of $\mathbf{E}$ is equal to the dimension of $\mathbf{F}$. Moreover, $\tilde{\mathbf{Q}}$ is any (not necessarily invertible) operator from $\mathbf{F}$ into $\mathbb{C}^2 \otimes \mathbb{C}^M / \mathbf{E}$.

2. If the support of the *Finite rank Perturbation* contains several points, then the index set (analogous to $\{0, \dots, M-1\}$) is determined by an integral divisor $D = \sum_{l=1}^L M_l z_l$. The analogous ansatz for the integral kernel of a *Finite rank Perturbation* is

$$\Psi_D(z, k) \mathbf{S}_D(k) \Phi_D(z', k) d^2 x',$$

where Ψ_D and Φ_D are the analogous row-vectors and column-vectors with $\deg(D)$ entries indexed by $m_l \in \bigcup_{l=1}^L \{0, \dots, M_l - 1\}$ and $\mathbf{S}_D$ is a $\deg(D) \times \deg(D)$ -matrix, all whose entries

are 2×2 -matrices. Let $\mathbf{R}_D(k)$ denote the $\deg(D) \times \deg(D)$ -matrix, whose entries indexed by $(m_i, n_j) \in \bigcup_{i=1}^L \{0, \dots, M_i - 1\} \times \bigcup_{j=1}^L \{0, \dots, M_j - 1\}$ are given by

$$\mathbf{R}_{m_i, n_j} = \begin{cases} \begin{pmatrix} m_i + n_i \\ m_i \end{pmatrix} \frac{(-1)^{n_i}}{\pi} \begin{pmatrix} 0 & \mathcal{K}_{1, m_i + n_i}(k) \\ -\mathcal{K}_{2, m_i + n_i}(k) & 0 \end{pmatrix} & \text{if } i = j \\ \frac{(-1)^{n_j}}{\pi m_i! n_j!} \begin{pmatrix} 0 & \mathcal{K}_1^{(m_i + n_j)}(z_i - z_j, k) \\ -\mathcal{K}_2^{(m_i + m_j)}(z_i - z_j, k) & 0 \end{pmatrix} & \text{if } i \neq j. \end{cases}$$

Therefore, the *Finite rank Perturbations* of $\mathbf{R}(0, 0, k, 0)$ are defined as the operators with integral kernels

$$\Psi_D(z, k) \mathbf{S}_D(k) \Phi_D(z', k) d^2 x'.$$

Here D is any integral divisor on $\mathbb{C}/\Lambda_{\mathbb{C}}$ and $\mathbf{S}_D(k)$ satisfies for all $k, k' \in \mathbb{C}^2$ the relation

$$\mathbf{S}_D(k) - \mathbf{S}_D(k') = \mathbf{S}_D(k) (\mathbf{R}_D(k) - \mathbf{R}_D(k')) \mathbf{S}_D(k').$$

3. The perturbed free resolvent $\mathbf{R}(0, 0, k, 0) + \mathbf{R}(0, 0, k, 0) \mathbf{S}(k) \mathbf{R}(0, 0, k, 0)$ induces the following perturbation of the resolvent $\mathbf{R}(V, W, k, 0)$:

$$\begin{aligned} & \mathbf{R}(0, 0, k, 0) (\mathbf{1} + \mathbf{S}(k) \mathbf{R}(0, 0, k, 0)) (\mathbf{1} - \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \mathbf{R}(0, 0, k, 0) (\mathbf{1} + \mathbf{S}(k) \mathbf{R}(0, 0, k, 0)))^{-1} \\ &= \mathbf{R}(0, 0, k, 0) ((\mathbf{1} + \mathbf{S}(k) \mathbf{R}(0, 0, k, 0))^{-1} - \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \mathbf{R}(0, 0, k, 0))^{-1} \\ &= \mathbf{R}(0, 0, k, 0) (\mathbf{1} - (\mathbf{1} + \mathbf{S}(k) \mathbf{R}(0, 0, k, 0))^{-1} \mathbf{S}(k) \mathbf{R}(0, 0, k, 0) - \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \mathbf{R}(0, 0, k, 0))^{-1} \\ &= \mathbf{R}(V, W, k, 0) (\mathbf{1} - (\mathbf{1} + \mathbf{S}(k) \mathbf{R}(0, 0, k, 0))^{-1} \mathbf{S}(k) \mathbf{R}(V, W, k, 0))^{-1}. \end{aligned}$$

This implies that the corresponding perturbation is equal to

$$\begin{aligned} & \mathbf{R}(V, W, k, 0) \mathbf{S}(V, W, k) \mathbf{R}(V, W, k, 0) \text{ with } \mathbf{S}(V, W, k) = \\ &= (\mathbf{1} - (\mathbf{1} + \mathbf{S}(k) \mathbf{R}(0, 0, k, 0))^{-1} \mathbf{S}(k) \mathbf{R}(V, W, k, 0))^{-1} (\mathbf{1} + \mathbf{S}(k) \mathbf{R}(0, 0, k, 0))^{-1} \mathbf{S}(k) \\ &= (\mathbf{1} + \mathbf{S}(k) \mathbf{R}(0, 0, k, 0) - \mathbf{S}(k) \mathbf{R}(V, W, k, 0))^{-1} \mathbf{S}(k) \\ &= \mathbf{S}(k) (\mathbf{1} + \mathbf{R}(0, 0, k, 0) \mathbf{S}(k) - \mathbf{R}(V, W, k, 0) \mathbf{S}(k))^{-1}. \end{aligned}$$

In the notation of the first two steps $\mathbf{S}(k)$ has the integral kernel $\check{\Psi}_D(z) \mathbf{S}_D(k) \check{\Phi}_D(z') d^2 x'$. With the regularization rules of the first step we have $\mathbf{R}_D(k) = \left\langle \left\langle \check{\Phi}_D, \mathbf{R}_D(0, 0, k, 0) \check{\Psi}_D \right\rangle \right\rangle$. For pairs of smooth potentials we may analogously define $\mathbf{R}_D(V, W, k) = \left\langle \left\langle \check{\Phi}_D, \mathbf{R}_D(V, W, k, 0) \check{\Psi}_D \right\rangle \right\rangle$. Then the perturbation $\mathbf{S}(V, W, k)$ has the integral kernel $\check{\Psi}_D(z) \mathbf{S}_D(V, W, k) \check{\Phi}_D(z') d^2 x'$ with

$$\begin{aligned} \mathbf{S}_D(V, W, k) &= (\mathbf{1} + \mathbf{S}_D(k) (\mathbf{R}(k_D) - \mathbf{R}_D(V, W, k)))^{-1} \mathbf{S}_D(k) \\ &= \mathbf{S}_D(k) (\mathbf{1} + (\mathbf{R}(k_D) - \mathbf{R}_D(V, W, k)) \mathbf{S}_D(k))^{-1}. \end{aligned}$$

Moreover, due to the relation $S_D(k) - S_D(k') = S_D(k) (R_D(k) - R_D(k')) S_D(k')$, the matrices $S_D(V, W, k)$ obey the relation

$$S_D(V, W, k) - S_D(V, W, k') = S_D(k) (R_D(V, W, k) - R_D(V, W, k')) S_D(V, W, k').$$

For pairs of non-smooth potentials the matrices $R_D(V, W, k)$ are not defined.

4. However, a reformulation of these perturbations makes sense for all pairs of potentials in L^2 . In fact, the perturbation $R(V, W, k, 0)S(V, W, k)R(V, W, k, 0)$ has the integral kernel

$$\Psi_D(z, k)S(V, W, k)\Phi_D(k, z')d^2x'$$

in terms of the row-vectors and column-vectors

$$\Psi_D(z, k) = R(V, W, k, 0)\check{\Psi}_D(z) \text{ and } \Phi_D^t(z, k) = R^t(V, W, k, 0)\check{\Phi}_D^t(z).$$

Furthermore, due to Lemma 2.22, the forms $\Phi_D(z, k) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \Psi_D(z, k')$ are closed on $\Delta \setminus \{z_1, \dots, z_L\}$, and for all small ε the integrals over the boundary of $S_\varepsilon = \bigcup_{l=1}^L B(z_l, \varepsilon)$ are equal to

$$\int_{\partial S_\varepsilon} \Phi_D(z, k) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \Psi_D(z, k') = 2\sqrt{-1} (R_D(V, W, k) - R_D(V, W, k')).$$

Hence we may renormalize $\Psi_D(z, k)$ and $\Phi_D(z, k)$ in such a way, that they have finite L^q -norm on the complement of S_ε . This leads to the following definition:

Finite rank Perturbations 3.22. *of $R(V, W, k, 0)$ are defined as perturbations by the operators with integral kernels*

$$\Psi_D(z, k)S_D(k)\Phi_D(z', k)d^2x'.$$

Here $D = \sum_{l=1}^L M_l z_l$ is any integral divisor on $\mathbb{C}/\Lambda_{\mathbb{C}}$, Ψ_D and Φ_D are finite row-vectors and column-vectors, whose entries belong on $\Delta \setminus \{z_1, \dots, z_L\}$ to the kernel and co-kernel of $D(V, W, k)$, respectively. Moreover, for all $l = 1, \dots, L$ these vectors obey on a small neighbourhood of z_l the equations

$$\begin{pmatrix} V(\bar{z} - \bar{z}_l)^{M_l} & \partial(\bar{z} - \bar{z}_l)^{M_l} \\ -\bar{\partial}(z - z_l)^{M_l} & W(z - z_l)^{M_l} \end{pmatrix} \Psi_D(z, k) = 0 \quad \begin{pmatrix} V(z - z_l)^{M_l} & \bar{\partial}(z - z_l)^{M_l} \\ -\partial(\bar{z} - \bar{z}_l)^{M_l} & W(\bar{z} - \bar{z}_l)^{M_l} \end{pmatrix} \Phi_D^t(z, k) = 0.$$

Furthermore, the representatives of these entries in the quotients of these kernels modulo the kernels and co-kernels of $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ (compare with Lemma 3.43), do not depend on k (i. e. locally there exist $\tilde{\Psi}_D(z, k)$ and $\tilde{\Phi}_D(z, k)$ in the kernels and co-kernel of $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$, such that the differences $\Psi_D(z, k) - \tilde{\Psi}_D(z, k)$ and $\Phi_D(z, k) - \tilde{\Phi}_D(z, k)$ do not depend on k). Finally, for all k, k' the matrices S_D obey the equation

$$S_D(k) - S_D(k') = \frac{1}{2\sqrt{-1}} \int_{\partial S_\varepsilon} S_D(k) \Phi_D(z, k) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \Psi_D(z, k') S_D(k').$$

Remark 3.23. The kernels of the operators $\begin{pmatrix} V(\bar{z}-\bar{z}_l)^{M_l} & \partial(\bar{z}-\bar{z}_l)^{M_l} \\ -\bar{\partial}(z-z_l)^{M_l} & W(z-z_l)^{M_l} \end{pmatrix}$ and $\begin{pmatrix} V(z-z_l)^{M_l} & \bar{\partial}(z-z_l)^{M_l} \\ -\partial(\bar{z}-\bar{z}_l)^{M_l} & W(\bar{z}-\bar{z}_l)^{M_l} \end{pmatrix}$ are those spinors ψ and ϕ , such that $\begin{pmatrix} z-z_l & 0 \\ 0 & \bar{z}-\bar{z}_l \end{pmatrix}^{M_l} \psi$ and $\begin{pmatrix} \bar{z}-\bar{z}_l & 0 \\ 0 & z-z_l \end{pmatrix}^{M_l} \phi$ belongs to the kernel of $\begin{pmatrix} V\left(\frac{\bar{z}-\bar{z}_l}{z-z_l}\right)^{M_l} & \partial \\ -\bar{\partial} & W\left(\frac{z-z_l}{\bar{z}-\bar{z}_l}\right)^{M_l} \end{pmatrix}$ and $\begin{pmatrix} V\left(\frac{z-z_l}{\bar{z}-\bar{z}_l}\right)^{M_l} & \bar{\partial} \\ -\partial & W\left(\frac{\bar{z}-\bar{z}_l}{z-z_l}\right)^{M_l} \end{pmatrix}$, respectively (compare with Section 3.4.4).

5. We shall generalize Cauchy's integral formula to these elements in the kernel of Dirac operators with potentials $V, W \in L^2$. If $\mathcal{K}_{\mathbb{R}^2}(V, W, z, z')$ denotes the integral kernel of the resolvent $R_{\mathbb{R}^2}(V, W, 0)$ on the bounded open set $\mathcal{O} \subset \mathbb{C}$, then we have

$$\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix} \mathcal{K}_{\mathbb{R}^2}(V, W, z, z') = \pi \delta(z - z') \mathbf{1} \quad \begin{pmatrix} V & \bar{\partial} \\ -\partial & W \end{pmatrix} \mathcal{K}_{\mathbb{R}^2}^t(V, W, z', z) = \pi \delta(z - z') \mathbf{1}.$$

Here the Dirac operator and his transposed acts on the integral kernel as a function depending on z for fixed z' . The arguments of the proof of Lemma 2.22 imply the

Generalized Cauchy's integral formula 3.24. All elements ψ and ϕ in the kernel and co-kernel of the Dirac operator $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ on a small open set $\mathcal{O}$ obey the formula

$$\begin{aligned} \psi(z') &= \frac{1}{2\pi\sqrt{-1}} \oint \mathcal{K}_{\mathbb{R}^2}(V, W, z', z) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \psi(z) \\ \phi(z') &= \frac{-1}{2\pi\sqrt{-1}} \oint \phi(z) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \mathcal{K}_{\mathbb{R}^2}(V, W, z, z'), \end{aligned}$$

as long as the integration path surrounds z' one times in the anti-clockwise-order, respectively.

Remark 3.25. At a first look it is not clear, whether the integral along the closed path is well defined. However, since on the complement of $\{z'\}$ the corresponding one-forms are closed, we may extend the integration over the closed path to an integration over a cylinder around z' . This extended closed integral can be considered as the evaluation of the action of the resolvent on a L^q -spinor. Moreover, since the spinors ψ and ϕ are in general not continuous, this equality should be understood as an equality of measurable functions.

6. Now we may extend the classification of these *Finite rank Perturbations* 3.22 of the free resolvents in the first two steps to pairs of L^2 -potentials. For all integral divisors $D = \sum_{l=1}^L M_l z_l$ we introduce the sheaves analogous to $\mathcal{O}_D$. On the complement of the support of D the corresponding sections belong to the kernels and co-kernels of $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$. Moreover, for all $l = 1, \dots, L$ these sections ψ and ϕ obey in a small neighbourhood of z_l

$$\begin{pmatrix} V(\bar{z}-\bar{z}_l)^{M_l} & \partial(\bar{z}-\bar{z}_l)^{M_l} \\ -\bar{\partial}(z-z_l)^{M_l} & W(z-z_l)^{M_l} \end{pmatrix} \psi = 0 \quad \begin{pmatrix} V(z-z_l)^{M_l} & \bar{\partial}(z-z_l)^{M_l} \\ -\partial(\bar{z}-\bar{z}_l)^{M_l} & W(\bar{z}-\bar{z}_l)^{M_l} \end{pmatrix} \phi^t = 0.$$

We chose a basis of the quotient of these sheaves modulo the kernel and co-kernel of $\begin{pmatrix} V & \partial \\ -\partial & W \end{pmatrix}$ (compare with Lemma 3.43), respectively. These basis should be represented by local elements ψ_{m_i} and ϕ_{m_i} of these sheaves nearby the singular points $z_1, \dots, z_L$. We arrange the former basis to a row-vector $\Psi_D(z)$ and the latter basis to a column-vector $\Phi_D(z)$ with $\deg(D)$ entries, respectively. Moreover, we assume that the corresponding matrix of residues vanishes

$$\text{Res}_D \Phi_D(z) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \Psi_D(z) = 0.$$

If $\mathcal{K}(V, W, k, z, z') \frac{d^2 x'}{\pi}$ denotes the integral kernel of the operator $R(V, W, k, 0)$, then the row-vectors and columns vectors

$$\begin{aligned} \Psi_D(z', k) &= \text{Res}_D \mathcal{K}(V, W, k, z', z) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \Psi_D(z) \\ \Phi_D(z', k) &= -\text{Res}_D \Phi_D(z) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \mathcal{K}(V, W, k, z, z') \end{aligned}$$

fulfill the conditions of *Finite rank Perturbations* 3.22. We remark that these functions are defined only for z' outside of the cycles around the support of D , which are used to calculate the residue, and for $k \notin \mathcal{F}(V, W)$. Due to the *Generalized Cauchy's integral formula* 3.24 the differences $\Psi_D(z, k) - \Psi_D(z)$ and $\Phi_D(z, k) - \Phi_D(z)$ belong on a small neighbourhood of the support of D to the kernel and co-kernel of $\begin{pmatrix} V & \partial \\ -\partial & W \end{pmatrix}$, respectively. Since the product $\Phi_D(z, k) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \Psi_D(z, k)$ is periodic the matrix of residues $\text{Res}_D \Phi_D(z, k) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \Psi_D(z, k)$ vanishes. Hence the following matrices of residues coincide:

$$R_D(V, W, k) = -\pi \text{Res}_D \Phi_D(z) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \Psi_D(z, k) = \pi \text{Res}_D \Phi_D(z, k) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \Psi_D(z).$$

In general we have

$$\text{Res}_D \Phi_D(z, k) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \Psi_D(z, k') = R_D(V, W, k) - R_D(V, W, k'),$$

and the classification of *Finite rank Perturbations* 3.22 of $R(0, 0, k, 0)$ in the first two steps carries over to a classification of *Finite rank Perturbations* 3.22 of $R(V, W, k, 0)$. Consequently, all *Finite rank Perturbations* 3.22 of $R(V, W, k, 0)$ have an integral kernel of the form $\Psi_D(z, k) S_D(k) \Phi_D(z', k) d^2 x'$ with

$$S_D(k) - S_D(k') = S_D(k) (R_D(V, W, k) - R_D(V, W, k')) S_D(k').$$

7. The foregoing considerations lead to another description of these *Finite rank Perturbations* 3.22. The corresponding Dirac operators yield holomorphic structures (in the sense of ‘quaternionic function theory’ [P-P, B-F-L-P-P, F-L-P-P]) with singularities at the points $\{z_1, \dots, z_L\}$. Let us first explain these singularities for one singular point z_l . We have seen in the first step that in these cases the matrices S are of the form

$$S(k) : \mathbb{C}^2 \otimes \mathbb{C}^M \rightarrow \mathbb{C}^2 \otimes \mathbb{C}^M / E \xrightarrow{(\tilde{Q} - \tilde{R}(k))^{-1}} F \hookrightarrow \mathbb{C}^2 \otimes \mathbb{C}^M.$$

Here F and E are subspaces of $\mathbb{C}^2 \otimes \mathbb{C}^M$ such that the co-dimension of E is equal to the dimension of F . Moreover, $\tilde{Q}$ is any operator from F into $\mathbb{C}^2 \otimes \mathbb{C}^M/E$. For smooth potentials we determined formally in the first step the corresponding perturbations of the Dirac operators. Surprisingly, we may define the kernels of these perturbed Dirac operators for all L^2 -potentials. In fact, by choice of the row-vectors $\Psi(z)$ and column-vectors $\Phi(z)$ in the sixth step these data $(E, F, \tilde{Q})$ are in one-to-one correspondence to the deformed kernels H_{sing, z_l} with the following properties:

Singularity (i) The space H_{sing, z_l} is a subspace of the kernel of $\begin{pmatrix} V(\bar{z}-\bar{z}_l)^{M_l} & \partial(\bar{z}-\bar{z}_l)^{M_l} \\ -\bar{\partial}(z-z_l)^{M_l} & W(z-z_l)^{M_l} \end{pmatrix}$ of co-dimension $2M_l$, i. e. the same co-dimension as the kernel of $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ (compare with Lemma 3.43).

Singularity (ii) The kernel of $\begin{pmatrix} V(\bar{z}-\bar{z}_l)^{-M_l} & \partial(\bar{z}-\bar{z}_l)^{-M_l} \\ -\bar{\partial}(z-z_l)^{-M_l} & W(z-z_l)^{-M_l} \end{pmatrix}$ is a subspace of the space H_{sing, z_l} of co-dimension $2M_l$, i. e. the same co-dimension as this kernel considered as a subspace of the kernel of $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ (compare with Lemma 3.43).

More precisely, the entries of $\Psi(z)$ fit together to a basis of the quotient of the kernel of $\begin{pmatrix} V(\bar{z}-\bar{z}_l)^{M_l} & \partial(\bar{z}-\bar{z}_l)^{M_l} \\ -\bar{\partial}(z-z_l)^{M_l} & W(z-z_l)^{M_l} \end{pmatrix}$ modulo the kernel of $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$. On the other hand the entries of $\Phi(z)$ are dual with respect to the pairing given by the residue to a basis of the quotient of the kernel of $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ modulo the kernel of $\begin{pmatrix} V(\bar{z}-\bar{z}_l)^{-M_l} & \partial(\bar{z}-\bar{z}_l)^{-M_l} \\ -\bar{\partial}(z-z_l)^{-M_l} & W(z-z_l)^{-M_l} \end{pmatrix}$. Due to the assumption that the matrix of residues of $\Phi(z) \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \Psi(z)$ vanishes, these two basis fit together to a basis of the quotient of the kernel of $\begin{pmatrix} V(\bar{z}-\bar{z}_l)^{M_l} & \partial(\bar{z}-\bar{z}_l)^{M_l} \\ -\bar{\partial}(z-z_l)^{M_l} & W(z-z_l)^{M_l} \end{pmatrix}$ modulo the kernel of $\begin{pmatrix} V(\bar{z}-\bar{z}_l)^{-M_l} & \partial(\bar{z}-\bar{z}_l)^{-M_l} \\ -\bar{\partial}(z-z_l)^{-M_l} & W(z-z_l)^{-M_l} \end{pmatrix}$. Therefore the spaces E and F are subspaces of this quotient, and the corresponding space H_{sing, z_l} is the sum of the the kernel of $\begin{pmatrix} V(\bar{z}-\bar{z}_l)^{-M_l} & \partial(\bar{z}-\bar{z}_l)^{-M_l} \\ -\bar{\partial}(z-z_l)^{-M_l} & W(z-z_l)^{-M_l} \end{pmatrix}$ plus E plus the graph of $\tilde{Q}$ considered as an operator from F into the quotient of the kernel of $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ modulo the sum of the kernel of $\begin{pmatrix} V(\bar{z}-\bar{z}_l)^{-M_l} & \partial(\bar{z}-\bar{z}_l)^{-M_l} \\ -\bar{\partial}(z-z_l)^{-M_l} & W(z-z_l)^{-M_l} \end{pmatrix}$ plus E .

In analogy to ‘quaternionic functions theory’ the Dirac operators are considered as holomorphic structures. Consequently, the *Finite rank Perturbations* 3.22 give rise to

Singularities of the holomorphic structure 3.26. *The kernel of a perturbed Dirac operators is locally in a neighbourhood $\mathcal{O}$ of a singular point z_1 a subspace H_{sing, z_l} of*

$$\left\{ \psi \mid \begin{pmatrix} z-z_l & 0 \\ 0 & \bar{z}-\bar{z}_l \end{pmatrix}^{M_l} \psi \in W_{\text{loc}}^{1,p}(\mathcal{O}) \times W_{\text{loc}}^{1,p}(\mathcal{O}) \right\} \text{ with } 1 < p < 2 \quad (\text{compare with Section 3.4.4}),$$

which fulfills conditions Singularity (i)–(ii).

Obviously, for a given pair of potentials and a given M_l the set of those spaces H_{sing, z_l} , which obey condition *Singularity* (i)–(ii), can be considered as a finite-dimensional Grassmannian [Gr-Ha, Chapter 1 §5.]. The resolvents of the *Finite rank Perturbations* 3.22 maps all spinors, whose support is disjoint from a small neighbourhood $\mathcal{O}$ of z_l , into the corresponding deformed kernel H_{sing, z_l} . Conversely, the corresponding *Finite rank Perturbations* 3.22 are uniquely defined by this property.

Remark 3.27. *From the very beginning it is not clear, whether for all these choices of spaces H_{sing, z_l} (or equivalently of data $(E, F, \tilde{Q})$) the corresponding matrices $S(k)$ are indeed meromorphic. More precisely, the finiteness of $S(k)$ for at least one value has to be guaranteed. We show this in the following step for all perturbations, which respect the involution η . The corresponding spaces H_{sing, z_l} are invariant under the (projective) involution $\psi \mapsto J\bar{\psi}$.*

Remark 3.28. *The results of Section 2.5.3 imply that these singularities contribute to the Willmore functional. More precisely, if $\mathcal{F}$ is the complex Fermi curve of a Finite rank Perturbation 3.22 of $R(U, \bar{U}, k, 0)$, then the Willmore functional is equal to*

$$\mathcal{W}(\mathcal{F}) = 4\|U\|_2^2 + \sum_{l=1}^L \mathcal{W}_{\text{sing}, z_l}.$$

In the correspondence of spaces H_{sing, z_l} and data $(E, F, \tilde{Q})$ we realized F as a space of ‘meromorphic’ spinors and E as a space of ‘holomorphic’ spinors. Let $0 < n_1 + 1 < \dots < n_J + 1$ denote the sequence of pole orders of the elements of F and $0 \leq m_1 < \dots < m_J$ denote the sequence of Orders of zeroes 3.41 of the elements of E . (In Section 3.4.3 and 3.4.5 we will see that these sequences of integers $0 \leq n_1 < \dots < n_J$ and $0 \leq m_1 < \dots < m_J$ are the Orders of zeroes 3.41 at ∞ of the spinors in the kernel and co-kernel of a blown up Dirac operator on $\mathbb{P}^1$.) The corresponding sets $\mathcal{N}$ introduced in Section 2.5.3 are given by

$$\mathcal{N} = \{-n_J - 1, \dots, -n_1 - 1\} \cup \mathbb{N}_0 \setminus \{m_1, \dots, m_J\}.$$

The corresponding local contribution to the Willmore functional is equal to

$$\mathcal{W}_{\text{sing}, z_l} = 4\pi (J + n_1 + \dots + n_J + m_1 + \dots + m_J).$$

For all $d \in \mathbb{N}$ let $\mathcal{L}_{(d-1)\infty}$ denote the sheaf of spinors in the kernel of the corresponding Dirac operator on $\mathbb{P}^1$, which may have poles of order $d - 1$ at ∞ , i. e. the sections obey $\text{ord}_\infty(\psi) \geq 1 - d$ (compare with Section 3.4.4). The corresponding quaternionic holomorphic line bundle has degree d . Due to extension of S erre Duality and the Riemann–Roch theorem to these quaternionic holomorphic line bundles (compare with [F-L-P-P, Theorem 2.2] and Section 3.4.4) for large d the quaternionic dimension of $\dim H^0(\mathbb{P}^1, \mathcal{L}_{(d-1)\infty})$ is equal to $d + 1$. The corresponding Orders of zeroes 3.41 at ∞ are equal to the numbers $\{n \in \mathbb{N}_0 \mid d - n - 1 \in \mathcal{N}\}$. The corresponding numbers, which in [F-L-P-P, Definition 4.2] are called $\text{ord}_\infty(\mathcal{L}_{(d-1)\infty})$, are for large d equal to

$$\text{ord}_\infty(\mathcal{L}_{(d-1)\infty}) = \sum_{\{n \in \mathbb{N}_0 \mid d - n - 1 \in \mathcal{N}\}} n - \frac{1}{2}d(d - 1) = \sum_{\{n \in \mathbb{N} \mid -n \in \mathcal{N}\}} n + \sum_{\{n \in \mathbb{N}_0 \mid n \notin \mathcal{N}\}} n = \frac{\mathcal{W}_{\text{sing}, z_l}}{4\pi}.$$

Hence the local contribution to the Willmore functional is exactly equal to the lower bound of the Pl ucker formula [F-L-P-P, Theorem 4.7] for the quaternionic holomorphic line bundle corresponding to $\mathcal{L}_{(d-1)\infty}$ for large d .

Remark 3.29. *The local structure of these perturbations yields a further condition for the Finite rank Perturbations 3.22. The data $(E_D, F_D, \tilde{Q}_D)$ have diagonal block form with respect to the decomposition into L blocks corresponding to the singular points $\{z_1, \dots, z_l\}$. Moreover, if the pair of potentials is invariant under one or several involutions introduced in Section 2.3, then the corresponding perturbations are covariant under the corresponding involutions (compare with Remark 3.60).*

Remark 3.30. *The local structure suggests the following generalization of Theorem 3.48: A subsequence of the sequence of kernels H_n of any sequence of Dirac operators $\begin{pmatrix} U_n & \partial \\ -\bar{\partial} & \bar{U}_n \end{pmatrix}$ with bounded potentials $U_n \in L^2(\mathcal{O})$ on a bounded open domain $\mathcal{O} \subset \mathbb{C}$ converges (in an appropriate sense) to a kernel with Singularities of the holomorphic structure 3.26. From this point of view the Quaternionic Singularity condition 1.6 states that there should exist a quaternionic holomorphic line bundle whose index of specialty is at least equal to one. Therefore, the extension to immersion of higher genus Riemann surfaces depends on an investigation of the subsets of the Picard groups of special divisors. For genus one, these are exactly the complex Fermi curves. For higher genus, the corresponding Bloch theory give rise to higher-dimensional analogs of complex Fermi curves.*

The spinors in the kernel of a given Dirac operator $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ are uniquely determined by the restrictions to the boundary $\partial B(z_l, \varepsilon)$ of a ball $B(z_l, \varepsilon)$. Moreover, the spaces H_{sing, z_l} may be described by the corresponding subspaces of the Banach spaces $L^q(\partial B(z_l, \varepsilon) \times L^q(\partial B(z_l, \varepsilon))$. The description of the latter subspaces as graphs of operators yields a suitable notion of convergence (compare with the description of the Grassmannian of a Hilbert space in [Pr-Se, Chapter 7.]). Obviously, this generalization extends the compactness property to immersions of arbitrary compact Riemann surfaces into $\mathbb{R}^3$ and $\mathbb{R}^4$.

8. In this step we shall extend the asymptotic analysis of Theorem 2.12 to those *Finite rank Perturbations* 3.22, which respect the involution η . Due to the proof of Lemma 2.11 the resolvents $R(U, \bar{U}, k, 0)$ obey the relations

$$R(U, \bar{U}, k + k_\kappa^+, 0) = R(U, \bar{U}, k + k_\kappa^-, 0) = \begin{pmatrix} \psi_{-k_\kappa^+} & 0 \\ 0 & \psi_{-k_\kappa^-} \end{pmatrix} R(\psi_{-\kappa} U, \psi_\kappa \bar{U}, k, 0) \begin{pmatrix} \psi_{k_\kappa^-} & 0 \\ 0 & \psi_{k_\kappa^+} \end{pmatrix}.$$

This implies that these transformations acts on the subspaces H_{sing, z_l} as the transformation

$$H_{\text{sing}, z_l} \mapsto \begin{pmatrix} \psi_{-k_\kappa^+} & 0 \\ 0 & \psi_{-k_\kappa^-} \end{pmatrix} H_{\text{sing}, z_l} = \left\{ \begin{pmatrix} \psi_{-k_\kappa^+} & 0 \\ 0 & \psi_{-k_\kappa^-} \end{pmatrix} \psi \mid \psi \in H_{\text{sing}, z_l} \right\}.$$

If H_{sing, z_l} obeys conditions *Singularity* (i)–(ii) and is invariant under the (projective) involution $\psi \mapsto J\bar{\psi}$, then in the limit $\|\kappa\| \rightarrow \infty$ the transformed spaces $\begin{pmatrix} \psi_{-k_\kappa^+} & 0 \\ 0 & \psi_{-k_\kappa^-} \end{pmatrix} H_{\text{sing}, z_l}$ converge to the kernel of the free Dirac operator. In fact, due to Theorem 2.12 the limit corresponds to a *Singularity of the holomorphic structure* 3.26 corresponding to the free Dirac operator. Moreover the limits are fixed points under the former transformations. Therefore they are invariant under $\begin{pmatrix} z - z_l & 0 \\ 0 & \mathbf{1} \end{pmatrix}$ and $\begin{pmatrix} \mathbf{1} & 0 \\ 0 & \bar{z} - \bar{z}_l \end{pmatrix}$. Obviously, the kernel of the free Dirac operator is the only subspace H_{sing, z_l} , which fulfills conditions *Singularity* (i)–(ii) (for the free Dirac

operator), and which is invariant with respect to the action of these two operators and with respect to the (projective) involution $\psi \mapsto J\bar{\psi}$. Now Theorem 2.12 carries over to the *complex Fermi curves* of *Finite rank Perturbations* 3.22 of $R(U, \bar{U}, k, 0)$ (with $U \in L^2(\mathbb{R}^2/\Lambda)$) corresponding to spaces H_{sing, z_l} , which are invariant with respect to the (projective) involution $\psi \mapsto J\bar{\psi}$. More precisely, in the estimates of the corresponding eigenfunctions the norms $\|\cdot\|_q$ has to be replaced by the norms of the corresponding restrictions to the relative complement of $S_\varepsilon = \bigcup_{l=1}^L B(z_l, \varepsilon)$. In particular, the matrices $S_D(k)$ corresponding to all these *Finite rank Perturbations* 3.22 are indeed meromorphic functions on $\mathbb{C}^2$ (compare with Remark 3.27), whose polar sets are the corresponding *complex Fermi curves*.

9. The methods of the fourth step yield another explanation of the relation in the definition *Finite rank Perturbations* 3.22. Let ψ and ϕ be spinors with support in S_ε . The proof of Lemma 2.22 implies the relation

$$\begin{aligned} \langle \langle R^t(V, W, k, 0)\phi, \psi \rangle \rangle - \langle \langle \phi, R(V, W, k', 0)\psi \rangle \rangle = \\ \frac{1}{2\sqrt{-1}} \int_{\partial S_\varepsilon} \left\langle \left\langle R^t(V, W, k, 0)\phi, \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} R(V, W, k', 0)\psi \right\rangle \right\rangle. \end{aligned}$$

The analog of this relation for the perturbed resolvents imply the relation in the definition *Finite rank Perturbations* 3.22. In fact, with the help of the row-vectors $\Psi_D(z)$ and the column-vectors $\Phi_D(z)$ introduced in the sixth step, the entries of Ψ_D and Φ_D can be distinguished by suitable ψ and ϕ . Therefore the analog to the former relation implies the latter.

10. In general the limits of the resolvents $R(U_n, \bar{U}_n, k, \lambda)$ depend non-continuously on λ , and do not obey the *first resolvent formula* [R-S-I, Theorem VI.5]. But as we already saw in Section 2.4, some modifications of the Dirac operators still reflect properties of *complex Fermi curves*. In fact, since the whole spectrum of the family of operators $\begin{pmatrix} 0 & \pm \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix} D(V, W, k)$ is determined by the *complex Fermi curves*, the limits of the corresponding resolvents might satisfy the corresponding *first resolvent formula*. More precisely, due to the relation

$$\begin{aligned} \begin{pmatrix} 0 & k'_2 - k_2 + \sqrt{-1}(k'_1 - k_1) \\ k'_2 - k_2 - \sqrt{-1}(k'_1 - k_1) & 0 \end{pmatrix} \tilde{D}(V, W, k') = \\ \pi g(k' - k, k' - k) \mathbf{1} + \begin{pmatrix} 0 & k'_2 - k_2 + \sqrt{-1}(k'_1 - k_1) \\ k'_2 - k_2 - \sqrt{-1}(k'_1 - k_1) & 0 \end{pmatrix} \tilde{D}(V, W, k) \end{aligned}$$

the corresponding resolvents obey the [R-S-I, Theorem VI.5]

Resolvent Formula 3.31.

$$\begin{aligned} \tilde{R}(V, W, k, 0) - \tilde{R}(V, W, k', 0) = \\ \pi \tilde{R}(V, W, k, 0) \begin{pmatrix} 0 & k_2 - k'_2 + \sqrt{-1}(k_1 - k'_1) \\ k_2 - k'_2 - \sqrt{-1}(k_1 - k'_1) & 0 \end{pmatrix} \tilde{R}(V, W, k', 0). \end{aligned}$$

At a first look condition *Finite rank Perturbations* (iii) and the corresponding relation in the definition *Finite rank Perturbations* 3.22 seems to be equivalent to the analog of the *Resolvent Formula* 3.31 for these *Finite rank Perturbations* 3.22. However, they do not obey the *Resolvent Formula* 3.31. More precisely, if the support of $\tilde{\psi}$ and $\tilde{\phi}$ are disjoint with small balls $B(z_1, \varepsilon), \dots, B(z_L, \varepsilon)$ around the support of the *Finite rank Perturbations*, then the action of $\tilde{D}(V, W, k)$ and $\tilde{D}^t(V, W, k)$ on $\tilde{R}(V, W, k, 0)\tilde{\psi}$ and $\tilde{R}^t(V, W, k, 0)\tilde{\phi}$ yields functions, which vanish on these balls, respectively. We decompose the integrals appearing in the matrix element

$$\begin{aligned} & \left\langle \left\langle \tilde{\phi}, \tilde{R}^t(V, W, k', 0) \begin{pmatrix} 0 & k'_2 - k_2 + \sqrt{-1}(k'_1 - k_1) \\ k'_2 - k_2 - \sqrt{-1}(k'_1 - k_1) & 0 \end{pmatrix} \tilde{R}(V, W, k, 0)\tilde{\psi} \right\rangle \right\rangle \\ & = 2\pi\sqrt{-1} \left\langle \left\langle \tilde{R}^t(V, W, k', 0)\tilde{\phi}, \tilde{R}(V, W, k, 0)\tilde{\psi} \right\rangle \right\rangle_{(k'_2 - k_2, k_1 - k'_1)} \end{aligned}$$

of the right hand side of the *Resolvent Formula* 3.31 into an integral over the complements of the small balls $B(z_1, \varepsilon) \cup \dots \cup B(z_L, \varepsilon)$ and an integral over these balls. Then, due to Lemma 2.25, the latter integral is equal to an integral of a one-form over the boundaries of these balls. An easy calculation shows that for all such $\tilde{\psi}$ and $\tilde{\phi}$ and all *Finite rank Perturbations* 3.22 the corresponding matrix elements of the left hand side of the *Resolvent Formula* 3.31 coincides with the corresponding matrix elements of the right hand side, if we replace the integral over the balls $B(z_1, \varepsilon), \dots, B(z_L, \varepsilon)$ by the integral of the corresponding one-form over the boundaries of these balls (described in Lemma 2.25). Furthermore, this condition does not result in any restriction on the functions $S(k)$. However, in the limit $\varepsilon \downarrow 0$ these boundary terms do not tend to zero for all k .

We shall see that the limits of resolvents are such *Finite rank Perturbations* 3.22. In a first step we investigate the limits $t \rightarrow \infty$ of the resolvents of a family $(U_t)_{t \in [1, \infty)}$ of potentials of the form $U_t(z) = tU_{t=1}(tz)$, where $U_{t=1}$ has support in $B(0, \varepsilon)$. Therefore, the potentials U_t have support in $B(0, t^{-1}\varepsilon)$ and the L^2 -norm does not depend on t . Moreover, for all $U_0 \in L^2(\mathbb{R}^2/\Lambda)$ the measures $(U_0 + U_t)(\bar{U}_0 + \bar{U}_t)d^2x$ converge in the limit $t \rightarrow \infty$ to the measure $U_0\bar{U}_0d^2x$ plus the point measure at $z = 0$ with mass equal to the square of the L^2 -norm of $U_{t=1}$. If the L^2 -norm of $U_{t=1}$ is smaller than the constant S_p^{-1} and k does not belong to the *complex Fermi curve* $\mathcal{F}(U_0, \bar{U}_0)$, then Lemma 2.6 implies that the family of resolvents $R(U_0 + U_t, \bar{U}_0 + \bar{U}_t, k, 0)$ converges to the resolvent $R(U_0, \bar{U}_0, k, 0)$. In general we shall see that this limit depends on the inverse of the operators $\mathbb{1} - \begin{pmatrix} U_{t=1} & \partial \\ -\bar{\partial} & \bar{U}_{t=1} \end{pmatrix} R_{\mathbb{R}^2}(0, 0, 0)$, where $R_{\mathbb{R}^2}(0, 0, 0)$ denotes the resolvent of the free Dirac operator on $\mathbb{R}^2$. In a next step the spectral theory of Dirac operators on $\mathbb{P}^1$ is used in order to investigate these inverse operators.

3.4.2 Spectral theory of Dirac operators on the sphere

For this purpose we use the covering $\mathbb{P}^1 = \mathbb{C} \cup \mathbb{P}^1 \setminus \{0\}$ and the affine coordinates z and $1/z$ on the members of this open covering. The spin bundle on $\mathbb{P}^1$ may be described as the trivial $\mathbb{C}^2$ bundles on both members of this covering together with the transition function $\begin{pmatrix} z & 0 \\ 0 & -\bar{z} \end{pmatrix}$, which transforms the sections on $\mathbb{C}$ into sections on $\mathbb{P}^1 \setminus \{0\}$. With respect to these

coordinates the free Dirac operator [Fr-1, Section 3.4] is equal to

$$(1 + z\bar{z}) \begin{pmatrix} 0 & \partial_z \\ -\bar{\partial}_z & 0 \end{pmatrix} \text{ on } \mathbb{C} \quad \text{and equal to}$$

$$(1 + z\bar{z}) \begin{pmatrix} z & 0 \\ 0 & -\bar{z} \end{pmatrix} \begin{pmatrix} 0 & \partial_z \\ -\bar{\partial}_z & 0 \end{pmatrix} \begin{pmatrix} z & 0 \\ 0 & -\bar{z} \end{pmatrix}^{-1} = (1 + 1/(z\bar{z})) \begin{pmatrix} 0 & \partial_{1/z} \\ -\bar{\partial}_{1/z} & 0 \end{pmatrix} \text{ on } \mathbb{P}^1 \setminus \{0\}.$$

We conclude that if $(1 + z\bar{z})U_z$ denotes the potential on $\mathbb{C}$, the corresponding potential on $\mathbb{P}^1 \setminus \{0\}$ is equal to $(1 + 1/(z\bar{z}))U_{1/z}$, where $U_{1/z}$ is given by

$$U_{1/z}(1/z) = \frac{1 + z\bar{z}}{1 + 1/(z\bar{z})} U_z(z) = z\bar{z}U_z(z).$$

The corresponding Dirac operators are given by

$$(1 + z\bar{z}) \begin{pmatrix} U_z & \partial_z \\ -\bar{\partial}_z & \bar{U}_z \end{pmatrix} \text{ on } \mathbb{C} \text{ and } (1 + 1/(z\bar{z})) \begin{pmatrix} U_{1/z} & \partial_{1/z} \\ -\bar{\partial}_{1/z} & \bar{U}_{1/z} \end{pmatrix} \text{ on } \mathbb{P}^1 \setminus \{0\}.$$

In particular, the integral over the density

$$U_z \bar{U}_z \frac{d\bar{z} \wedge dz}{2\sqrt{-1}} = U_{1/z} \bar{U}_{1/z} \frac{d(1/\bar{z}) \wedge d(1/z)}{2\sqrt{-1}}$$

is equal to the L^2 -norm of U_z on $\mathbb{C}$ (or of $U_{1/z}$ on $\mathbb{P}^1 \setminus \{0\}$). Therefore, all potentials $U \in L^2(\mathbb{R}^2)$ may be considered as potentials U_z of Dirac operators on $\mathbb{P}^1$. Due to Dolbeault's Lemma [Gu-Ro, Chapter I Section D 2. Lemma] the integral kernel of the corresponding free resolvent is equal to

$$\begin{pmatrix} 0 & (z - z')^{-1} \\ (\bar{z}' - \bar{z})^{-1} & 0 \end{pmatrix} \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}(1 + z'\bar{z}')} \text{ on } \mathbb{C} \quad \text{and equal to}$$

$$\begin{pmatrix} z & 0 \\ 0 & -\bar{z} \end{pmatrix} \begin{pmatrix} 0 & (z - z')^{-1} \\ (\bar{z}' - \bar{z})^{-1} & 0 \end{pmatrix} \begin{pmatrix} z' & 0 \\ 0 & -\bar{z}' \end{pmatrix}^{-1} \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}(1 + z'\bar{z}')}$$

$$= \begin{pmatrix} 0 & (1/z - 1/z')^{-1} \\ (1/\bar{z}' - 1/\bar{z})^{-1} & 0 \end{pmatrix} \frac{d(1/\bar{z}') \wedge d(1/z')}{2\pi\sqrt{-1}(1 + 1/(z'\bar{z}'))} \text{ on } \mathbb{P}^1 \setminus \{0\}.$$

We define L^p -spinors as $\mathbb{C}^2$ -valued functions on $\mathbb{C}$, whose components have bounded norm

$$\|f\| = \left(\int_{\mathbb{R}^2} |f(z)|^p (1 + z\bar{z})^{(\frac{p}{2}-2)} d^2x \right)^{1/p}.$$

Due to the transformation property of spinors this norm for spinors on $\mathbb{C}$ is equal to the norm of the corresponding spinors on $\mathbb{P}^1 \setminus \{0\}$ with respect to the coordinates $1/z$. Since the multiplication with $(1 + z\bar{z})^{\frac{1}{2}-\frac{2}{p}}$ is an isometry from the space of L^p -spinors onto $L^p(\mathbb{R}^2) \times$

$L^p(\mathbb{R}^2)$, the free resolvent, $R_{\mathbb{P}^1}(0, 0, 0)$ considered as an operator from $L^p(\mathbb{R}^2) \times L^p(\mathbb{R}^2)$ ($\simeq L^p$ -spinors) into $L^q(\mathbb{R}^2) \times L^q(\mathbb{R}^2)$ ($\simeq L^q$ -spinors), has the integral kernel

$$\begin{pmatrix} 0 & (z - z')^{-1} \\ (\bar{z}' - \bar{z})^{-1} & 0 \end{pmatrix} (1 + z\bar{z})^{\frac{1}{2} - \frac{2}{q}} (1 + z'\bar{z}')^{\frac{2}{p} - \frac{3}{2}} \frac{d^2 x'}{\pi}.$$

If $\frac{1}{q} = \frac{1}{p} - \frac{1}{2}$, then the potential U represents the potential $(1 + z\bar{z})U$ on $\mathbb{P}^1$ considered as an operator from $L^q(\mathbb{R}^2) \times L^q(\mathbb{R}^2)$ ($\simeq L^q$ -spinors) into $L^p(\mathbb{R}^2) \times L^p(\mathbb{R}^2)$ ($\simeq L^p$ -spinors). Therefore, in this case the resolvent of the Dirac operator on $\mathbb{P}^1$ with potential $(1 + z\bar{z}) \begin{pmatrix} U & 0 \\ 0 & \bar{U} \end{pmatrix}$ considered as an operator from $L^p(\mathbb{R}^2) \times L^p(\mathbb{R}^2)$ ($\simeq L^p$ -spinors) into $L^q(\mathbb{R}^2) \times L^q(\mathbb{R}^2)$ ($\simeq L^q$ -spinors) is equal to

$$R_{\mathbb{P}^1}(U, \bar{U}, \lambda) = R_{\mathbb{P}^1}(0, 0, \lambda) \left(\mathbf{1} - \begin{pmatrix} U & 0 \\ 0 & \bar{U} \end{pmatrix} R_{\mathbb{P}^1}(0, 0, \lambda) \right)^{-1}.$$

Fortunately, if $p = \frac{4}{3}$ and $q = 4$, the resolvent $R_{\mathbb{P}^1}$ considered as an operator from $L^{\frac{4}{3}}(\mathbb{R}^2) \times L^{\frac{4}{3}}(\mathbb{R}^2)$ into $L^4(\mathbb{R}^2) \times L^4(\mathbb{R}^2)$ is equal to the resolvent $R_{\mathbb{R}^2}(0, 0, 0)$ of the free Dirac operator on $\mathbb{R}^2$.

In order to carry over the spectral theory developed in Section 2.1 we shall estimate the free resolvent $R_{\mathbb{P}^1}(0, 0, \sqrt{-1}\lambda)$ in the limit $\lambda \rightarrow \infty$. Due to the invariance of the Dirac operator under the action of $SU(2, \mathbb{C})$ on $\mathbb{P}^1$, it suffices to estimate the Green's function $\begin{pmatrix} a_{\mathbb{P}^1, \lambda}(x) & b_{\mathbb{P}^1, \lambda}(x) \\ c_{\mathbb{P}^1, \lambda}(x) & d_{\mathbb{P}^1, \lambda}(x) \end{pmatrix}$, which satisfies the equation

$$\begin{pmatrix} \sqrt{-1}\lambda & -(1 + z\bar{z})\partial \\ (1 + z\bar{z})\bar{\partial} & \sqrt{-1}\lambda \end{pmatrix} \begin{pmatrix} a_{\mathbb{P}^1, \lambda}(x) & b_{\mathbb{P}^1, \lambda}(x) \\ c_{\mathbb{P}^1, \lambda}(x) & d_{\mathbb{P}^1, \lambda}(x) \end{pmatrix} = \begin{pmatrix} \delta(x) & 0 \\ 0 & \delta(x) \end{pmatrix}.$$

Here $\delta(x)$ denotes the two-dimensional δ -function on $\mathbb{R}^2$, i. e. $\delta(x)d^2x$ denotes the point measure at $x = 0$ of mass 1. Due to the relation

$$\begin{aligned} & \begin{pmatrix} \sqrt{-1}\lambda & -(1 + z\bar{z})\partial \\ (1 + z\bar{z})\bar{\partial} & \sqrt{-1}\lambda \end{pmatrix} \begin{pmatrix} \sqrt{-1}\lambda & (1 + z\bar{z})\partial \\ -(1 + z\bar{z})\bar{\partial} & \sqrt{-1}\lambda \end{pmatrix} \\ &= \begin{pmatrix} -\lambda^2 + (1 + z\bar{z})^2 \partial \bar{\partial} + (1 + z\bar{z}) \bar{z} \bar{\partial} & 0 \\ 0 & -\lambda^2 + (1 + z\bar{z})^2 \bar{\partial} \partial + (1 + z\bar{z}) z \partial \end{pmatrix} \end{aligned}$$

it suffices to calculate the Green's functions of the two operators in the diagonal entries of the right hand side. These Green's functions depend only on $r = \sqrt{z\bar{z}} = \sqrt{x_1^2 + x_2^2}$. In fact, on functions depending only on r , both operators in the diagonal entries of the right hand side act as the operator

$$-\lambda^2 + \left(\frac{1 + r^2}{2} \right)^2 \left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} \right) + \frac{1 + r^2}{2} r \frac{d}{dr}.$$

The substitution $r = \tan(\varphi/2)$ transforms this operator into

$$-\lambda^2 + \frac{d^2}{d\varphi^2} + \frac{1}{\sin(\varphi)} \frac{d}{d\varphi}.$$

The two dimensional δ -function is equal to the one-dimensional δ -function with respect to the measure $\sin(\varphi)d\varphi$. Therefore, we shall calculate the unique Green's function $\mathcal{G}_{\mathbb{P}^1, \lambda}$ depending on $\varphi \in \mathbb{R}/(2\pi\mathbb{Z})$, which satisfies

$$-\lambda^2 \sin(\varphi) \mathcal{G}_{\mathbb{P}^1, \lambda}(\varphi) + \sin(\varphi) \mathcal{G}_{\mathbb{P}^1, \lambda}''(\varphi) + \mathcal{G}_{\mathbb{P}^1, \lambda}'(\varphi) = \delta(\varphi),$$

where δ denotes the one-dimensional δ -function on $\varphi \in \mathbb{R}/(2\pi\mathbb{Z})$ with respect to the measure $d\varphi$. Obviously this Green's function $\mathcal{G}_{\mathbb{P}^1, \lambda}$ is odd and has a Fourier series of the form

$$\mathcal{G}_{\mathbb{P}^1, \lambda}(\varphi) = \sum_{n \in \mathbb{N}} a_n \sin(n\varphi).$$

Since the Fourier series of the δ -functions is equal to

$$\delta(\varphi) = \frac{1}{2\pi} + \frac{1}{\pi} \sum_{n \in \mathbb{N}} \cos(n\varphi),$$

the coefficients a_n satisfy the recursion relation

$$a_{n+1} = \frac{2na_n + (\lambda^2 + (n-1)^2)a_{n-1} - 2/\pi}{\lambda^2 + (n+1)^2} \text{ for } n \in \mathbb{N} \text{ with } a_1 = -\frac{1}{\pi(\lambda^2 + 1)} \text{ and } a_0 = 0.$$

Lemma 3.32. *If $\lambda \in \mathbb{R}$, then the functions $(1 - \cos(\varphi))\mathcal{G}_{\mathbb{P}^1, \lambda}(\varphi)$ and $(1 - \cos(\varphi))\mathcal{G}_{\mathbb{P}^1, \lambda}'(\varphi)$ are continuous functions depending on $\varphi \in \mathbb{R}/(2\pi\mathbb{Z})$, which converge uniformly to zero in the limit $\lambda \rightarrow \infty$.*

Proof. The functions $(1 - \cos(\varphi))\mathcal{G}_{\mathbb{P}^1, \lambda}(\varphi)$ and $(1 - \cos(\varphi))\mathcal{G}_{\mathbb{P}^1, \lambda}'(\varphi)$ have the Fourier series

$$\begin{aligned} (1 - \cos(\varphi))\mathcal{G}_{\mathbb{P}^1, \lambda}(\varphi) &= \sum_{n \in \mathbb{N}} \frac{2a_n - a_{n-1} - a_{n+1}}{2} \sin(n\varphi) \text{ and} \\ (1 - \cos(\varphi))\mathcal{G}_{\mathbb{P}^1, \lambda}'(\varphi) &= -\frac{a_1}{2} + \sum_{n \in \mathbb{N}} \frac{2na_n - (n-1)a_{n-1} - (n+1)a_{n+1}}{2} \cos(n\varphi), \end{aligned}$$

respectively. It suffices to show that the sum of $|2a_n - a_{n-1} - a_{n+1}|$ and the sum of $|2na_n - (n-1)a_{n-1} - (n+1)a_{n+1}|$ exist and converge in the limit $\lambda \rightarrow \infty$ to zero. We shall estimate the absolute values of the coefficients a_n .

1. First we claim that for all $1 < \alpha < 2$ there exists a constant $C_\alpha > 0$, such that all absolute values $|a_n|$ are bounded by $C_\alpha \frac{n^\alpha}{\lambda^2 + n^2}$. In fact, due to the recursion relation it suffices to show the inequalities

$$C_\alpha \frac{2n^{(1+\alpha)}}{\lambda^2 + n^2} + C_\alpha(n-1)^\alpha + \frac{2}{\pi} \leq C_\alpha(n+1)^\alpha \quad \forall n \in \mathbb{N} \setminus \{1\},$$

$$|a_1| \leq C_\alpha \frac{1}{\lambda^2 + 1} \text{ and } |a_2| \leq C_\alpha \frac{2^\alpha}{\lambda^2 + 4}.$$

Since $a_2 = -\frac{2}{\pi(\lambda^2+4)}$, the latter inequalities are equivalent to $\frac{1}{\pi} \leq C_\alpha$ and $\frac{2}{\pi} \leq C_\alpha 2^\alpha$. Obviously we have for all $1 < \alpha < 2$ and all $n \in \mathbb{N}$

$$(n-1)^\alpha \leq \left(1 - \frac{2}{n+1}\right)^\alpha (n+1)^\alpha \leq \left(1 - \frac{2}{n+1}\right) \left(1 - \frac{2(\alpha-1)}{n+1}\right) (n+1)^\alpha \leq (n+1)^\alpha - 2\alpha(n+1)^{\alpha-1} + 4(\alpha-1)(n+1)^{\alpha-2}.$$

If we insert this inequality in the expression on the left hand side of the first inequality above, we obtain

$$\begin{aligned} C_\alpha \frac{2n^{(1+\alpha)}}{\lambda^2 + n^2} + C_\alpha(n-1)^\alpha + \frac{2}{\pi} &\leq \\ &\leq 2C_\alpha(n+1)^{\alpha-1} + C_\alpha((n+1)^\alpha - 2\alpha(n+1)^{\alpha-1} + 4(\alpha-1)(n+1)^{\alpha-2}) + \frac{2}{\pi}. \end{aligned}$$

Therefore, it suffices to establish the inequalities

$$\frac{1}{\pi} \leq C_\alpha(\alpha-1)(n+1)^{\alpha-1} \left(1 - \frac{2}{n+1}\right) \quad \forall n \in \mathbb{N} \setminus \{1\}, \quad \frac{1}{\pi} \leq C_\alpha \text{ and } \frac{2}{\pi} \leq C_\alpha 2^\alpha.$$

Obviously there exist such $C_\alpha < \infty$ for all $1 < \alpha < 2$.

2. In a second step we claim that for all $1 < \alpha < 2$ there exists a constant C'_α , such that the absolute values $|a_{n+1} - a_{n-1}|$ are bounded by $C'_\alpha \frac{n^{\alpha+1}}{(\lambda^2 + (n+1)^2)(\lambda^2 + (n-1)^2)}$. In fact, the recursion relation implies

$$|a_{n+1} - a_{n-1}| \leq \frac{2n}{\lambda^2 + (n+1)^2} (|a_n| + 2|a_{n-1}|).$$

Hence the second claim follows from what has already been proven, with $C'_\alpha = 6C_\alpha$.

3. In a third step we prove that for all $1 < \alpha < 2$ and $|\lambda| \geq \sqrt{2}$ (this assumption on λ is not essential and can be removed) there exists a constant C''_α such that the absolute values $|2a_n - a_{n+1} - a_{n-1}|$ are bounded by $C''_\alpha \frac{n^\alpha}{(\lambda^2 + (n+1)^2)(\lambda^2 + (n-1)^2)}$. To see this set $b_n = 2a_n - a_{n+1} - a_{n-1}$ for $n \in \mathbb{N}$. If we subtract from two times the recursion relation above the two relations, which are obtained by reducing and enlarging the indices by one, then we obtain the following recursion relation:

$$\begin{aligned} b_{n+1} &= \frac{-2nb_n + (\lambda^2 + (n-2)^2)b_{n-1} + 4(a_{n+1} - a_{n-1})}{\lambda^2 + (n+2)^2} \quad \forall n \in \mathbb{N} \setminus \{1\} \\ \text{with } b_1 &= -\frac{6}{\pi(\lambda^2 + 1)(\lambda^2 + 4)} \text{ and } b_2 = -\frac{4(\lambda^2 - 5)}{\pi(\lambda^2 + 1)(\lambda^2 + 4)(\lambda^2 + 9)}. \end{aligned}$$

Therefore, it suffices to show for all $n \in \mathbb{N} \setminus \{1\}$

$$C''_\alpha \frac{2n^{(\alpha+1)}}{(\lambda^2 + (n-1)^2)(\lambda^2 + (n+1)^2)} + C''_\alpha \frac{(n-1)^\alpha}{\lambda^2 + n^2} + 4|a_{n+1} - a_{n-1}| \leq C''_\alpha \frac{(n+1)^\alpha}{\lambda^2 + n^2}$$

together with the two inequalities $\frac{6}{\pi} \leq C''_\alpha$ and $\frac{4|\lambda^2-5|}{\pi(\lambda^2+4)} \leq C''_\alpha 2^\alpha$. If $\lambda^2 \geq 2$, then we have

$$\frac{(\lambda^2 + n^2)n^2}{(\lambda^2 + (n-1)^2)(\lambda^2 + (n+1)^2)} \leq 1 - \frac{\lambda^4 + \lambda^2(n^2 + 2) - 2n^2 + 1}{(\lambda^2 + (n-1)^2)(\lambda^2 + (n+1)^2)} \leq 1.$$

With the help of the estimates in the first and the second step we arrive at the estimates

$$2C'_\alpha n^{\alpha-1} \leq 2C'_\alpha (n+1)^{\alpha-1} \leq C''_\alpha (\alpha-1)(n+1)^{\alpha-1} \left(1 - \frac{2}{n+1}\right) \quad \forall n \in \mathbb{N} \setminus \{1\},$$

$$\frac{6}{\pi} \leq C''_\alpha \text{ and } \frac{4|\lambda^2-5|}{\pi(\lambda^2+4)} \leq C''_\alpha 2^\alpha.$$

Again it is obvious that there exist such C''_α .

Due to the proven third claim the sum $\sum_{n \in \mathbb{N}} |b_n|$ exists and converges in the limit $\lambda \rightarrow \infty$ to zero. Due to the proven second and the third claim the sum

$$\sum_{n \in \mathbb{N}} |2na_n - (n+1)a_{n+1} - (n-1)a_{n-1}| \leq \sum_{n \in \mathbb{N}} n|b_n| + \sum_{n \in \mathbb{N}} |a_{n+1} - a_{n-1}|$$

exists and converges in the limit $\lambda \rightarrow \infty$ to zero.

q.e.d.

The Green's functions of the Dirac operator on $\mathbb{P}^1$ may be expressed in terms of $\mathcal{G}_{\mathbb{P}^1, \lambda}$ as

$$\begin{pmatrix} a_{\mathbb{P}^1, \lambda}(x) & b_{\mathbb{P}^1, \lambda}(x) \\ c_{\mathbb{P}^1, \lambda}(x) & d_{\mathbb{P}^1, \lambda}(x) \end{pmatrix} = \begin{pmatrix} \sqrt{-1}\lambda \mathcal{G}_{\mathbb{P}^1, \lambda}(2 \arctan(\sqrt{z\bar{z}})) & (1+z\bar{z})\partial \mathcal{G}_{\mathbb{P}^1, \lambda}(2 \arctan(\sqrt{z\bar{z}})) \\ -(1+z\bar{z})\bar{\partial} \mathcal{G}_{\mathbb{P}^1, \lambda}(2 \arctan(\sqrt{z\bar{z}})) & \sqrt{-1}\lambda \mathcal{G}_{\mathbb{P}^1, \lambda}(2 \arctan(\sqrt{z\bar{z}})) \end{pmatrix}.$$

Hence, due to the invariance under the action of $SU(2, \mathbb{C})$ on $\mathbb{P}^1$, this lemma implies that outside of the diagonal the integral kernel of the resolvent $\begin{pmatrix} \sqrt{-1}\lambda & -(1+z\bar{z})\partial \\ (1+z\bar{z})\bar{\partial} & \sqrt{-1}\lambda \end{pmatrix}^{-1}$ is a continuous function, which converges in the limit $\lambda \rightarrow \infty$ on the complement of arbitrary small discs (with respect to the invariant metric of $\mathbb{P}^1$) uniformly to zero. Therefore, the arguments of Theorem 2.3 and Lemma 2.6 carry over to the resolvents of Dirac operators on $\mathbb{P}^1$.

Corollary 3.33. *The resolvents define meromorphic maps $(U, \lambda) \mapsto R_{\mathbb{P}^1}(U, \bar{U}, \lambda)$ from the Hilbert space $L^2(\mathbb{R}^2) \times \mathbb{C}$ into the compact operators on the L^2 -spinors. On all subsets of $L^2(\mathbb{R}^2)$, where for some $C_p < S_p^{-1}$ and $\varepsilon > 0$ the L^2 -norms of the restrictions of the potentials to all ε -balls of $\mathbb{C} \subset \mathbb{P}^1$ (with respect to the natural metric of $\mathbb{P}^1$) are not larger than C_p , this map is weakly continuous. Finally, for $1 < p < 2$ the operators $\mathbf{1} - \begin{pmatrix} U & 0 \\ 0 & \bar{U} \end{pmatrix} R_{\mathbb{P}^1}(0, 0, \lambda)$ on $L^p(\mathbb{R}^2) \times L^p(\mathbb{R}^2)$ are Fredholm operators of index zero, and invertible if and only if λ is no eigenvalue of the corresponding Dirac operator on $\mathbb{P}^1$.*

Proof. It remains to prove the last statement. Due to the first resolvent formula [R-S-I, Theorem VI.5] for all vectors ψ and all λ and λ' in the complement of the spectrum the corresponding resolvents R_λ and $R_{\lambda'}$ satisfy the equation

$$R_{\lambda'}(\psi + (\lambda' - \lambda)R_\lambda\psi) = R_\lambda\psi.$$

We conclude that the range of the resolvent R_λ does not depend on λ as long as λ does not belong to the spectrum. Since the Riesz–Schauder Theorem is also true for compact operators on Banach spaces [Yo, Chapter X.5.], the Dirac operators on $\mathbb{P}^1$ may be considered as closed operators on the space of L^p -spinors with compact resolvents. The inverse mapping theorem [R-S-I, Theorem III.11] together with the formula

$$R_{\mathbb{P}^1}(U, \bar{U}, \lambda) = R_{\mathbb{P}^1}(0, 0, \lambda) \left(\mathbf{1} - \begin{pmatrix} U & 0 \\ 0 & \bar{U} \end{pmatrix} R_{\mathbb{P}^1}(0, 0, \lambda) \right)^{-1}$$

for the resolvents of these Dirac operators acting on the L^p -spinors shows that the operator $\mathbf{1} - \begin{pmatrix} U & 0 \\ 0 & \bar{U} \end{pmatrix} R_{\mathbb{P}^1}(0, 0, \lambda)$ on the L^p spinors is invertible if and only if λ does not belong to the spectrum of the corresponding Dirac operator. This follows also from the following variant of the first resolvent formula [R-S-I, Theorem VI.5]:

$$(\lambda - \lambda') R_{\lambda'} = R_\lambda \left(\frac{\mathbf{1}}{\lambda - \lambda'} - R_\lambda \right)^{-1} = \left(\frac{\mathbf{1}}{\lambda - \lambda'} - R_\lambda \right)^{-1} R_\lambda,$$

since, due to the arguments of the proof of Lemma 2.6 for all $U \in L^2(\mathbb{R}^2)$ and large λ the operators $\mathbf{1} - \begin{pmatrix} U & 0 \\ 0 & \bar{U} \end{pmatrix} R_{\mathbb{P}^1}(0, 0, \sqrt{-1}\lambda)$ on $L^p(\mathbb{R}^2) \times L^p(\mathbb{R}^2)$ are invertible. Since the range of the spectral projections $\check{P}$ are contained in the domain, the operators

$$\check{Q} = R_{\mathbb{P}^1}^{-1}(0, 0, \lambda) \check{P} R_{\mathbb{P}^1}(0, 0, \lambda)$$

are well defined projections, whenever λ does not belong to the spectrum of the free Dirac operator. In general the operators $\mathbf{1} - \begin{pmatrix} U & 0 \\ 0 & \bar{U} \end{pmatrix} R_{\mathbb{P}^1}(0, 0, \lambda)$ are invertible operators from the range of the projection $\mathbf{1} - \check{Q}$ onto the range of the projection $\mathbf{1} - \check{P}$. Since the spectral projections of the Dirac operators on $\mathbb{P}^1$ are finite-dimensional, these operators are Fredholm operators of index zero, and the kernel and the co-kernel coincides with the corresponding eigenspaces and transposed eigenspaces, respectively. **q.e.d.**

Remark 3.34. In particular, for $p = \frac{4}{3}$ and $q = 4$ and for all $U \in L^2(\mathbb{R}^2)$ the operators $\mathbf{1} - \begin{pmatrix} U & 0 \\ 0 & \bar{U} \end{pmatrix} R_{\mathbb{R}^2}(0, 0, 0)$ on $L^{\frac{4}{3}}(\mathbb{R}^2) \times L^4(\mathbb{R}^2)$ are also Fredholm operators of index zero, and invertible, if the corresponding Dirac operator on $\mathbb{P}^1$ has a trivial kernel. The relation of these kernels to the spectral theory of the Dirac operators on $\mathbb{R}^2$ is not so obvious.

3.4.3 An illustrative family of potentials

Now we are prepared to treat the limits $t \rightarrow \infty$ of the resolvents $R(U_t, \bar{U}_t, k, 0)$ of the family $(U_t)_{t \in [1, \infty)}$ of potentials of the form $U_t(z) = tU_{t=1}(tz)$, where $U_{t=1}$ has support in a ball $B(0, \varepsilon)$ contained in Δ .

Lemma 3.35. If the Dirac operator $(1 + z\bar{z}) \begin{pmatrix} U_{t=1} & \partial \\ -\bar{\partial} & \bar{U}_{t=1} \end{pmatrix}$ on $\mathbb{P}^1$ does not have a kernel, then the resolvents $R(U_t, \bar{U}_t, k, 0)$ considered as operators from $L^p(\Delta) \times L^p(\Delta)$ into $L^{q'}(\Delta) \times L^{q'}(\Delta)$ with $1 < p \leq 2$ and $2 \leq q' < q = \frac{2p}{2-p}$ converges in the limit $t \rightarrow \infty$ to $R(0, 0, k, 0)$. In general,

for all $\varepsilon > 0$ the restrictions of these resolvents to the complement of $B(0, \varepsilon)$ considered as operators from $L^p(\Delta \setminus B(0, \varepsilon)) \times L^p(\Delta \setminus B(0, \varepsilon))$ into $L^{\frac{2p}{2-p}}(\Delta \setminus B(0, \varepsilon)) \times L^{\frac{2p}{2-p}}(\Delta \setminus B(0, \varepsilon))$ with $1 < p < 2$ converges in the limit $t \rightarrow \infty$ to a Finite rank Perturbation 3.22 of $R(0, 0, k, 0)$. Moreover, the rank of the perturbation is bounded by the dimension of the kernel of the Dirac operator $(1 + z\bar{z}) \begin{pmatrix} U_{t=1} & \partial \\ -\bar{\partial} & \bar{U}_{t=1} \end{pmatrix}$ on $\mathbb{P}^1$, and the support of this perturbation is contained in $0 \in \Delta$.

Remark 3.36. If $\mathcal{B}$ is a subset of $\mathcal{A}$, then the functions on $\mathcal{B}$ may be considered as functions on $\mathcal{A}$, which vanish on the relative complement of $\mathcal{B}$. Vice versa, the restriction to $\mathcal{B}$ of any function on $\mathcal{A}$ yields a function on $\mathcal{B}$. For a given operator on a space of functions on $\mathcal{A}$, we call the operator, which associates to any function on $\mathcal{B}$ the restriction to $\mathcal{B}$ of the action of the operator on this function considered as a function on $\mathcal{A}$, the restriction of the operator to $\mathcal{B}$ or to the space of functions on $\mathcal{B}$.

Proof. We proof this within three steps. In the first step we investigate the family of resolvents transformed under the scaling transformations $z \rightarrow tz$. The limit $t \rightarrow \infty$ of these transformations is a kind of blowing up. Consequently, it yields a Dirac operator on $\mathbb{P}^1$. In the second step we show that if this blown up Dirac operator on $\mathbb{P}^1$ has a trivial kernel, then the family of resolvents $R(U_t, \bar{U}, k, 0)$ converge to the free resolvents $R(0, 0, k, 0)$. Finally, in the third step we show that a kernel of the Dirac operator blown up to $\mathbb{P}^1$ yields a *Finite rank Perturbation* 3.22 of the free resolvents.

1. The family of transformed resolvents. For all $t \in \mathbb{C}$ the transformations

$$l_{p,t} : f \mapsto l_{p,t}f \text{ with } (l_{p,t}f)(z) = |t|^{2/p}f(tz)$$

define isometries from $L^p(\mathbb{R}^2)$ onto $L^p(\mathbb{R}^2)$ and from $L^p(B(0, \varepsilon))$ onto $L^p(B(0, t^{-1}\varepsilon))$. Due to the explicit form of the integral kernel of $R(0, 0, k, 0)$ in Lemma 3.20 this resolvent is the convolution with a meromorphic and a anti-meromorphic function, respectively. The Laurent series of these functions around $z = 0$ converge on small discs. Therefore, for all $k \notin \mathcal{F}(0, 0)$ the restrictions of the operators

$$\begin{pmatrix} l_{\frac{2p}{2-p},t} & 0 \\ 0 & l_{\frac{2p}{2-p},t} \end{pmatrix}^{-1} R(0, 0, k, 0) \begin{pmatrix} l_{p,t} & 0 \\ 0 & l_{p,t} \end{pmatrix}$$

to $L^p(B(0, \varepsilon)) \times L^p(B(0, \varepsilon))$ define a real analytic family of operators depending on the parameter $t \in [1, \infty)$. Furthermore, since in the limit $t \rightarrow \infty$ only the pole of the convolution function at $z = 0$ contributes, this family converges in this limit to the restriction of the operator $R_{\mathbb{R}^2}(0, 0, 0)$ to $L^p(B(0, \varepsilon)) \times L^p(B(0, \varepsilon))$. Hence, due to Lemma 3.20, this family extends to a holomorphic family

$$\frac{t}{|t|} \begin{pmatrix} l_{\frac{2p}{2-p},t} & 0 \\ 0 & l_{\frac{2p}{2-p},t} \end{pmatrix}^{-1} R(0, 0, k, 0) \begin{pmatrix} l_{p,\bar{t}} & 0 \\ 0 & l_{p,t} \end{pmatrix}$$

depending on $\{t \in \mathbb{P}^1 \mid |t| \geq 1\}$. More precisely, if

$$\mathcal{K}_1(z, k) = z^{-1} + \sum_{n=0}^{\infty} \mathcal{K}_{1,n} z^n \quad \text{and} \quad \mathcal{K}_2(z, k) = \bar{z}^{-1} + \sum_{n=0}^{\infty} \mathcal{K}_{2,n} \bar{z}^n$$

denotes the Laurent series of the two functions $\mathcal{K}_1$ and $\mathcal{K}_2$ in the integral kernel of $R(0, 0, k, 0)$ (compare with Lemma 3.20), then the Taylor series at $t = \infty$ of the integral kernel of this family is given by

$$\begin{pmatrix} 0 & (z - z')^{-1} \\ (\bar{z}' - \bar{z})^{-1} & 0 \end{pmatrix} \frac{d^2 x'}{\pi} + \sum_{n \in \mathbb{N}} t^{-n} \begin{pmatrix} 0 & \mathcal{K}_{1,n-1} (z - z')^{n-1} \\ -\mathcal{K}_{2,n-1} (\bar{z} - \bar{z}')^{n-1} & 0 \end{pmatrix} \frac{d^2 x'}{\pi}.$$

Since the convolution with a polynomial with respect to z is a finite rank operator, with the exception of the zero-th Taylor coefficients of this family at $t = \infty$ (which is equal to the restriction of $R_{\mathbb{R}^2}(0, 0, 0)$ to $B(0, \varepsilon)$) all Taylor coefficients at $t = \infty$ are finite rank operators from $L^p(B(0, \varepsilon)) \times L^p(B(0, \varepsilon))$ to $L^{\frac{2p}{2-p}}(B(0, \varepsilon)) \times L^{\frac{2p}{2-p}}(B(0, \varepsilon))$.

Therefore, we have also a real analytic family of operators

$$\begin{pmatrix} \mathbf{I}_{p,t} & 0 \\ 0 & \mathbf{I}_{p,t} \end{pmatrix}^{-1} \left(\mathbf{1} - \begin{pmatrix} U_t & 0 \\ 0 & \bar{U}_t \end{pmatrix} R(0, 0, k, 0) \right) \begin{pmatrix} \mathbf{I}_{p,t} & 0 \\ 0 & \mathbf{I}_{p,t} \end{pmatrix}$$

on $L^p(B(0, \varepsilon)) \times L^p(B(0, \varepsilon))$ depending on $t \in [1, \infty)$, which extends to a holomorphic family

$$\begin{aligned} \mathbf{1} - \frac{t}{|t|} \begin{pmatrix} U_{t=1} & 0 \\ 0 & \bar{U}_{t=1} \end{pmatrix} \begin{pmatrix} \mathbf{I}_{\frac{2p}{2-p}, t} & 0 \\ 0 & \mathbf{I}_{\frac{2p}{2-p}, \bar{t}} \end{pmatrix}^{-1} R(0, 0, k, 0) \begin{pmatrix} \mathbf{I}_{p, \bar{t}} & 0 \\ 0 & \mathbf{I}_{p, t} \end{pmatrix} = \\ \begin{pmatrix} \mathbf{I}_{p, \bar{t}} & 0 \\ 0 & \mathbf{I}_{p, t} \end{pmatrix}^{-1} \left(\mathbf{1} - \begin{pmatrix} U_t & 0 \\ 0 & \bar{U}_t \end{pmatrix} R(0, 0, k, 0) \right) \begin{pmatrix} \mathbf{I}_{p, \bar{t}} & 0 \\ 0 & \mathbf{I}_{p, t} \end{pmatrix} \end{aligned}$$

depending on $\{t \in \mathbb{P}^1 \mid |t| \geq 1\}$, if we extend the family U_t by the formula

$$U_t = \mathbf{I}_{p, \bar{t}} \frac{t}{|t|} U_{t=1} \mathbf{I}_{\frac{2p}{2-p}, t}^{-1} \quad \bar{U}_t = \mathbf{I}_{p, t} \frac{t}{|t|} \bar{U}_{t=1} \mathbf{I}_{\frac{2p}{2-p}, \bar{t}}^{-1}.$$

This definition does not depend on $p \in (1, 2)$. If t is not real, then these operators are not multiplication operators and therefore non-local, but for real t this definition coincides with the definition of U_t above. At $t = \infty$ the value of this family is equal to the restriction of the operator $\mathbf{1} - \begin{pmatrix} U_{t=1} & 0 \\ 0 & \bar{U}_{t=1} \end{pmatrix} R_{\mathbb{R}^2}(0, 0, 0)$ to $L^p(B(0, \varepsilon)) \times L^p(B(0, \varepsilon))$.

2. A trivial kernel of the blown up Dirac operator. Due to Corollary 3.33 the foregoing operator has an inverse, if and only if the Dirac operator $(1 + z\bar{z}) \begin{pmatrix} U_{t=1} & \partial \\ -\bar{\partial} & \bar{U}_{t=1} \end{pmatrix}$ on $\mathbb{P}^1$ does not have a kernel. At a first sight this is only true for $p = \frac{4}{3}$ and $q = 4$. But since the function $(1 + z\bar{z})$ and its inverse are bounded on the domain of $U_{t=1}$, this is true for all $1 < p < 2$. We conclude that in this case and for all sufficiently large real t and $k \notin \mathcal{F}(0, 0)$ the restrictions of the operators $\mathbf{1} - \begin{pmatrix} U_t & 0 \\ 0 & \bar{U}_t \end{pmatrix} R(0, 0, k, 0)$ to $L^p(B(0, t^{-1}\varepsilon)) \times L^p(B(0, t^{-1}\varepsilon))$ have an inverse,

which is bounded uniformly with respect to t . Now we apply Lemma 3.21 and conclude that under the same assumptions the resolvents $R(U_t, \bar{U}_t, k, 0)$ converge in the limit $t \rightarrow \infty$ to the corresponding limit of the operators

$$R(0, 0, k, 0) + R(0, 0, k, 0) \left(\mathbf{1} - \begin{pmatrix} U_t & 0 \\ 0 & \bar{U}_t \end{pmatrix} R_{\mathbb{R}^2}(0, 0, 0) \right)^{-1} \begin{pmatrix} U_t & 0 \\ 0 & \bar{U}_t \end{pmatrix} R(0, 0, k, 0).$$

If in this formula we view the resolvent on the left hand side as a bounded operator from $L^{p'}(\Delta) \times L^{p'}(\Delta)$ into $L^{q'}(\Delta) \times L^{q'}(\Delta)$ with $1 < p' < 2$ and $p \leq q' \leq 2p'/(2 - p')$ and $k \notin \mathcal{F}(0, 0)$ (compare with Lemma 2.4), then the inverse operator may be considered as an operator on $L^{p'}(B(0, t^{-1}\varepsilon)) \times L^{p'}(B(0, t^{-1}\varepsilon))$ instead of $L^p(B(0, t^{-1}\varepsilon)) \times L^p(B(0, t^{-1}\varepsilon))$. If $p' < p$, then the restriction is an operator from $L^p(\Delta)$ into $L^{p'}(B(0, t^{-1}\varepsilon))$, which due to Hölder's inequality [R-S-I, Theorem III.1 (c)] is bounded by $(\pi\varepsilon^2)^{(p-p')/(pp')} |t|^{2(p'-p)/(pp')}$. Hence, if the Dirac operator $\begin{pmatrix} U_{t=1} & \partial \\ -\partial & \bar{U}_{t=1} \end{pmatrix}$ on $\mathbb{P}^1$ has no kernel, then in the limit $t \rightarrow \infty$ the resolvents $R(U_t, \bar{U}_t, k, 0)$ converge as operators from $L^p(\Delta) \times L^p(\Delta)$ into $L^{q'}(\Delta) \times L^{q'}(\Delta)$ with $1 < p \leq 2$ and $2 \geq q' < q = \frac{2p}{2-p}$ for all $k \notin \mathcal{F}(0, 0)$ in the limit $t \rightarrow \infty$ to the operator $R(0, 0, k, 0)$.

3. A non-trivial kernel of the blown up Dirac operator. In the third step we extend our arguments to the case, where the Dirac operator $(1 + z\bar{z}) \begin{pmatrix} U_{t=1} & \partial \\ -\partial & \bar{U}_{t=1} \end{pmatrix}$ on $\mathbb{P}^1$ has a non-trivial kernel. First we shall calculate the singular Laurent coefficients at $t = \infty$ of the family of inverse operators

$$\begin{pmatrix} \mathbf{l}_{p,\bar{t}} & 0 \\ 0 & \mathbf{l}_{p,t} \end{pmatrix}^{-1} \left(\mathbf{1} - \begin{pmatrix} U_t & 0 \\ 0 & \bar{U}_t \end{pmatrix} R(0, 0, k, 0) \right)^{-1} \begin{pmatrix} \mathbf{l}_{p,\bar{t}} & 0 \\ 0 & \mathbf{l}_{p,t} \end{pmatrix}$$

on $L^p(B(0, \varepsilon)) \times L^p(B(0, \varepsilon))$ of the holomorphic family introduced above. The original holomorphic family has a Taylor series at $t = \infty$, whose zero coefficient is equal to the restriction $\mathbf{A}$ of the operator $\mathbf{1} - \begin{pmatrix} U_{t=1} & 0 \\ 0 & \bar{U}_{t=1} \end{pmatrix} R_{\mathbb{R}^2}(0, 0, 0)$ to $L^p(B(0, \varepsilon)) \times L^p(B(0, \varepsilon))$, which due to Corollary 3.33 is a Fredholm operator of index zero. Furthermore, all other Taylor coefficients are finite rank operators, since the corresponding Taylor coefficients of the resolvents are finite rank operators. Finally, the difference $\mathbf{B}_t$ of the original holomorphic family minus $\mathbf{A}$ (the value at $t = \infty$) are compact operators.

Lemma 3.37. *Let $\mathbf{A}$ be a Fredholm operator of index zero from the Banach space $\mathbf{E}$ into the Banach space $\mathbf{G}$. Let $\mathbf{E}_{\text{finite}} \subset \mathbf{E}$ and $\mathbf{F}_{\text{finite}} \subset \mathbf{G}^*$ denote the kernel and co-kernel of $\mathbf{A}$, respectively. Furthermore, let $\mathbf{G}_{\text{infinite}}$ denote the range of $\mathbf{A}$ (i. e. the orthogonal complement of $\mathbf{F}_{\text{finite}}$). If $\mathbf{B}$ is another operator from $\mathbf{E}$ into $\mathbf{G}$ with the property, that the operator*

$$\mathbf{B}_{\text{finite}} : \mathbf{E}_{\text{finite}} \hookrightarrow \mathbf{E} \xrightarrow{\mathbf{B}} \mathbf{G} \hookrightarrow \mathbf{G}^{**} \twoheadrightarrow \mathbf{F}_{\text{finite}}^*$$

is invertible, then there exist unique decompositions

$$\mathbf{E} = \mathbf{E}_{\text{finite}} \oplus \mathbf{E}_{\text{infinite}} \quad \mathbf{G} = \mathbf{G}_{\text{finite}} \oplus \mathbf{G}_{\text{infinite}},$$

where $\mathbf{E}_{\text{infinite}}$ is the orthogonal complement of the image of $\mathbf{F}_{\text{finite}}$ under the operator $\mathbf{B}^ : \mathbf{G}^* \rightarrow \mathbf{E}^*$, and $\mathbf{G}_{\text{finite}}$ is the image of $\mathbf{E}_{\text{finite}}$ under the operator $\mathbf{B}$. Moreover, both operators $\mathbf{A}$ and $\mathbf{B}$ have diagonal block form with respect to these decompositions.*

Here the map $G^{**} \rightarrow F_{\text{finite}}^*$ on the right hand side in the definition of B_{finite} is the dual map of the natural inclusion $F_{\text{finite}} \hookrightarrow G^*$.

Proof. If B_{finite} is invertible, the operator

$$E \xrightarrow{B} G \hookrightarrow G^{**} \rightarrow F_{\text{finite}}^* \xrightarrow{B_{\text{finite}}^{-1}} E_{\text{finite}} \hookrightarrow E$$

is a projection onto E_{finite} . Since E_{infinite} is the kernel of the map

$$E \xrightarrow{B} G \hookrightarrow G^{**} \rightarrow F_{\text{finite}}^*,$$

this space is also the kernel of the former projection. Therefore, we have a decomposition $E = E_{\text{finite}} \oplus E_{\text{infinite}}$. On the other hand, the operator

$$G \hookrightarrow G^{**} \rightarrow F_{\text{finite}}^* \xrightarrow{B_{\text{finite}}^{-1}} E_{\text{finite}} \xrightarrow{B|_{E_{\text{finite}}}} G_{\text{finite}} \hookrightarrow G$$

is a projection onto G_{finite} . Since G_{infinite} is the orthogonal complement of F_{finite} , this subspace is the kernel of the latter projection, and we have a decomposition $G = G_{\text{finite}} \oplus G_{\text{infinite}}$.

Moreover, the operator A vanishes on E_{finite} and maps E_{infinite} onto G_{infinite} . By definition of G_{finite} , the operator B maps E_{finite} onto G_{finite} . Finally, since E_{infinite} is the kernel of the map

$$E \xrightarrow{B} G \hookrightarrow G^{**} \rightarrow F_{\text{finite}}^*,$$

the operator B maps E_{infinite} into G_{infinite} .

q.e.d.

This lemma implies that the operator $A + B$ is invertible, if B_{finite} is invertible, and the component of B from E_{infinite} into G_{infinite} has norm smaller than the inverse of the same component of A . In fact, in this case the direct sum of these two components are invertible. In the application we have in mind, A is equal to the value at $t = \infty$ of the family of Fredholm operators introduced above, and B_t is equal to the difference of this family minus the value at $t = \infty$. In the limit $t \rightarrow \infty$ the operators B_t become arbitrarily small.

Remark 3.38. *In the sequel we use a notation, which is slightly different from the notation used in Lemma 3.37. In fact, the spaces, which are denoted by capital F in Lemma 3.37, are in the sequel denoted by $R_{\mathbb{R}^2}^t(0, 0, 0)F$. The latter spaces F have the advantage, that the support of their elements is contained in the support of $U_{t=1}$.*

Let $E_{\text{finite}} \subset E = L^p(B(0, \varepsilon)) \times L^p(B(0, \varepsilon))$ denote the kernel of the restriction of the operator $\mathbf{1} - \begin{pmatrix} U_{t=1} & 0 \\ 0 & \bar{U}_{t=1} \end{pmatrix} R_{\mathbb{R}^2}(0, 0, 0)$ to $L^p(B(0, \varepsilon)) \times L^p(B(0, \varepsilon))$, and $F_{\text{finite}} \subset F = L^{\frac{2p}{3p-2}}(B(0, \varepsilon)) \times L^{\frac{2p}{3p-2}}(B(0, \varepsilon))$ the co-kernel of the restriction of the operator $\mathbf{1} - R_{\mathbb{R}^2}(0, 0, 0) \begin{pmatrix} U_{t=1} & 0 \\ 0 & \bar{U}_{t=1} \end{pmatrix}$ to $L^{\frac{2p}{2-p}}(B(0, \varepsilon)) \times L^{\frac{2p}{2-p}}(B(0, \varepsilon))$ (i. e. the kernel of the restriction of the operator $\mathbf{1} - \begin{pmatrix} U_{t=1} & 0 \\ 0 & \bar{U}_{t=1} \end{pmatrix} R_{\mathbb{R}^2}^t(0, 0, 0)$ to $L^{\frac{2p}{3p-2}}(B(0, \varepsilon)) \times L^{\frac{2p}{3p-2}}(B(0, \varepsilon))$). Obviously, the supports of the elements of E_{finite} and F_{finite} are contained in $B(0, \varepsilon)$. Therefore, these subspaces does not depend on ε , as long as $B(0, \varepsilon)$ contains the support of $U_{t=1}$. Furthermore, the resolvent

$R_{\mathbb{R}^2}(0, 0, 0)$ and the transposed resolvent map these subspaces E_{finite} and F_{finite} onto the kernel and the co-kernel of the corresponding Dirac operator on $\mathbb{P}^1$, respectively. Finally, due to the statement analogous to (iii) of Corollary 2.16, E_{finite} and F_{finite} are invariant under the anti-linear (projective) involutions $\psi \mapsto J\bar{\psi}$ and $\phi \mapsto J\bar{\phi}$, respectively. As we have seen in Corollary 2.17, these involutions cannot have fixed points, and the dimensions of E_{finite} and F_{finite} are even, say $2K$. Now let $n_1, \dots, n_J$ denote the unique sequence of numbers in $\mathbb{N}_0$ with the property,

that for all $j = 1, \dots, J$ n_j is the smallest index such that the span of $\begin{pmatrix} \bar{z}^{n_1} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ z^{n_1} \end{pmatrix}, \dots, \begin{pmatrix} \bar{z}^{n_j} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ z^{n_j} \end{pmatrix}$ does not contain any non-trivial element in the orthogonal complement of E_{finite} .

Furthermore, let $m_1, \dots, m_J$ be the analogous sequence with the property,

that for all $j = 1, \dots, J$ m_j is the smallest index such that the span of $\begin{pmatrix} z^{m_1} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ \bar{z}^{m_1} \end{pmatrix}, \dots, \begin{pmatrix} z^{m_j} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ \bar{z}^{m_j} \end{pmatrix}$ does not contain any non-trivial element in the orthogonal complement of F_{finite} .

In order to ensure the existence of these sequences we remark that for all $\psi \in E_{\text{finite}}$ and $\phi \in F_{\text{finite}}$ the Taylor coefficients at $z = \infty$ of the corresponding elements $R_{\mathbb{R}^2}(0, 0, 0)\psi$ and $R_{\mathbb{R}^2}^t(0, 0, 0)\phi$ in the kernel and the co-kernel of the corresponding Dirac operator on $\mathbb{P}^1$ are multiples of the numbers $\langle\langle \begin{pmatrix} \bar{z}^n \\ 0 \end{pmatrix}, \psi \rangle\rangle$, $\langle\langle \begin{pmatrix} 0 \\ z^n \end{pmatrix}, \psi \rangle\rangle$ and $\langle\langle \phi, \begin{pmatrix} z^n \\ 0 \end{pmatrix} \rangle\rangle$, $\langle\langle \phi, \begin{pmatrix} 0 \\ \bar{z}^n \end{pmatrix} \rangle\rangle$ with $n \in \mathbb{N}_0$, respectively. Hence, due to the *Strong unique continuation property* 2.10, not all of these numbers can vanish, and the sequences $n_1, \dots, n_J$ and $m_1, \dots, m_J$ are well defined (compare with Remark 3.28).

By definition of these sequences, the kernel E_{finite} and the co-kernel F_{finite} have unique basis $\psi_1, \dots, \psi_{2J}$ and $\phi_1, \dots, \phi_{2J}$ dual to the basis

$$\begin{pmatrix} \bar{z}^{n_1} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ z^{n_1} \end{pmatrix}, \dots, \begin{pmatrix} \bar{z}^{n_J} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ z^{n_J} \end{pmatrix} \text{ and } \begin{pmatrix} z^{m_1} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ \bar{z}^{m_1} \end{pmatrix}, \dots, \begin{pmatrix} z^{m_J} \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ \bar{z}^{m_J} \end{pmatrix}, \text{ respectively.}$$

Coming back to the notation of Section 3.4.1 let D be the divisor equal to $(\max\{n_J, m_J\} + 1)$ times the point 0 on $\mathbb{C}/\Lambda_{\mathbb{C}}$ and Ψ_D , Φ_D and $R_D(k)$ the corresponding matrices (compare with the first step of Section 3.4.1). Moreover, let N and M denote the following $(\max\{n_J, m_J\} + 1) \times J$ - and $J \times (\max\{n_J, m_J\} + 1)$ -matrices, whose entries are the following 2×2 -matrices indexed by $j = 1, \dots, J$ and $n, m = 0, \dots, \max\{n_J, m_J\}$:

$$N_{n,j} = \begin{cases} \mathbf{1} & \text{if } n = n_j \\ 0 & \text{if } n \neq n_j \end{cases} \text{ and } M_{j,m} = \begin{cases} \mathbf{1} & \text{if } m = m_j \\ 0 & \text{if } m \neq m_j \end{cases}, \text{ respectively.}$$

The transposed resolvent $R_{\mathbb{R}^2}^t(0, 0, 0)$ maps the dual of the Banach space $L^{\frac{2p}{2-p}}(B(0, \varepsilon)) \times L^{\frac{2p}{2-p}}(B(0, \varepsilon))$, which is isomorphic to $L^{\frac{2p}{3p-2}}(B(0, \varepsilon)) \times L^{\frac{2p}{3p-2}}(B(0, \varepsilon))$, onto the dual of the Banach space $L^p(B(0, \varepsilon)) \times L^p(B(0, \varepsilon))$, which is isomorphic to $L^{\frac{p}{p-1}}(B(0, \varepsilon)) \times L^{\frac{p}{p-1}}(B(0, \varepsilon))$. The range of the restriction of the operator $\mathbf{1} - \begin{pmatrix} U_{t=1} & 0 \\ 0 & \bar{U}_{t=1} \end{pmatrix} R_{\mathbb{R}^2}(0, 0, 0)$ to $L^p(B(0, \varepsilon)) \times$

$L^p(B(0, \varepsilon))$ is the orthogonal complement of $\mathbf{R}_{\mathbb{R}^2}^t(0, 0, 0)\mathbf{F}_{\text{finite}}$. Let $\mathbf{G}_{\text{infinite}} \subset L^{\frac{2p}{2-p}}(B(0, \varepsilon)) \times L^{\frac{2p}{2-p}}(B(0, \varepsilon))$ ($= \mathbf{G}$) denote this orthogonal complement. By construction the operator $\mathbf{1} - \begin{pmatrix} U_{t=1} & 0 \\ 0 & \bar{U}_{t=1} \end{pmatrix} \mathbf{R}_{\mathbb{R}^2}(0, 0, 0)$ has the kernel $\mathbf{E}_{\text{finite}}$ and the co-kernel $\mathbf{R}_{\mathbb{R}^2}^t(0, 0, 0)\mathbf{F}_{\text{finite}}$. With respect to the basis $\psi_1, \dots, \psi_{2J}$ and the dual basis of $\mathbf{R}_{\mathbb{R}^2}^t(0, 0, 0)\phi_1, \dots, \mathbf{R}_{\mathbb{R}^2}^t(0, 0, 0)\phi_{2J}$ the holomorphic family

$$\mathbf{B}_{t,\text{finite}} : \mathbf{E}_{\text{finite}} \hookrightarrow \mathbf{E} \xrightarrow{\mathbf{B}_t} \mathbf{G} \hookrightarrow \mathbf{G}^{**} \twoheadrightarrow (\mathbf{R}_{\mathbb{R}^2}^t(0, 0, 0)\mathbf{F}_{\text{finite}})^*$$

takes the matrix form:

$$\frac{-1}{t} \begin{pmatrix} t^{-m_1-n_1} \mathbf{R}_{m_1, n_1} & \dots & t^{-m_1-n_J} \mathbf{R}_{m_1, n_J} \\ \vdots & \dots & \vdots \\ t^{-m_J-n_1} \mathbf{R}_{m_J, n_1} & \dots & t^{-m_J-n_J} \mathbf{R}_{m_J, n_J} \end{pmatrix}$$

Here $\mathbf{R}_{m,n}$ denotes the entries of the matrix $\mathbf{R}_D(k)$. In the sequel we will assume that this matrix is invertible for $t \neq \infty$. This is equivalent to the statement that the determinant of the $J \times J$ -matrix $\mathbf{M}\mathbf{R}_D(k)\mathbf{N}$ does not vanish. In the eighth step of Section 3.4.1 we showed that the result of Section 2.2 carry over to these *complex Fermi curves*.

If $n_J = J - 1$ ($\iff n_j = j - 1 \ \forall j = 1, \dots, J$), then the projection onto $\mathbf{E}_{\text{finite}}$

$$\mathbf{E} \xrightarrow{\mathbf{B}_t} \mathbf{G} \hookrightarrow \mathbf{G}^{**} \twoheadrightarrow (\mathbf{R}_{\mathbb{R}^2}^t(0, 0, 0)\mathbf{F}_{\text{finite}})^* \xrightarrow{\mathbf{B}_{t,\text{finite}}^{-1}} \mathbf{E}_{\text{finite}} \hookrightarrow \mathbf{E}$$

has no singularity at $t = \infty$. Unfortunately for all other values of n_J it has a singularity. In fact, if $\psi \in \mathbf{E}$ has the property that

$$\langle \langle \begin{pmatrix} \bar{z}^n \\ 0 \end{pmatrix}, \psi \rangle \rangle = \begin{cases} 0 & \text{for } n = 0, \dots, N-1 \\ a & \text{for } n = N \end{cases} \quad \langle \langle \begin{pmatrix} 0 \\ z^n \end{pmatrix}, \psi \rangle \rangle = \begin{cases} 0 & \text{for } n = 0, \dots, N-1 \\ b & \text{for } n = N, \end{cases}$$

with $N \in \mathbb{N}_0$ and $(a, b) \in \mathbb{P}^1$, then the image of ψ under

$$\mathbf{E} \xrightarrow{\mathbf{B}_t} \mathbf{G} \hookrightarrow \mathbf{G}^{**} \twoheadrightarrow (\mathbf{R}_{\mathbb{R}^2}^t(0, 0, 0)\mathbf{F}_{\text{finite}})^*$$

has with respect to the dual basis of $\mathbf{R}_{\mathbb{R}^2}^t(0, 0, 0)\phi_1, \dots, \mathbf{R}_{\mathbb{R}^2}^t(0, 0, 0)\phi_{2J}$ and for large t the form

$$\left(\frac{1}{\pi t} + \mathbf{O}(t^{-2}) \right) \begin{pmatrix} \binom{m_1+N}{m_1} \frac{(-1)^N}{t^{m_1+N}} \begin{pmatrix} -b\mathcal{K}_{1, m_1+N} \\ a\mathcal{K}_{2, m_1+N} \end{pmatrix} \\ \vdots \\ \binom{m_J+N}{m_J} \frac{(-1)^N}{t^{m_J+N}} \begin{pmatrix} -b\mathcal{K}_{1, m_J+N} \\ a\mathcal{K}_{2, m_J+N} \end{pmatrix} \end{pmatrix}.$$

Obviously the inverse of $\mathbf{B}_{t,\text{finite}}$ applied on this vector can have a singularity at $t = \infty$, only if $N \in \{0, \dots, n_J\} \setminus \{n_1, \dots, n_J\}$. Analogously, if $m_J = J - 1$ ($\iff m_j = j - 1 \ \forall j = 1, \dots, J$), then the projection of $\mathbf{F}$ onto $\mathbf{F}_{\text{finite}}$ has no singularity at $t = \infty$.

We conclude that the resolvents of the family $(U_t, \bar{U}_{\bar{t}})$ differ from the free resolvent by

$$\begin{aligned} R(U_t, \bar{U}_{\bar{t}}, k, 0) - R(0, 0, k, 0) &= \\ &= R(0, 0, k, 0) \begin{pmatrix} l_{p, \bar{t}} & 0 \\ 0 & l_{p, t} \end{pmatrix} (A + B_t)^{-1} \begin{pmatrix} l_{p, \bar{t}} & 0 \\ 0 & l_{p, t} \end{pmatrix}^{-1} \begin{pmatrix} U_t & 0 \\ 0 & \bar{U}_{\bar{t}} \end{pmatrix} R(0, 0, k, 0) \\ &= R(0, 0, k, 0) \begin{pmatrix} l_{p, \bar{t}} & 0 \\ 0 & l_{p, t} \end{pmatrix} (A + B_t)^{-1} \begin{pmatrix} U_{t=1} & 0 \\ 0 & \bar{U}_{\bar{t}=1} \end{pmatrix} \begin{pmatrix} l_{\frac{2p}{2-p}, t} & 0 \\ 0 & l_{\frac{2p}{2-p}, \bar{t}} \end{pmatrix}^{-1} R(0, 0, k, 0). \end{aligned}$$

We shall calculate these differences in the limit $t \rightarrow \infty$ with the arguments used in the first two steps combined with Lemma 3.37. By definition of the sequence $n_1, \dots, n_J$ and due to Lemma 3.20 we have

$$\begin{aligned} R(0, 0, k, 0) \begin{pmatrix} l_{p, t} & 0 \\ 0 & l_{p, t} \end{pmatrix} \psi_{2j-1} &= \frac{|t|^{2/p} (-1)^{n_j}}{\pi n_j! t^{n_j+2}} \left(\begin{pmatrix} 0 \\ -\mathcal{K}_2^{(n_j)} \end{pmatrix} + \mathbf{O}(t^{-1}) \right) \text{ and} \\ R(0, 0, k, 0) \begin{pmatrix} l_{p, t} & 0 \\ 0 & l_{p, t} \end{pmatrix} \psi_{2j} &= \frac{|t|^{2/p} (-1)^{n_j}}{\pi n_j! t^{n_j+2}} \left(\begin{pmatrix} \mathcal{K}_1^{(n_j)} \\ 0 \end{pmatrix} + \mathbf{O}(t^{-1}) \right) \end{aligned}$$

on the complement of the support of U_t and for all real t and $j = 1, \dots, J$. Here $\mathcal{K}_1^{(n_j)}$ and $\mathcal{K}_2^{(n_j)}$ denotes the holomorphic and anti-holomorphic n_j -th derivatives of the functions $\mathcal{K}_1$ and $\mathcal{K}_2$, respectively. Moreover, the transpose of $l_{\frac{2p}{2-p}, t}^{-1}$ is $l_{\frac{2p}{3p-2}, t}$. Therefore we also have

$$\begin{aligned} R^t(0, 0, k, 0) \begin{pmatrix} l_{\frac{2p}{3p-2}, t} & 0 \\ 0 & l_{\frac{2p}{3p-2}, t} \end{pmatrix} \phi_{2k-1} &= \frac{|t|^{(1-2/p)}}{\pi m_j! t^{m_j}} \left(\begin{pmatrix} 0 \\ -\mathcal{K}_1^{(m_j)} \end{pmatrix} + \mathbf{O}(t^{-1}) \right) \text{ and} \\ R^t(0, 0, k, 0) \begin{pmatrix} l_{\frac{2p}{3p-2}, t} & 0 \\ 0 & l_{\frac{2p}{3p-2}, t} \end{pmatrix} \phi_{2k} &= \frac{|t|^{(1-2/p)}}{\pi m_j! t^{m_j}} \left(\begin{pmatrix} \mathcal{K}_2^{(m_j)} \\ 0 \end{pmatrix} + \mathbf{O}(t^{-1}) \right), \end{aligned}$$

on the complement of the support of U_t and for all real t and for all $j = 1, \dots, J$. If we endow E/E_{finite} with the topology of the dual Banach space of the orthogonal complement of E_{finite} in the dual of the Banach space $L^p(B(0, \varepsilon)) \times L^p(B(0, \varepsilon))$, which is isomorphic to $L^{\frac{p}{p-1}}(B(0, \varepsilon)) \times L^{\frac{p}{p-1}}(B(0, \varepsilon))$, then due to Corollary 3.33 the operator A induces an invertible operator from E/E_{finite} onto G_{infinite} . The identity of E minus the projection of E onto the subspace E_{finite} , which was introduced in Lemma 3.37, defines an operator from E/E_{finite} onto the corresponding complementary subspace E_{infinite} . We have seen above that this operator is not bounded in the limit $t \rightarrow \infty$, if $n_J \neq J - 1$. Nevertheless, this divergence is overcompensated by the convergence of the corresponding limits of $R(0, 0, k, 0) \begin{pmatrix} l_{p, t} & 0 \\ 0 & l_{p, t} \end{pmatrix} \psi_j$. Hence the composition of the projection onto E_{infinite} , considered as an operator from E/E_{finite} into E , with the operator $R(0, 0, k, 0) \begin{pmatrix} l_{p, t} & 0 \\ 0 & l_{p, t} \end{pmatrix}$ converges to zero in the limit $t \rightarrow \infty$. On the other hand, the operator $\begin{pmatrix} U_{t=1} & 0 \\ 0 & \bar{U}_{\bar{t}=1} \end{pmatrix}$ maps G_{infinite} onto the dual space of F/F_{finite} , since $\begin{pmatrix} U_{t=1} & 0 \\ 0 & \bar{U}_{\bar{t}=1} \end{pmatrix} R_{\mathbb{R}^2}^t(0, 0, 0)$ acts on F_{finite} as the identity. The analogous projection of F onto the complementary subspace F_{infinite} defines an operator from F/F_{infinite} onto F , which is not

bounded in the limit $t \rightarrow \infty$, if $m_j \neq J - 1$. Analogously this divergence is overcompensated by the convergence of the corresponding limits $R^t(0, 0, k, 0) \begin{pmatrix} I_{\frac{2p}{3p-2}, t} & 0 \\ 0 & I_{\frac{2p}{3p-2}, t} \end{pmatrix} \phi_j$. To sum up, in all cases regardless of whether n_j or m_j is equal to $J - 1$ or not, the arguments of the first two steps combined with the application of Lemma 3.37 yields that only the part

$$G \hookrightarrow G^{**} \twoheadrightarrow (R_{\mathbb{R}^2}^t(0, 0, 0)F)^* \xrightarrow{B_{t, \text{finite}}^{-1}} E_{\text{finite}} \hookrightarrow E$$

of $(A + B_t)^{-1}$ contributes in the limit $t \rightarrow \infty$ to $R(U_t, \bar{U}_t, k, 0)$. Therefore, and with the notation of Section 3.4.1, the operator $\lim_{t \rightarrow \infty} R(U_t, \bar{U}_t, k, 0) - R(0, 0, k, 0)$ has the integral kernel

$$\Psi_D(z) S_D(k) \Phi_D(z') d^2 x' \quad \text{with} \quad S_D(k) = -N (M R_D(k) N)^{-1} M.$$

Obviously this is an integral kernel of a *Finite rank Perturbation* 3.22.

More precisely, we shall show that for all $\varepsilon > 0$, the restrictions (compare with Remark 3.36) of the resolvents $R(U_t, \bar{U}_t, k, 0)$ to the complements of $B(0, \varepsilon)$, considered as operators from $L^p(\Delta \setminus B(0, \varepsilon)) \times L^p(\Delta \setminus B(0, \varepsilon))$ to $L^{\frac{2p}{2-p}}(\Delta \setminus B(0, \varepsilon)) \times L^{\frac{2p}{2-p}}(\Delta \setminus B(0, \varepsilon))$ with $1 < p < 2$, converges in the limit $t \rightarrow \infty$ to the corresponding restriction of the operator above. In fact, due to Lemma 3.20 the restriction of the resolvent $R(0, 0, k, 0)$ considered as an operator from $L^p(B(0, t^{-1}\varepsilon)) \times L^p(B(0, t^{-1}\varepsilon))$ into $L^\infty(\Delta \setminus B(0, \varepsilon)) \times L^\infty(\Delta \setminus B(0, \varepsilon))$ is for all $k \notin \mathcal{F}(0, 0)$ and large t bounded. Therefore, the arguments of the first two steps carry over and prove the claim, provided the Dirac operator $\begin{pmatrix} U_{t=1} & \partial \\ -\bar{\partial} & \bar{U}_{t=1} \end{pmatrix}$ on $\mathbb{P}^1$ has no kernel. Consequently the arguments of the third step imply the claim for general potentials $U_{t=1} \in L^2(B(0, \varepsilon))$. **q.e.d.**

3.4.4 The order of zeroes of spinors

For any potential $U \in L^2(\mathbb{R}^2)$ the restriction U_t of $z \mapsto tU(tz)$ to any small ball $B(0, \varepsilon)$ yields an analogous family of potentials. In order to carry over Lemma 3.35 to these families, we shall investigate the limits $\lim_{t \rightarrow \infty} \begin{pmatrix} I_{p, t} & 0 \\ 0 & I_{p, t} \end{pmatrix} \psi$ of spinors ψ in the kernel of a Dirac operator on $\mathbb{P}^1$ with L^2 -potentials. Obviously these limits depend only on the restriction of ψ to arbitrary small neighbourhoods of $z = \infty$. Moreover, if we use the coordinates z^{-1} instead of z this limit is the same as $\lim_{t \rightarrow 0} \begin{pmatrix} I_{p, t} & 0 \\ 0 & I_{p, t} \end{pmatrix} \psi$ of a spinor ψ in the kernel of $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ on some neighbourhood $0 \ni \mathcal{O} \subset \mathbb{C}$.

In the following discussion concerning these limits we make use of the *Lorentz spaces* $L^{p, q}$. These rearrangement invariant Banach spaces are an extension of the family of the usual Banach spaces L^p indexed by an additional parameter $1 \leq q \leq \infty$ for $1 < p < \infty$. For $p = 1$ or $p = \infty$ we consider only the *Lorentz spaces* $L^{p, \infty}$, which in these cases are isomorphic to L^p ([S-W, Chapter V. §3.], [B-S, Chapter 4 Section 4.] and [Zi, Chapter 1. Section 8.]). We recall some properties of these Banach spaces:

- (i) For $1 < p \leq \infty$ the *Lorentz spaces* $L^{p, p}$ coincide with the usual L^p -spaces. Moreover, the *Lorentz space* $L^{1, \infty}$ coincides with the usual Banach space L^1 .

- (ii) On a finite measure space the *Lorentz space* $L^{p,q}$ is contained in $L^{p',q'}$ either if $p > p'$ or if $p = p'$ and $q \leq q'$.

In [O] Hölder's inequality and Young's inequality are generalized to these *Lorentz spaces* ([B-S, Chapter 4 Section 7.] and [Zi, Chapter 2. Section 10.]):

Generalized Hölder's inequality 3.39. *Either for $1/p_1 + 1/p_2 = 1/p_3 < 1$ and $1/q_1 + 1/q_2 \geq 1/q_3$ or for $1/p_1 + 1/p_2 = 1$, $1/q_1 + 1/q_2 \geq 1$ and $(p_3, q_3) = (1, \infty)$ there exists some constant $C > 0$ with*

$$\|fg\|_{(p_3, q_3)} \leq C \|f\|_{(p_1, q_1)} \|g\|_{(p_2, q_2)}.$$

Generalized Young's inequality 3.40. *Either for $1/p_1 + 1/p_2 - 1 = 1/p_3 > 0$ and $1/q_1 + 1/q_2 \geq 1/q_3$ or for $1/p_1 + 1/p_2 = 1$, $1/q_1 + 1/q_2 \geq 1$ and $(p_3, q_3) = (\infty, \infty)$ there exists some constant $C > 0$ with*

$$\|f * g\|_{(p_3, q_3)} \leq C \|f\|_{(p_1, q_1)} \|g\|_{(p_2, q_2)}.$$

Therefore, the resolvents of the Dirac operators on $\mathbb{R}^2/\Lambda$ or $\mathbb{P}^1$ or $\mathbb{R}^2$ are bounded operators from the L^1 -spinors into the $L^{2,\infty}$ -spinors, from the $L^{2,1}$ -spinors into the continuous spinors, from the L^p -spinors into the $L^{q,p}$ -spinors, and finally from the $L^{p,q}$ -spinors into the L^q -spinors, with $1 < p < 2$ and $q = 2p/(2-p)$. Moreover, the Sobolev constant S_p (compare with Lemma 2.4 and Remark 2.5) is equal to the corresponding norm $\|f\|_{2,\infty}$ times the corresponding constant of the *Generalized Young's inequality* 3.40.

Obviously the restriction of $z \mapsto \psi(tz)$ to any ball $B(0, \varepsilon)$ depends for small t only on the restriction of ψ to an arbitrary small neighbourhood of 0. Let $\mathcal{O} \subset \mathbb{C}$ be a bounded open neighbourhood of 0 and let $V, W \in L^2(\mathcal{O})$ be two potentials on this domain. If ψ is any element in the kernel of the corresponding Dirac operator $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ on this domain, then $(\mathbf{1} - R_{\mathbb{R}^2}(0, 0, 0)) \begin{pmatrix} V & \partial \\ 0 & W \end{pmatrix} \psi$ defines on this domain an element of the kernel of the free Dirac operator $\begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix}$. Vice versa, if we diminish the neighbourhood $\mathcal{O} \ni 0$ and consequently the L^2 -norms of V and W , such that the sum

$$\sum_{n=0}^{\infty} (R_{\mathbb{R}^2}(0, 0, 0)) \begin{pmatrix} V & \partial \\ 0 & W \end{pmatrix}^n = (\mathbf{1} - R_{\mathbb{R}^2}(0, 0, 0)) \begin{pmatrix} V & \partial \\ 0 & W \end{pmatrix}^{-1}$$

converges to a bounded operator on $L^q(\mathcal{O}) \times L^q(\mathcal{O})$ with $2 < q < \infty$, then this operator transforms all elements in the kernel of the free Dirac operator on $\mathcal{O}$ into the kernel of the Dirac operator $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ on this domain. Hence the latter kernel is isomorphic to the former kernel. Moreover, the former kernel contains all functions ψ , whose components are holomorphic and anti-holomorphic functions on $\mathcal{O}$, respectively. This shows that any spinor $\psi \in W^{1,p}(\mathcal{O}) \times W^{1,p}(\mathcal{O})$ with $1 < p < 2$ in the kernel of the Dirac operator $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ on $\mathcal{O}$ with potentials $V, W \in L^2(\mathcal{O})$ are on small neighbourhood equal to $(\mathbf{1} - R_{\mathbb{R}^2}(0, 0, 0)) \begin{pmatrix} V & \partial \\ 0 & W \end{pmatrix}^{-1}$ applied on an element in the kernel of the free Dirac operator on $\mathcal{O}$. Due to Weyl's Lemma [Co-II, §18.4. Lemma 4.10] all elements in the kernel of the free Dirac operator are continuous functions. This implies that the spinor belongs to

$$\psi \in \bigcap_{1 < p < 2} W_{\text{loc}}^{1,p}(\mathcal{O}) \times W_{\text{loc}}^{1,p}(\mathcal{O}) \subset \bigcap_{2 < q < \infty} L_{\text{loc}}^q(\mathcal{O}) \times L_{\text{loc}}^q(\mathcal{O}).$$

Moreover, for $1 < p < 2$ and $\frac{1}{p} = \frac{1}{2} + \frac{1}{q}$ the $L^q(\mathcal{O} \times L^q(\mathcal{O}))$ -norms and the $W^{1,p}(\mathcal{O}) \times W^{1,p}(\mathcal{O})$ -norms of a spinor in the kernel of a Dirac operator with potentials $V, W \in L^2(\mathcal{O})$ are equivalent. Since the kernel of the free Dirac operator is a direct sum of Bergman spaces, i. e. the closed subspaces of $L^q(B(0, \varepsilon))$ of holomorphic and anti-holomorphic functions, respectively ([H-K-Z, Chapter 1.] and [Zh, Chapter 4.]), all kernels of Dirac operators with potentials $V, W \in L^2(\mathcal{O})$ are closed subspaces of $L^q(\mathcal{O}) \times L^q(\mathcal{O})$ ($2 < q < \infty$). In the sequel these kernels are considered as subspaces of these Banach spaces.

If a spinor $\psi \in W^{1,p}(\mathcal{O}) \times W^{1,p}(\mathcal{O})$ with $1 < p < 2$ belongs to the kernel of the Dirac operator $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ on $\mathcal{O}$ with potentials $V, W \in L^2(\mathcal{O})$, then obviously the spinor $\begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix} \psi$ belongs to the kernel of the Dirac operator $\begin{pmatrix} Vz/\bar{z} & \partial \\ -\bar{\partial} & Wz/\bar{z} \end{pmatrix}$.

Order of zeroes 3.41. *The order of a zero of ψ at $z = 0$ is defined as the largest integer m , such that $\begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix}^{-m} \psi \in W^{1,p}(\mathcal{O}) \times W^{1,p}(\mathcal{O})$ with $1 < p < 2$ belongs to the kernel of the Dirac operator $\begin{pmatrix} V(z/\bar{z})^m & \partial \\ -\bar{\partial} & W(\bar{z}/z)^m \end{pmatrix}$. Due to the Strong unique continuation property 2.10 this number is finite and denoted by $\text{ord}_0(\psi)$.*

If the potentials V and W belong to $L^{2,1}(\mathcal{O})$, then the eigen-spinors of the corresponding Dirac operators are continuous spinors. We claim that in this case ψ vanishes at $z = 0$ if and only if $\tilde{\psi} = \begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix}^{-1} \psi$ belongs to the kernel of the Dirac operator $\begin{pmatrix} \tilde{V} & \partial \\ -\bar{\partial} & \tilde{W} \end{pmatrix} = \begin{pmatrix} Vz/\bar{z} & \partial \\ -\bar{\partial} & Wz/\bar{z} \end{pmatrix}$. In fact, since the resolvent $R_{\mathbb{R}^2}(0, 0, 0)$ has the integral kernel

$$\begin{pmatrix} 0 & (z - z')^{-1} \\ (\bar{z}' - \bar{z})^{-1} & 0 \end{pmatrix} \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}},$$

the operator $\begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix} R_{\mathbb{R}^2}(0, 0, 0) \begin{pmatrix} \bar{z} & 0 \\ 0 & z \end{pmatrix}^{-1} - R_{\mathbb{R}^2}(0, 0, 0)$ has the integral kernel $\begin{pmatrix} 0 & 1/z' \\ -1/\bar{z}' & 0 \end{pmatrix} \frac{d\bar{z}' \wedge dz'}{2\pi\sqrt{-1}}$.

Consequently, the operator $\mathbf{1} - R_{\mathbb{R}^2}(0, 0, 0) \begin{pmatrix} \tilde{V} & \partial \\ 0 & \tilde{W} \end{pmatrix}$ maps $\tilde{\psi}$ into the kernel of the free Dirac operator on $\mathcal{O}$, if ψ vanishes at $z = 0$. Vice versa, if $\tilde{\psi}$ belongs to the kernel of the Dirac operator $\begin{pmatrix} \tilde{V} & \partial \\ -\bar{\partial} & \tilde{W} \end{pmatrix}$, then $\tilde{\psi}$ is continuous, since the potentials $\tilde{V}$ and $\tilde{W}$ belong to $L^{2,1}(\mathcal{O})$. Therefore, the spinor $\psi = \begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix} \tilde{\psi}$ vanishes at $z = 0$ and obviously belongs to the kernel of the Dirac operator $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ on $\mathcal{O}$. This proves

Lemma 3.42. *Let V and W be potentials in $L^{2,1}(\mathcal{O})$ on a bounded open neighbourhood of $0 \in \mathcal{O} \subset \mathbb{C}$ and ψ a non-trivial spinor in the kernel of $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$ on $\mathcal{O}$. Then there exists a unique $N_0 \ni m = \text{ord}_0(\psi)$ such that $\begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix}^{-m} \psi$ is a continuous spinor in the kernel of $\begin{pmatrix} V(z/\bar{z})^m & \partial \\ -\bar{\partial} & W(\bar{z}/z)^m \end{pmatrix}$, which does not vanish in a small neighbourhood of $0 \in \mathcal{O}$. In particular, all zeroes are isolated.* **q.e.d.**

If the potentials V and W belong to $L^2(\mathcal{O})$, then the space of all spinors, which have at $0 \in \mathcal{O}$ a zero of positive order, is a subspace of the kernel of the Dirac operator $\begin{pmatrix} V & \partial \\ -\bar{\partial} & W \end{pmatrix}$. Due to our considerations above, this subspace is for small open neighbourhoods $\mathcal{O} \ni 0$ the image

of the kernel of the free Dirac operator on $\mathcal{O}$ under the operator

$$\begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix} \sum_{n=0}^{\infty} \left(R_{\mathbb{R}^2}(0, 0, 0) \begin{pmatrix} Vz/\bar{z} & 0 \\ 0 & W\bar{z}/z \end{pmatrix} \right)^n.$$

The composition of this operator with the operator $\mathbf{1} - R_{\mathbb{R}^2}(0, 0, 0) \begin{pmatrix} V & 0 \\ -\bar{\partial} & W \end{pmatrix}$ is an operator on $L^q(\mathcal{O}) \times L^q(\mathcal{O})$. For small $\mathcal{O}$ this composition belongs to small neighbourhoods of $\begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix}$ in the space of operators on $L^q(\mathcal{O}) \times L^q(\mathcal{O})$. Therefore, for all spinors $\psi \in W^{1,p}(\mathcal{O}) \times W^{1,p}(\mathcal{O})$ in the kernel of $\begin{pmatrix} V & 0 \\ -\bar{\partial} & W \end{pmatrix}$ there exists a unique $\psi' \in W^{1,p}(\mathcal{O}) \times W^{1,p}(\mathcal{O})$ in the same kernel, which has a zero of positive order at 0, such that $(\mathbf{1} - R_{\mathbb{R}^2}(0, 0, 0) \begin{pmatrix} V & 0 \\ -\bar{\partial} & W \end{pmatrix})(\psi + \psi')$ is a constant spinor in the kernel of the free Dirac operator. In particular, the quotient of this kernel modulo the subspace of all spinors, which have at 0 a zero of positive order, is a complex two-dimensional space (for potentials of the form $(U, \bar{U})$ a quaternionic one-dimensional space).

Lemma 3.43. *The quotient spaces of the kernel of the Dirac operator $\begin{pmatrix} V & 0 \\ -\bar{\partial} & W \end{pmatrix}$ with potentials $V, W \in L^2(\mathcal{O})$ on a bounded open neighbourhood of $0 \in \mathcal{O} \subset \mathbb{C}$ divided by the subspace of all spinors, which have at 0 a zero of at least n -th order, is a complex $2n$ -dimensional space (for potentials of the form $(U, \bar{U})$ a quaternionic n -dimensional space). **q.e.d.***

Actually these arguments show that if a spinor in the kernel of $\begin{pmatrix} V & 0 \\ -\bar{\partial} & W \end{pmatrix}$ has no zero at 0, then it has no zero in a neighbourhood of 0. By definition of *Order of zeroes* 3.41 this implies

Corollary 3.44. *The zeroes of any element of the kernel of the Dirac operator $\begin{pmatrix} V & 0 \\ -\bar{\partial} & W \end{pmatrix}$ with potentials $V, W \in L^2(\mathcal{O})$ on a bounded open set $\mathcal{O} \subset \mathbb{C}$ are isolated. **q.e.d.***

From the topological point of view the continuity of a cross section is indispensable in order to determine the topological invariants of the corresponding bundles. But the spinors in the kernels of Dirac operators with potentials $V, W \in L^2$ are in general not continuous. We shall overcome this problem with the help of sheaf theory. In fact, the elements of these kernels of a Dirac operators with potentials $V, W \in L^2$ define sheaves, which are very similar to sheaves of holomorphic functions. In fact, due to the *Strong unique continuation property* 2.10, all local germs have at most one global continuation. Furthermore, due to Lemma 3.43 the corresponding local sheaves have the same properties as the local sheaves of holomorphic functions. Finally, due to Lemma 2.22 the residue can be extended to a pairing of ‘meromorphic’ elements in the kernel of Dirac operator with potentials $V, W \in L^2$. In fact, if locally near $z_l \in \mathcal{O} \subset \mathbb{C}$ ψ and ϕ obey

$$\begin{pmatrix} V(\bar{z} - \bar{z}_l)^{M_l} & \partial(\bar{z} - \bar{z}_l)^{M_l} \\ -\bar{\partial}(z - z_l)^{M_l} & W(z - z_l)^{M_l} \end{pmatrix} \psi = 0 \quad \begin{pmatrix} V(z - z_l)^{M_l} & \bar{\partial}(z - z_l)^{M_l} \\ \partial(\bar{z} - \bar{z}_l)^{M_l} & W(\bar{z} - \bar{z}_l)^{M_l} \end{pmatrix} \phi^t = 0,$$

respectively, then the residue of ψ and ϕ at $z = z_l$ is defined as

$$\text{Res}_{z=z_l} \left(\phi \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \psi \right) = \frac{1}{2\pi\sqrt{-1}} \int_{\partial B(0,\varepsilon)} \phi \begin{pmatrix} 0 & d\bar{z} \\ dz & 0 \end{pmatrix} \psi.$$

Due to Remark 2.7 and the Sobolev embedding theorem [Ad, 5.4 Theorem] the restriction of ψ and ϕ to one-dimensional submanifolds belong to all L^q with $q < \infty$. Therefore, the integral on the right hand side is well defined. Moreover, due to Lemma 2.22 this residue does not depend on ε or more generally on the path around $z = 0$. As a consequence the sheaf-theoretic proofs of the Riemann–Roch theorem [Fo, Theorem 16.9] and Serre Duality [Fo, §17] carry over to these sheaves of elements in the kernel of Dirac operators with potentials $V, W \in L^2$.

In order to establish the relation between Dirac operators with potentials and quaternionic holomorphic structures [F-L-P-P, Definition 2.1] on quaternionic line bundles, we identify the quaternions with all complex 2×2 -matrices of the form $\begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix}$. If we consider a $\mathbb{C}^2$ -valued function $\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$ on an open set $\mathcal{O} \subset \mathbb{C}$ as a quaternionic-valued function $\begin{pmatrix} \psi_1 & -\bar{\psi}_2 \\ \psi_2 & \bar{\psi}_1 \end{pmatrix}$, then $-J$ times the Dirac operator

$$-J \begin{pmatrix} U & \partial \\ -\bar{\partial} & \bar{U} \end{pmatrix} = \begin{pmatrix} \bar{\partial} & -\bar{U} \\ U & \partial \end{pmatrix}$$

yields a quaternionic holomorphic structure on the trivial quaternionic line bundle on $\mathcal{O}$ endowed with the complex structure of multiplication on the left with complex numbers $\mathbb{C} \subset \mathbb{Q}$. In particular, the action of $\sqrt{-1}$ is given by left-multiplication with $\begin{pmatrix} \sqrt{-1} & 0 \\ 0 & -\sqrt{-1} \end{pmatrix}$. In fact, we have

$$\begin{pmatrix} \bar{\partial}\psi_1 & -\bar{\partial}\bar{\psi}_2 \\ \partial\psi_2 & \partial\bar{\psi}_1 \end{pmatrix} + \begin{pmatrix} 0 & -\bar{U} \\ U & 0 \end{pmatrix} \begin{pmatrix} \psi_1 & -\bar{\psi}_2 \\ \psi_2 & \bar{\psi}_1 \end{pmatrix} = \begin{pmatrix} \bar{\partial}\psi_1 - \bar{U}\psi_2 & -\overline{\partial\psi_2 + U\psi_1} \\ \partial\psi_2 + U\psi_1 & \overline{\bar{\partial}\psi_1 - \bar{U}\psi_2} \end{pmatrix}.$$

Hence the corresponding Hopf field is equal to $Q = -\bar{U}d\bar{z}$. Lemma 3.42 now implies, that if U belongs to $L^{2,1}(\mathcal{O})$, then any holomorphic section of the corresponding quaternionic holomorphic line bundle defines a non-vanishing section of another quaternionic holomorphic line bundle, whose degree differs from the degree of the former line bundle by the number of zeroes of the section of the former line bundle. Moreover, the Hopf field $\tilde{Q}$ of the latter line bundle belongs locally also to $L^{2,1}(\mathcal{O})$ and the Willmore energy [F-L-P-P, Definition 2.3] of the latter is the same as the Willmore energy of the former. If on $\mathcal{O}$ ψ is a non-vanishing spinor of the kernel of $\begin{pmatrix} U & \partial \\ -\bar{\partial} & \bar{U} \end{pmatrix}$, then the corresponding quaternionic flat connection may be decomposed into a holomorphic structure and an anti-holomorphic structure [F-L-P-P, (19)]. The holomorphic structure is equal to the holomorphic structure, which is induced by the composition of the Dirac operator with $-J$. Furthermore, the anti-holomorphic structure has the form

$$\begin{pmatrix} \partial\psi_1 & -\partial\bar{\psi}_2 \\ \bar{\partial}\psi_2 & \bar{\partial}\bar{\psi}_1 \end{pmatrix} + \begin{pmatrix} B & A \\ -\bar{A} & \bar{B} \end{pmatrix} \begin{pmatrix} \psi_1 & -\bar{\psi}_2 \\ \psi_2 & \bar{\psi}_1 \end{pmatrix} = 0$$

with $B = -\frac{\bar{\psi}_1\partial\psi_1 + \psi_2\partial\bar{\psi}_2}{\psi_1\bar{\psi}_1 + \psi_2\bar{\psi}_2} A = -\frac{\bar{\psi}_2\partial\psi_1 - \psi_1\partial\bar{\psi}_2}{\psi_1\bar{\psi}_1 + \psi_2\bar{\psi}_2}.$

Due to the decomposition [F-L-P-P, (20)] the operator $\partial + B$ defines an anti-holomorphic structure on the corresponding complex line bundle, and A yields the part of the corresponding anti-holomorphic structure on the quaternionic line bundle, which anti-commutes with

the action of $\sqrt{-1}$ (i. e. the left multiplication with $\sqrt{-1} \in \mathbb{C} \subset \mathbb{Q}$). If the Hopf field belongs locally to $L^{2,1}(\mathcal{O})$, then the eigen-spinors are continuous and belong to the Sobolev space $W^{1,2}(\mathcal{O}) \times W^{1,2}(\mathcal{O})$. Hence for non-vanishing ψ the potentials A and B belong to $L^2(\mathcal{O})$.

Lemma 3.45. *Let $U \in L^2(\mathcal{O})$ be a potential with small L^2 -norm, such that the spinor*

$$\psi = (\mathbf{1} - \mathbf{R}_{\mathbb{R}^2}(0, 0, 0)) \begin{pmatrix} U & 0 \\ 0 & \bar{U} \end{pmatrix}^{-1} \begin{pmatrix} a \\ b \end{pmatrix} \quad \text{with } (a, b) \in \mathbb{P}^1$$

is a well defined non-vanishing spinor on $\mathcal{O}$. Then the parts A and B of the corresponding anti-holomorphic structure, which anti-commute and commute with the action of $\sqrt{-1}$, have finite L^2_{loc} - and $L^{2,\infty}_{\text{loc}}$ -norms, respectively. Finally, the derivative $\bar{\partial}B$ of B defines a real signed measure on $\mathbb{P}^1$, which contains no point measures.

Proof. We just showed that for non-vanishing spinors ψ in the kernel of the Dirac operator $\begin{pmatrix} U & \bar{\partial} \\ -\bar{\partial} & \bar{U} \end{pmatrix}$ on $\mathcal{O}$ with a potential $U \in L^{2,1}(\mathcal{O})$ the potentials A and B of the corresponding anti-holomorphic structure belong to $L^2(\mathcal{O})$. The zero curvature equation of the connection may be written as

$$\begin{pmatrix} \bar{\partial}B & \bar{\partial}A \\ -\bar{\partial}\bar{A} & \bar{\partial}\bar{B} \end{pmatrix} - \begin{pmatrix} B & 0 \\ 0 & \bar{B} \end{pmatrix} \begin{pmatrix} 0 & -\bar{U} \\ U & 0 \end{pmatrix} - \begin{pmatrix} 0 & A \\ -\bar{A} & 0 \end{pmatrix} \begin{pmatrix} B & A \\ -\bar{A} & \bar{B} \end{pmatrix} = \begin{pmatrix} 0 & -\partial\bar{U} \\ \bar{\partial}U & 0 \end{pmatrix} - \begin{pmatrix} 0 & -\bar{U} \\ U & 0 \end{pmatrix}^2.$$

The diagonal and off-diagonal terms of this equation describes the vanishing of the parts of the curvature of the flat connection, which commutes and anti-commutes with the action of $\sqrt{-1}$ [F-L-P-P, proof of Theorem 4.4]:

$$\bar{\partial}B = U\bar{U} - A\bar{A} \quad \bar{\partial}A + \partial\bar{U} = \bar{B}A - B\bar{U}.$$

More invariantly these equations may be written as

$$d(Bdz) = Ud\bar{z} \wedge \overline{Ud\bar{z}} + Adz \wedge \overline{Adz} \quad d(Adz - \bar{U}d\bar{z}) = \overline{Bdz} \wedge Adz + \bar{U}d\bar{z} \wedge Bdz.$$

Hence for $U \in L^{2,1}(\mathcal{O})$ the statement follows.

We shall extend this proof with a limiting argument to small potentials $U \in L^2(\mathcal{O})$. In fact, for any small $U \in L^2(\mathcal{O})$ we choose a sequence of potentials in $L^{2,1}(\mathcal{O})$ with limit U . We extend all potentials to a slightly larger open domain $\mathcal{O}'$ which contains the closure $\bar{\mathcal{O}}$, so that they vanish on the relative complement of $\mathcal{O}$ in $\mathcal{O}'$. Obviously, the corresponding sequence of spinors ψ_n defined in the lemma extend to $\mathcal{O}'$. By definition these spinors are smooth on $\mathcal{O}' \setminus \bar{\mathcal{O}}$. Furthermore, the sequence of integrals of the corresponding one-forms $B_n dz$ along a closed path in $\mathcal{O}' \setminus \bar{\mathcal{O}}$ around $\mathcal{O}$ converges. Since the sequence of measures $U_n \bar{U}_n d^2x$ converges, this implies that the sequence A_n is a bounded sequence in $L^2(\mathcal{O})$. Due to the Banach–Alaoglu theorem [R-S-I, Theorem IV.21], this sequence A_n has a weakly convergent subsequence with limit A . Also the sequence of real signed measures $(U_n \bar{U}_n - A_n \bar{A}_n) d^2x$ on $\mathcal{O}$ has a weakly convergent subsequence. Finally, due to the equations $\bar{\partial}B = U\bar{U} - A\bar{A}$, the sequence of functions B_n is bounded in the Lorentz space $L^{2,\infty}(\mathcal{O})$. Due to [B-S, Chapter 2 Theorem 2.7. and Chapter 4 Corollary 4.8.] this Lorentz space is the dual space of the corresponding Lorentz space $L^{2,1}$. The sequence B_n has also a weakly convergent subsequence with limit B

and $\bar{\partial}B$ considered as a measure is equal to the limit of the measures $(U_n\bar{U}_n - A_n\bar{A}_n) d^2x$. Since the sequence of spinors ψ_n converges in the Banach space of L^q -spinors, and since the sequences A_n and B_n both converge weakly, the limit ψ is anti-holomorphic with respect to the anti-holomorphic structure defined by the limits A and B .

It remains to prove that the function $\bar{\partial}B = -\bar{\partial}\partial \ln(\psi_1\bar{\psi}_1 + \psi_2\bar{\psi}_2)$ considered as a measure contains no point measures. If this finite Baire measure contains a point measure at $z = z'$ of negative mass smaller or equal to $-n\pi$, then, due to the following Lemma 3.46, the spinor $\tilde{\psi} = \begin{pmatrix} z-z' & 0 \\ 0 & \bar{z}-\bar{z}' \end{pmatrix}^{-n} \psi$ belongs to $\bigcap_{q<\infty} L^q_{\text{loc}}(\mathcal{O}) \times L^q_{\text{loc}}(\mathcal{O})$. This implies that ψ has a zero of order n at z' . Hence, due to our assumptions, the masses of all point measures are larger than $-\pi$. Again the following Lemma 3.46 implies that $\psi/(\psi_1\bar{\psi}_1 + \psi_2\bar{\psi}_2)$ is a L^2_{loc} -spinor in the kernel of $\begin{pmatrix} \partial & A \\ -\bar{A} & \bar{\partial} \end{pmatrix}$. Since these kernels are contained in $\bigcap_{q<\infty} L^q_{\text{loc}}(\mathcal{O}) \times L^q_{\text{loc}}(\mathcal{O})$, Lemma 3.46 implies that this measure contains no point measures. **q.e.d.**

Lemma 3.46. *If $\mathcal{O}$ denotes a bounded open subset of $\mathbb{C}$, then for all finite signed Baire measures $d\mu$ on $\mathcal{O}$ [Ro-2, Chapter 13 Section 5] there exists a function h in the Zygmund space $L_{\text{exp}}(\mathcal{O})$ [B-S, Chapter 4 Section 6.] such that $-\bar{\partial}\partial h = d\mu$ (in the sense of distributions). Moreover, if for a suitable $\varepsilon > 0$ all ε -balls of $\mathcal{O}$ have measure smaller than π/q with respect to the positive part $d\mu^+$ of the Hahn decomposition of the finite signed Baire measure $d\mu$ on $\mathcal{O}$ [Ro-2, Chapter 11 Section 5], then the exponentials $\exp(h)$ of all $h \in L_{\text{exp}}(\mathcal{O})$ with $-\bar{\partial}\partial h = d\mu$ belong to $L^q_{\text{loc}}(\mathcal{O})$. Conversely, if the positive part $d\mu^+$ contains a point measure with mass π/q , then the corresponding functions $h \in L_{\text{exp}}(\mathcal{O})$ with $-\bar{\partial}\partial h = d\mu$ does not belong to $L^q_{\text{loc}}(\mathcal{O})$.*

Proof. Due to Dolbeault's Lemma [Gu-Ro, Chapter I Section D 2. Lemma] the convolution with the function $-\frac{2}{\pi} \ln |z|$ defines a right inverse of the operator $-\bar{\partial}\partial$. Now we claim that the restriction of this convolution operator defines a bounded operator from $L^1(\mathcal{O})$ into the Zygmund space $L_{\text{exp}}(\mathcal{O})$. Since the domain $\mathcal{O}$ is bounded, the claim is equivalent to the analogous statement about the restriction to $\mathcal{O}$ of the convolution with the non-negative function

$$f(z) = \begin{cases} -\frac{2}{\pi} \ln |z| & \text{if } |z| < 1 \\ 0 & \text{if } 1 \leq |z| \end{cases}.$$

Associated to this function f is its distribution function μ_f and its non-increasing rearrangement f^* ([S-W, Chapter II §3. Chapter V §3.], [B-S, Chapter 2 Section 1.] and [Zi, Chapter 1. Section 8.]):

$$\mu_f(s) = \pi \exp(-\pi s) \qquad f^*(t) = \begin{cases} -\frac{\ln(t/\pi)}{\pi} & \text{if } 0 \leq t \leq \pi \\ 0 & \text{if } \pi \leq t \end{cases}.$$

If $g \in L^1(\mathcal{O})$, then $g^{**}(t) = \frac{1}{t} \int_0^t g^*(s) ds$ is bounded by $\|g\|_1/t$, since the $L^{1,\infty}$ -norm $\|g\|_{(1,\infty)} = \sup\{tg^{**}(t) \mid t > 0\} = \int_0^\infty g^*(t) dt$ coincides with the L^1 -norm [S-W, Chapter V (3.9)]. Therefore, [Zi, (1.8.14) and (1.8.15)] in the proof of [Zi, 1.8.8. Lemma] (borrowed from [O,

Lemma 1.5.] implies that the non-increasing rearrangement $h^*(t)$ of the convolution $h = f * g$ is bounded by

$$\begin{aligned} h^*(t) \leq h^{**}(t) \leq h_2^{**}(t) + h_1^{**}(t) &\leq g^{**}(t) \int_{f^*(t)}^{\infty} \mu_f(s) ds - \int_t^{\infty} s g^{**}(s) df^*(s) \\ &\leq \frac{\|g\|_1}{t} \exp(-\pi f^*(t)) - \|g\|_1 \int_t^{\infty} df^*(s) \\ &\leq \frac{\|g\|_1}{\pi} + \|g\|_1 f^*(t). \end{aligned}$$

Since by definition the non-increasing rearrangement $h^*(t)$ vanishes for all arguments, which are larger than the Lebesgue measure of $\mathcal{O}$, we conclude the validity of the following estimate:

$$\int_0^{\infty} (\exp(qh^*(t)) - 1) dt \leq \int_0^{|\mathcal{O}|} \exp(qh^*(t)) dt \leq |\mathcal{O}| \exp(q\|g\|_1/\pi) \int_0^{|\mathcal{O}|} (t/\pi)^{-\|g\|_1 q/\pi} dt,$$

with an obvious modification when $\pi < |\mathcal{O}|$. Due to a standard argument [B-S, Chapter 2 Exercise 3.] the finiteness of this integral is equivalent to the statement that $\exp|h|$ belongs to $L^q(\mathcal{O})$. To sum up, the exponential $\exp(h)$ of the convolution $h = f * g$ belongs to $L^q(\mathcal{O})$, if $q < \frac{\pi}{\|g\|_1}$. This proves the claim. In particular, for all $g \in L^1(\mathcal{O})$ there exists an element $h \in L_{\exp}(\mathcal{O})$ with $-\bar{\partial}\partial h = g$.

Due to [B-S, Chapter 4 Theorem 6.5] $L_{\exp}(\mathcal{O})$ is the dual space of the Zygmund space $L \log L(\mathcal{O})$. Hence we shall improve the previous estimate and show that the convolution with $-\frac{2}{\pi} \ln|z|$ defines a bounded operator from $L \log L(\mathcal{O}) \subset L^1(\mathcal{O})$ into $C(\mathcal{O}) \subset L_{\exp}(\mathcal{O})$. By definition of the norm [B-S, Chapter 4 Definition 6.3.]

$$\|g\|_{L \log L} = -\frac{1}{|\mathcal{O}|} \int_0^{|\mathcal{O}|} g^*(t) \ln(t/|\mathcal{O}|) dt = \int_0^{|\mathcal{O}|} g^{**}(t) dt$$

we may improve the previous estimate to [Zi, (1.8.14) and (1.8.15)]

$$h^{**}(t) \leq g^{**}(t) \int_{f^*(t)}^{\infty} \mu_f(s) ds - \int_t^{\infty} s g^{**}(s) df^*(s) \leq \frac{1}{\pi} \int_0^t g^*(s) ds + \int_t^{\pi} g^{**}(s) ds \leq \|g\|_{L \log L}.$$

This implies that in this case $h^{**}(t)$ is bounded, and consequently $h \in L^{\infty}(\mathcal{O})$. Furthermore, since the function $\ln|z|$ is continuous for $z \neq 0$, the convolution with $-\frac{2}{\pi} \ln|z|$ is a bounded operator from $L \log L(\mathcal{O})$ into the Banach space $C(\mathcal{O})$. Finally, the dual of this operator yields a bounded operator from the Banach space of finite signed Baire measures on $\mathcal{O}$ [Ro-2, Chapter 13 Section 5 25. Riesz Representation Theorem] into $L_{\exp}(\mathcal{O})$. More precisely, if

the measure of $\mathcal{O}$ with respect to a finite positive measure $d\mu$ is smaller than π/q , then the exponential $\exp(h)$ of the corresponding function $h = f * d\mu$ belongs to $L^q(\mathcal{O})$.

Due to Weyl's Lemma [Co-II, §18.4. Lemma 4.10] the difference of two arbitrary functions h_1 and h_2 with $-\bar{\partial}\partial h_1 = -\bar{\partial}\partial h_2 = d\mu$ is analytic. Therefore, it suffices to show the second and third statement of the lemma for the convolution of $-\frac{2}{\pi}\ln|z|$ with $d\mu$. Due to the boundedness of $\mathcal{O}$ we may neglect that part of this convolution, where the former function is negative. Therefore, we may neglect the negative part of $d\mu$ in order to bound the exponential $\exp(h)$. The decomposition of the convolution into an ε -near and an ε -distant part analogous to the decomposition in the proof of Lemma 2.6 completes the proof. **q.e.d.**

Proposition 3.47. *Let U be a potential in $L^2(\mathcal{O})$ on a bounded open neighbourhood of $0 \in \mathcal{O} \subset \mathbb{C}$ and ψ a non-trivial spinor in the kernel of the Dirac operator $\begin{pmatrix} U & \bar{\partial} \\ -\bar{\partial} & U \end{pmatrix}$ on $\mathcal{O}$. For any sequence of parameters $t_n \in (0, 1]$ with limit 0 the sequence ψ_{t_n} of the family*

$$\psi_t = \frac{\begin{pmatrix} l_{q,t} & 0 \\ 0 & l_{q,t} \end{pmatrix} \psi}{\left\| \begin{pmatrix} l_{q,t} & 0 \\ 0 & l_{q,t} \end{pmatrix} \psi \right\|_{L^q(B(0,1)) \times L^q(B(0,1))}} \quad (\text{with } 2 < q < \infty)$$

has a subsequence, whose restriction to any $L^q(B(0, \varepsilon)) \times L^q(B(0, \varepsilon))$ with $0 < \varepsilon < \infty$ converges to $\begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix}^{\text{ord}_0(\psi)} \begin{pmatrix} a \\ b \end{pmatrix}$ with $(a, b) \in \mathbb{P}^1$.

Proof. Let us first prove the statement for the spinors considered in Lemma 3.45. These spinors belong locally to the kernels of the operators

$$\begin{pmatrix} \partial & 0 \\ 0 & \bar{\partial} \end{pmatrix} + \begin{pmatrix} B & A \\ -\bar{A} & \bar{B} \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} \bar{\partial} & 0 \\ 0 & \partial \end{pmatrix} + \begin{pmatrix} 0 & -\bar{U} \\ U & 0 \end{pmatrix}.$$

Due to the *Generalized Hölder's inequality* 3.39 the operators $\begin{pmatrix} B & A \\ -\bar{A} & \bar{B} \end{pmatrix}$ and $\begin{pmatrix} 0 & -\bar{U} \\ U & 0 \end{pmatrix}$ define bounded operators from $L_{\text{loc}}^q \times L_{\text{loc}}^q$ into $L_{\text{loc}}^{p,q} \times L_{\text{loc}}^{p,q}$, with $2 < q < \infty$ and $1/p = 1/q + 1/2$. Furthermore, if we restrict these operators to arbitrary small balls $B(0, \varepsilon)$, then we may achieve that the norms of these operators become arbitrarily small. On the other hand, due to the *Generalized Young's inequality* 3.40, the restrictions of the resolvents $-\begin{pmatrix} \partial & 0 \\ 0 & \bar{\partial} \end{pmatrix}^{-1} = \mathbf{J}\mathbf{R}_{\mathbb{R}^2}(0, 0, 0)$ and $-\begin{pmatrix} \bar{\partial} & 0 \\ 0 & \partial \end{pmatrix}^{-1} = \mathbf{R}_{\mathbb{R}^2}(0, 0, 0)\mathbf{J}$ define bounded operators from $L^{p,q}(B(0, \varepsilon)) \times L^{p,q}(B(0, \varepsilon))$ into $L^q(B(0, \varepsilon)) \times L^q(B(0, \varepsilon))$. Therefore, the discussion of spinors in the kernel of Dirac operators previous to of Lemma 3.42 carries over and show that the spinors

$$\left(\mathbf{1} - \mathbf{J}\mathbf{R}_{\mathbb{R}^2}(0, 0, 0) \right) \begin{pmatrix} B & A \\ -\bar{A} & \bar{B} \end{pmatrix} \psi \quad \left(\mathbf{1} - \mathbf{R}_{\mathbb{R}^2}(0, 0, 0)\mathbf{J} \right) \begin{pmatrix} 0 & -\bar{U} \\ U & 0 \end{pmatrix} \psi$$

belong to the kernel of $\begin{pmatrix} \partial & 0 \\ 0 & \bar{\partial} \end{pmatrix}$ and $\begin{pmatrix} \bar{\partial} & 0 \\ 0 & \partial \end{pmatrix}$, respectively. The components of spinors in these kernels belong to the Bergman spaces, i. e. the closed subspaces of $L^q(B(0, \varepsilon))$ of holomorphic and anti-holomorphic functions, respectively ([H-K-Z, Chapter 1.] and [Zh, Chapter 4.]). When ε tends to zero, then these spinors converge to ψ . This implies that the components of ψ converge to the corresponding Bergman projections of ψ ([H-K-Z, Chapter 1.2.] and [Zh, §4.3])

and the analogous projections of ψ onto the anti-holomorphic functions, respectively. These Bergman projections define bounded operators onto the closed subspaces of $L^q(B(0, \varepsilon))$ of holomorphic functions ([H-K-Z, Theorem 1.10] and [Zh, 4.2.3. Theorem]). Since the Bergman projections map the anti-holomorphic functions onto the constant functions (and analogously the projections onto the anti-holomorphic functions maps the holomorphic functions onto the constant functions), ψ converges for $\varepsilon \downarrow 0$ to a constant spinor. The compactness of the space of all constant spinors $\begin{pmatrix} a \\ b \end{pmatrix}$ with $\|\begin{pmatrix} a \\ b \end{pmatrix}\|_{L^q(B(0,1)) \times L^q(B(0,1))}^q = \pi(|a|^q + |b|^q) = 1$ proves the statement for those spinors, which were considered in Lemma 3.45.

If the norms of the restriction of the potentials U , A and B to $B(0, t)$ are very small compared with $\varepsilon^{2/q'-2/q}$ with $0 < \varepsilon < 1$ and $q' < q < \infty$, then these spinors obey the estimate

$$\varepsilon^{2/q'} \|\psi\|_{L^q(B(0,t)) \times L^q(B(0,t))} \leq \|\psi\|_{L^q(B(0,t\varepsilon)) \times L^q(B(0,t\varepsilon))}$$

uniformly with respect to U , A and B and for ε bounded from below. In fact, this follows from the arguments above, since the constant spinors obey this estimate for $q' = q$. Reiterated application implies that these spinors fulfill for all $q' < q$ in the limit $\varepsilon \downarrow 0$ the estimate $\mathbf{O}(\varepsilon^{2/q'}) \leq \|\psi\|_{L^q(B(0,\varepsilon)) \times L^q(B(0,\varepsilon))}$. On the other hand, since all spinors ψ in the kernel of $\begin{pmatrix} U & \partial \\ -\bar{\partial} & \bar{U} \end{pmatrix}$ belong to $\bigcap_{2 < q < \infty} L_{\text{loc}}^q(\mathcal{O}) \times L_{\text{loc}}^q(\mathcal{O})$, Hölder's inequality [R-S-I, Theorem III.1 (c)]

implies for all $q < q''$ the estimate $\|\psi\|_{L^q(B(0,\varepsilon)) \times L^q(B(0,\varepsilon))} \leq \mathbf{O}(\varepsilon^{\text{ord}_0(\psi)+2/q''})$. Now the transformation $\psi \mapsto \tilde{\psi} = \begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix}^{-\text{ord}_0(\psi)} \psi$ together with Lemma 3.43 shows that the statement for general ψ follows from the statement for those ψ considered in Lemma 3.45. **q.e.d.**

Summing up, we have the following equivalent definitions of the *order of a zero* of ψ :

- (i) $\text{ord}_0(\psi)$ is equal to the largest $m \in \mathbb{N}_0$, such that $\begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix}^{-m} \psi$ belongs to the kernel of $\begin{pmatrix} \bar{\partial} & -\bar{U}(\bar{z}/z)^m \\ U(z/\bar{z})^m & \partial \end{pmatrix}$ ($\iff$ Order of zeroes 3.41).
- (ii) For all $2 \leq q' < q < q'' \leq \infty$ we have in the limit $\varepsilon \downarrow 0$

$$\mathbf{O}(\varepsilon^{\text{ord}_0(\psi)+2/q'}) \leq \|\psi\|_{L^q(B(0,\varepsilon)) \times L^q(B(0,\varepsilon))} \leq \mathbf{O}(\varepsilon^{\text{ord}_0(\psi)+2/q''}).$$

- (iii) $\text{ord}_0(\psi)$ is equal to the order of the zero of the limit constructed in Proposition 3.47

We do not know, whether in general these limits of ψ_{t_n} coincide for all sequences t_n converging to zero.

3.4.5 The general case

Now we are prepared to treat the general case. We shall construct a convergent subsequence of a sequence of resolvents $R(U_n, \bar{U}_n, k, 0)$ corresponding to a bounded sequence of complex potentials $U_n \in L^2(\mathbb{R}^2/\Lambda)$. Firstly, due to the Banach-Alaoglu theorem [R-S-I, Theorem IV.21], we may assume that the sequence of potentials is weakly convergent, and that the sequence of non-negative L^1 -functions $U_n \bar{U}_n$ considered as finite Baire measures on $\mathbb{R}^2/\Lambda$ [Ro-2, Chapter 13] is also weakly convergent. If the latter limit does not contain point measures with mass larger or equal to S_p^{-2} then due to Lemma 2.6 the corresponding resolvents converge.

Theorem 3.48. *Any bounded sequence U_n of complex potentials in $L^2(\mathbb{C}/\Lambda_{\mathbb{C}})$ has a subsequence, whose resolvents $R(U_n, \bar{U}_n, k, 0)$ converge weakly to a Finite rank Perturbation 3.22 of the resolvent $R(U, \bar{U}, k, 0)$ of a weak limit of the sequence. More precisely, the support of the Finite rank Perturbation 3.22 is contained in the finite set of singular points $\{z_1, \dots, z_L\}$, which have measure larger or equal to π with respect to a weak limit of the sequence of measures $U_n \bar{U}_n d^2x$. Moreover, for all $\varepsilon > 0$ the restrictions of the subsequence of resolvents to the complements of $\mathcal{S}_{D,\varepsilon} = B(z_1, \varepsilon) \cup \dots \cup B(z_L, \varepsilon)$ converge when considered as operators from $L^p(\Delta \setminus \mathcal{S}_{D,\varepsilon}) \times L^p(\Delta \setminus \mathcal{S}_{D,\varepsilon})$ to $L^{q'}(\Delta \setminus \mathcal{S}_{D,\varepsilon}) \times L^{q'}(\Delta \setminus \mathcal{S}_{D,\varepsilon})$ with $1 < p \leq 2$ and $2 \leq q' < q = \frac{2p}{2-p}$.*

Proof. We prove this theorem in four steps. Since we shall prove the limits of the sequences of resolvents to be *Finite rank Perturbations* 3.22, whose support is contained in the finite set of singular points, we shall decompose the sequence of potentials into a sum of sequences of potentials with disjoint support. More precisely, the limit of one sequence, which we call regular, yields the ordinary weak limit. The limits of the other sequences, which we call singular, and whose domains converge to the singular points, yields the *Finite rank Perturbations* 3.22. Hence in the first step we set up this decomposition. In the second step we treat the cases of single decompositions, in which for each singular point we have one sequence of singular potentials. In the third step we treat the more complicated cases, in which for some of the sequences of singular potentials the procedure of decomposition has to be iterated.

Since we consider only local potentials (i. e. multiplication operators) the corresponding limits are also local in the sense of Remark 3.29. Therefore we treat only the case of one singular point (i. e. $L = 1$). The interested reader should fill in the straight forward modifications of our arguments for the case of several singular points. The crucial issue is the replacement of the Taylor coefficients of the family of resolvents in the beginning of the proof of Lemma 3.35 by the Taylor coefficients of a *Finite rank Perturbation* 3.22, whose support is disjoint from $z = 0$.

1. The decomposition. We shall decompose the sequence of potentials U_n into a sum

$$U_n = U_{\text{reg},n} + \sum_{l=1}^L U_{\text{sing},n,l}$$

of potentials with disjoint support. Here $U_{\text{sing},n,1}, \dots, U_{\text{sing},n,L}$ are the restrictions of U_n to small disjoint balls $B(z_1, \varepsilon_{n,1}), \dots, B(z_L, \varepsilon_{n,L})$, whose radii $\varepsilon_{n,l}$ tend to zero. Consequently, $U_{\text{reg},n}$ are the restrictions of U_n to the complements of the union of these balls. More precisely, we assume

Decomposition (i) For all $l = 1, \dots, L$ the weak limit of the sequence of finite Baire measures $U_{\text{sing},n,l} \bar{U}_{\text{sing},n,l} d^2x$ [Ro-2, Chapter 13] is equal to the point measures of the weak limit of $U_n \bar{U}_n d^2x$ at z_l . Consequently, the weak limit of the measures $U_{\text{reg},n} \bar{U}_{\text{reg},n} d^2x$ is equal to the weak limit of the measures $U_n \bar{U}_n d^2x$ minus the corresponding point measures at $z_1, \dots, z_L$.

Obviously there are many possible choices of the sequences $\varepsilon_{n,l}$ with this property (e. g. for a unique choice of $\varepsilon_{n,l}$ the square of the L^2 -norm of $U_{\text{sing},n,l}$ is equal to the mass of the point

measure at z_l of the weak limit of $U_n \bar{U}_n d^2 x$). Locally we may consider the potentials $U_{\text{sing},n,l}$ as potentials on $\mathbb{P}^1$. We want to treat each of these L sequences of potentials analogously to a subsequence U_{t_n} of an *illustrative family of potentials* 3.4.3 with $t_n \rightarrow \infty$. In general we should use arbitrary Möbius transformations instead of the scaling transformations $z \mapsto tz$. The action of the Möbius group $SL(2, \mathbb{C})/\mathbb{Z}_2$ on $\mathbb{P}^1$

$$SL(2, \mathbb{C}) \ni \begin{pmatrix} a & b \\ c & d \end{pmatrix} : \mathbb{P}^1 \rightarrow \mathbb{P}^1, z \mapsto \frac{az + b}{cz + d}$$

induces a unitary representation π of the Möbius group on $L^2(\mathbb{C})$ [Kn, Chapter II. §4.]

$$\pi \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) : L^2(\mathbb{C}) \rightarrow L^2(\mathbb{C}), \text{ with } \pi \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) U(z) = \frac{1}{|a - cz|^2} U \left(\frac{dz - b}{a - cz} \right).$$

In doing so we transform each of these L sequences of potentials (considered as potentials on $\mathbb{P}^1$) by a sequence of Möbius transformations $g_{n,l}$ in such a way that the transformed sequence of potentials has the following property:

Decomposition (ii) There exists some $\varepsilon > 0$, such that the L^2 -norm of the restrictions of the transformed potentials $\pi(g_{n,l}) U_{\text{sing},n,l}$ to all ε -balls (with respect to the usual metric of $\mathbb{P}^1$) is bounded by the constant $C_p < S_p^{-1}$.

Such decompositions do not always exist. But we shall see now that, if all points of $\mathbb{P}^1$ have measure smaller than $2S_p^{-2}$ with respect to the weak limit of the finite Baire measures $U_n \bar{U}_n d^2 x$, then these decompositions indeed exist. The free Dirac operator on $\mathbb{P}^1$ is invariant under the compact subgroup $SU(2, \mathbb{C})/\mathbb{Z}_2$ of the Möbius group ($\simeq SL(2, \mathbb{C})/\mathbb{Z}_2$) as well as the usual metric on $\mathbb{P}^1$. Therefore, due to the global Iwasawa decomposition [He, Chapter VI Theorem 5.1], it suffices to consider in place of the whole Möbius group only the semidirect product of the scaling transformations ($z \mapsto \exp(t)z$ with $t \in \mathbb{R}$) with the translations ($z \mapsto z + z_0$ with $z_0 \in \mathbb{C}$). In the sequel we assume that all $g_{n,l}$ belong to this subgroup of the Möbius group.

Lemma 3.49. *If the square of the L^2 -norm of a complex potential U on $\mathbb{P}^1$ is smaller than $2S_p^{-2}$, then there exists a constant $C_p < S_p^{-1}$ and a Möbius transformation such that the L^2 -norm of the restrictions of the transformed potentials to all balls of radius $\pi/6$ is not larger than C_p .*

Proof. Let $r_{\max}(U)$ be the maximum of the set

$$\{r \mid \text{the } L^2\text{-norms of the restrictions of } U \text{ to all balls of radius } r \text{ are not larger than } C_p\}.$$

Since the L^2 -norm of the restriction of U to a ball depends continuously on the center and the diameter of the ball, this set has indeed a maximum. Moreover, there exist balls with radius $r_{\max}(U)$, on which the restricted potential has L^2 -norm equal to C_p .

We claim that there exists a Möbius transformation h , such that $r_{\max}(\pi(h)U)$ is the supremum of the set of all $r_{\max}(\pi(g)U)$, where g runs through the Möbius group. Let g_n

be a maximizing sequence of this set, i. e. the limit of the sequence $r_{\max}(\pi(g_n)U)$ is equal to the supremum of the former set. Since $r_{\max}(\pi(g)U)$ is equal to $r_{\max}(U)$, if g belongs to the subgroup $SU(2, \mathbb{C})/\mathbb{Z}_2$ of isometries of the Möbius group, and due to the global Iwasawa decomposition [He, Chapter VI Theorem 5.1], the sequence g_n may be chosen in the semidirect product of the scaling transformations $z \mapsto \exp(t)z$ with the translations $z \mapsto z + z_0$. It is quite easy to see, that if the values of t and z_0 corresponding to a sequence g_n of such Möbius transformations are not bounded, then there exist arbitrary small balls, on which the L^2 -norms of the restrictions of $\pi(g_n)U$ have subsequences converging to the L^2 -norm of U . Hence if the L^2 -norm of U is larger than C_p , then the maximizing sequence of Möbius transformations can be chosen to be bounded and therefore has a convergent subsequence. In this case the continuity implies the claim. If the L^2 -norm of U is not larger than C_p , then $r_{\max}(\pi(g)U)$ does not depend on g and the claim is obvious.

If the L^2 -norm of U is smaller than $\sqrt{2}C_p$, then all $r_{\max}(U)$ -balls, on which the restriction of U has L^2 -norm equal to C_p , have pairwise non-empty intersection. In particular, all of them have non-empty intersection with one of these balls. Consequently, if $r_{\max}(U)$ is smaller than $\pi/6$, then these $r_{\max}(U)$ -balls are contained in one hemisphere. In this case there exists a Möbius transformation g , which enlarges $r_{\max}(U)$ (i. e. $r_{\max}(U) < r_{\max}(\pi(g)U)$). We conclude that there exist a Möbius transformation g , such that $r_{\max}(\pi(g)U)$ is smaller than $\pi/6$. **q.e.d.**

The upper bound $2S_p^{-2}$ is sharp, because for a sequence of L^2 -potentials on $\mathbb{P}^1$, whose square of the absolute values considered as a sequence of finite Baire measures converges weakly to the sum of two point measures of mass S_p^{-2} at opposite points, the corresponding sequence of maxima of $r_{\max}(\pi(h)\cdot)$ converges to zero. But the lower bound $\pi/6$ is of course not optimal.

If the L^2 -norms of the potentials $U_{\text{sing},n,l}$ are smaller than $\sqrt{2}C_p$, then this lemma ensures the existence of Möbius transformations $g_{n,l}$ with the property *Decomposition (ii)*. In general we showed the existence of a sequence of Möbius transformations $h_{n,l}$, which maximizes $r_{\max}(\pi(g)U_{\text{sing},n,l})$. If for one $l = 1, \dots, L$ the corresponding sequences $\pi(h_{n,l})U_{\text{sing},n,l}$ do not obey condition *Decomposition (ii)*, then we apply this procedure of decomposition to the corresponding sequence of potentials $\pi(h_{n,l})U_{\text{sing},n,l}$ on $\mathbb{P}^1$. Consequently, we decompose the sequence $U_{\text{sing},n,l}$ into a finite sum of potentials with disjoint support, such that the corresponding potentials $\pi(g_{n,l})U_{\text{sing},n,l}$ on $\mathbb{P}^1$ obey the analogous conditions *Decomposition (i)*. Due to Lemma 3.49 after finitely many iterations of this procedure of decomposing the potentials into finite sums of potentials with disjoint support, we arrive at a decomposition

$$U_n = U_{\text{reg},n} + \sum_{l=1}^{L'} U_{\text{sing},n,l}$$

of potentials with disjoint support. More precisely, the potentials $U_{\text{sing},n,1}, \dots, U_{\text{sing},n,L'}$ are restrictions of U_n either to small balls or to the relative complements of finitely many small balls inside of small balls. In particular, the domains of these potentials are excluded either from the domain of $U_{\text{reg},n}$, or from the domain of another $U_{\text{sing},n,l}$. The former potentials obey condition *Decomposition (i)* and the latter obey condition

Decomposition (i') If the domain of $U_{\text{sing},n,l'}$ is excluded from the domain of $U_{\text{sing},n,l}$, then the weak limit of the sequence of finite Baire measures $\pi(g_{n,l}) U_{\text{sing},n,l'} \pi(g_{n,l}) \bar{U}_{\text{sing},n,l'} d^2x$ on $\mathbb{C} \subset \mathbb{P}^1$ converges weakly to the point measure of the weak limit of the sequence $\pi(g_{n,l}) U_{\text{sing},n,l} \pi(g_{n,l}) \bar{U}_{\text{sing},n,l} d^2x$ at some point of $\mathbb{C}$, whose measure with respect to the latter limit is not smaller than S_p^{-2} .

All these potentials $U_{\text{sing},n,1}, \dots, U_{\text{sing},n,L'}$ obey condition *Decomposition* (ii). We remark that if the weak limit of the finite Baire measures $\pi(h_{n,l}) U_{\text{sing},n,l} \pi(h_{n,l}) \bar{U}_{\text{sing},n,l} d^2x$ on $\mathbb{C} \subset \mathbb{P}^1$, where $h_{n,l}$ denotes the sequence of Möbius transformations minimizing $R(\pi(g) U_{\text{sing},n,l})$, contains at $z = \infty$ a point measure, whose mass is not smaller than S_p^{-2} , then we decompose from the sequence $\pi(h_{n,l}) U_{\text{sing},n,l}$ potentials, whose domains are the complement of a large ball in the domains of these potentials. In these cases the domains of the analog to the regular sequence of the decomposition are excluded from the domains of the analog to the singular sequence, whose L^2 -norm accumulates at $z = \infty$. Since the Möbius transformations corresponding to the former are faster divergent than the Möbius transformations of the latter, the latter should be considered as less singular than the former. Therefore, also in this case the domains of the more singular sequences are excluded from the domains of the less singular sequences.

By construction and due to Lemma 2.6 the sequence of resolvents $R(U_{\text{reg},n}, \bar{U}_{\text{reg},n}, k, 0)$ converges in the norm limit to the resolvent $R(U, \bar{U}, k, 0)$ of the weak limit U of the sequence of potentials. We shall include the perturbations of the sequences $U_{\text{sing},n,1}, \dots, U_{\text{sing},n,L'}$. In doing so we start with the most singular of these sequences (i. e. those $U_{\text{sing},n,l}$, whose domains are small balls) and include subsequently those sequences, from whose domains the domains of the former sequences are excluded. In the last step we include the sequences $U_{\text{reg},n}$.

2. The case of single decompositions. Here we assume that the decomposition obeys conditions *Decomposition* (i)–(ii) (i. e. the domains of all $U_{\text{sing},n,l}$ are small balls) and $L = 1$. Due to Corollary 3.33 and condition *Decomposition* (ii) a subsequence of the sequence of resolvents $R_{\mathbb{P}^1}(\pi(g_{n,1}) U_{\text{sing},n,1}, \pi(g_{n,1}) \bar{U}_{\text{sing},n,1}, 0)$ converges to the resolvent $R_{\mathbb{P}^1}(U_{\text{sing},1}, \bar{U}_{\text{sing},1}, 0)$ of the weak limit $U_{\text{sing},1}$ of $\pi(g_{n,1}) U_{\text{sing},n,1}$. Therefore, the arguments in Lemma 3.35 concerning the case that the Dirac operator $(1 + z\bar{z}) \begin{pmatrix} U_{t=1} & \partial \\ -\bar{\partial} & \bar{U}_{t=1} \end{pmatrix}$ on $\mathbb{P}^1$ has a trivial kernel, carry over to the case that the Dirac operator $(1 + z\bar{z}) \begin{pmatrix} U_{\text{sing},1} & \partial \\ -\bar{\partial} & \bar{U}_{\text{sing},1} \end{pmatrix}$ on $\mathbb{P}^1$ has a trivial kernel.

In case that the latter Dirac operator has a non-trivial kernel, we follow again the lines of the proof of Lemma 3.35 concerning the analogous case. In the notation of the proof of Lemma 3.35 we use the formula (compare with Remark 3.34)

$$R(U_{\text{sing},n,1}, \bar{U}_{\text{sing},n,1}, k, 0) - R(0, 0, k, 0) = R(0, 0, k, 0) \begin{pmatrix} l_{\frac{4}{3},n} & 0 \\ 0 & l_{\frac{4}{3},n} \end{pmatrix} (A_n + B_n)^{-1} \begin{pmatrix} \pi(g_{n,1}) U_{\text{sing},n,1} & 0 \\ 0 & \pi(g_{n,1}) \bar{U}_{\text{sing},n,1} \end{pmatrix} \begin{pmatrix} l_{4,n} & 0 \\ 0 & l_{4,n} \end{pmatrix}^{-1} R(0, 0, k, 0).$$

Here $l_{p,n}$ is the analog sequence of isometries induced by the sequence of Möbius transformations $g_{n,1}$. Moreover, $E_{\text{finite}} \subset L^{\frac{4}{3}}(\mathbb{R}^2) \times L^{\frac{4}{3}}(\mathbb{R}^2)$ is the kernel of the operator $\mathbb{1} - \begin{pmatrix} U_{\text{sing},1} & 0 \\ 0 & \bar{U}_{\text{sing},1} \end{pmatrix} R_{\mathbb{R}^2}(0, 0, 0)$ and $F_{\text{finite}} \subset L^{\frac{4}{3}}(\mathbb{R}^2) \times L^{\frac{4}{3}}(\mathbb{R}^2)$ is the co-kernel of the operator

$\mathbb{1} - R_{\mathbb{R}^2}(0, 0, 0) \begin{pmatrix} U_{\text{sing},1} & 0 \\ 0 & \bar{U}_{\text{sing},1} \end{pmatrix}$ (compare with Remark 3.34). Furthermore, the two sequences of operators A_n and B_n are the corresponding decompositions introduced in Lemma 3.37 of the restriction of the sequence of operators

$$\begin{pmatrix} I_{\frac{4}{3},n} & 0 \\ 0 & I_{\frac{4}{3},n} \end{pmatrix}^{-1} \left(\mathbb{1} - \begin{pmatrix} U_{\text{sing},n,1} & 0 \\ 0 & \bar{U}_{\text{sing},n,1} \end{pmatrix} R(0, 0, k, 0) \right) \begin{pmatrix} I_{\frac{4}{3},n} & 0 \\ 0 & I_{\frac{4}{3},n} \end{pmatrix}$$

to the domains of $\pi(g_{n,1})U_{\text{sing},n,1}$. More precisely, let $B(z_1, \varepsilon_{n,1})$ denote the sequence of domains of $U_{\text{sing},n,1}$. Then these restrictions to the Möbius transformed domains $g_{n,1}B(z_1, \varepsilon_{n,1}) \subset \mathbb{R}^2$ are considered as operators on $L^{\frac{4}{3}}(\mathbb{R}^2) \times L^{\frac{4}{3}}(\mathbb{R}^2)$. Also the subspaces, which are denoted in Lemma 3.37 by capital F, are here again denoted by $R_{\mathbb{R}^2}^t(0, 0, 0)F$ (compare with Remark 3.38). By definition of E_{finite} and F_{finite} the spaces $R_{\mathbb{R}^2}(0, 0, 0)E_{\text{finite}}$ and $R_{\mathbb{R}^2}^t(0, 0, 0)F_{\text{finite}}$ are the kernel and co-kernel of the Diarc operator $\begin{pmatrix} U_{\text{sing},1} & \partial \\ -\bar{\partial} & \bar{U}_{\text{sing},1} \end{pmatrix}$, respectively. So the numbers $n_1, \dots, n_J$ and $m_1, \dots, m_J$ here are defined as those numbers, which occur as *Order of zeroes* 3.41 of the elements of the latter two spaces, respectively. In the situation of Lemma 3.35 this definition gives the same numbers as the definition therein.

With the help of Lemma 3.21 we may now again determine the limits of the resolvents $R(U_{\text{sing},n,l}, \bar{U}_{\text{sing},n,l}, k, 0)$. Here the family B_t in the proof of Lemma 3.35 is replaced by the sequence of finite rank operators B_n . We remark that the Möbius transformations $g_{n,1}$ belong to the semidirect product of the scaling transformations ($z \mapsto \exp(t)z$ with $t \in \mathbb{R}$) with the translations ($z \mapsto z + z_0$ with $z_0 \in \mathbb{C}$). Since both resolvents $R(0, 0, k, \lambda)$ and $R_{\mathbb{R}^2}(0, 0, \lambda)$ are invariant under the translations, it suffices to take the dilatations into account, and that was done in Lemma 3.35. However, in addition to the arguments in the proof of that Lemma we encounter two complications. Firstly, the domains $g_{n,1}B(z_1, \varepsilon_{n,1})$ depend on n . For all elements $\psi \in E_{\text{finite}}$ and elements $\phi \in F_{\text{finite}}$ let

$$\begin{aligned} \psi_{n,\text{ren}} &= \frac{\psi|_{g_{n,1}B(z_1, \varepsilon_{n,1})}}{\|R_{\mathbb{R}^2}(0, 0, 0)\psi\|_{L^4(\mathbb{R}^2 \setminus g_{n,1}B(z_1, \varepsilon_{n,1})) \times L^4(\mathbb{R}^2 \setminus g_{n,1}B(z_1, \varepsilon_{n,1}))}} \\ \phi_{n,\text{ren}} &= \frac{\phi|_{g_{n,1}B(z_1, \varepsilon_{n,1})}}{\|R_{\mathbb{R}^2}^t(0, 0, 0)\phi\|_{L^4(\mathbb{R}^2 \setminus g_{n,1}B(z_1, \varepsilon_{n,1})) \times L^4(\mathbb{R}^2 \setminus g_{n,1}B(z_1, \varepsilon_{n,1}))}}, \end{aligned}$$

denote two sequences of restricted and renormalized spinors. In order to carry over the proof of Lemma 3.35 to the present situation we replace the pairings of the elements $\psi \in E_{\text{finite}}$ with the functions $\begin{pmatrix} \bar{z}^m \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ z^m \end{pmatrix}$ and of the elements $\phi \in F_{\text{finite}}$ with the functions $\begin{pmatrix} z^m \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ \bar{z}^m \end{pmatrix}$. So we use the renormalized pairings

$$\lim_{n \rightarrow \infty} \langle \langle \psi_{n,\text{ren}}, \begin{pmatrix} \bar{z}^m \\ 0 \end{pmatrix} \rangle \rangle, \lim_{n \rightarrow \infty} \langle \langle \psi_{n,\text{ren}}, \begin{pmatrix} 0 \\ z^m \end{pmatrix} \rangle \rangle \quad \text{and} \quad \lim_{n \rightarrow \infty} \langle \langle \phi_{n,\text{ren}}, \begin{pmatrix} z^m \\ 0 \end{pmatrix} \rangle \rangle, \lim_{n \rightarrow \infty} \langle \langle \phi_{n,\text{ren}}, \begin{pmatrix} 0 \\ \bar{z}^m \end{pmatrix} \rangle \rangle.$$

We remark that due to Proposition 3.47 the limits

$$\lim_{n \rightarrow \infty} R_{\mathbb{R}^2}(0, 0, 0) \begin{pmatrix} I_{\frac{4}{3},n} & 0 \\ 0 & I_{\frac{4}{3},n} \end{pmatrix} \psi_{n,\text{ren}} \quad \text{and} \quad \lim_{n \rightarrow \infty} R_{\mathbb{R}^2}^t(0, 0, 0) \begin{pmatrix} I_{\frac{4}{3},n} & 0 \\ 0 & I_{\frac{4}{3},n} \end{pmatrix} \phi_{n,\text{ren}}$$

exist and are non-trivial for non-trivial ψ and ϕ .

Secondly, we have to take the differences of $\pi(g_{n,l}) U_{\text{sing},n,l}$ from $U_{\text{sing},l}$ into account. Therefore in the present situation the sequence $\mathbf{B}_n$ differs from the corresponding sequence in Lemma 3.35 by a sequence of finite rank operators. The corresponding data $(E, F, \bar{Q})$ belong to a finite-dimensional Grassmannian (compare with the seventh step of Section 3.4.1). These Grassmannians are compact [Gr-Ha, Chapter 1 §5.]. Consequently those subsequences of resolvents, which correspond to convergent sequences, converge to *rank one Perturbations* 3.22.

Now we shall include into our considerations the sequence $U_{\text{reg},n}$. For this purpose we use the tools of the third step of Section 3.4.1. More precisely, we replace in the corresponding formulas the operators $\mathbf{S}(k)$ by the sequence of operators

$$\mathbf{S}_n(k) = \begin{pmatrix} \mathbf{l}_{\frac{4}{3},n} & 0 \\ 0 & \mathbf{l}_{\frac{4}{3},n} \end{pmatrix} (\mathbf{A}_n + \mathbf{B}_n)^{-1} \begin{pmatrix} \pi(g_{n,1}) U_{\text{sing},n,1} & 0 \\ 0 & \pi(g_{n,1}) \bar{U}_{\text{sing},n,1} \end{pmatrix} \begin{pmatrix} \mathbf{l}_{4,n} & 0 \\ 0 & \mathbf{l}_{4,n} \end{pmatrix}^{-1}.$$

Then the corresponding perturbations of $\mathbf{R}(U_{\text{reg},n}, \bar{U}_{\text{reg},n}, k, 0)$ are equal to

$$\mathbf{R}(U_{\text{reg},n}, \bar{U}_{\text{reg},n}, k, 0) \mathbf{S}_n(U_{\text{reg},n}, \bar{U}_{\text{reg},n}, k) \mathbf{R}(U_{\text{reg},n}, \bar{U}_{\text{reg},n}, k, 0), \text{ with} \\ \mathbf{S}_n(U_{\text{reg},n}, \bar{U}_{\text{reg},n}, k) = (\mathbf{1} + \mathbf{S}_n(k) \mathbf{R}(0, 0, k, 0) - \mathbf{S}_n(k) \mathbf{R}(U_{\text{reg},n}, \bar{U}_{\text{reg},n}, k, 0))^{-1} \mathbf{S}_n(k).$$

Moreover, in the present situation for all $\psi \in E_{\text{finite}}$ and $\phi \in F_{\text{finite}}$ we redefine the sequences

$$\psi_{n,\text{ren}} = \frac{\psi|_{g_{n,1}B(z_1, \varepsilon_{n,1})}}{\left\| \mathbf{R}_{\mathbb{R}^2}(U_{\text{reg},n}|_{\mathcal{O}}, \bar{U}_{\text{reg},n}|_{\mathcal{O}}, 0) \begin{pmatrix} \mathbf{l}_{\frac{4}{3},n} & 0 \\ 0 & \mathbf{l}_{\frac{4}{3},n} \end{pmatrix} \psi \right\|_{L^4(\mathcal{O} \setminus B(z_1, \varepsilon)) \times L^4(\mathcal{O} \setminus B(z_1, \varepsilon))}} \\ \phi_{n,\text{ren}} = \frac{\phi|_{g_{n,1}B(z_1, \varepsilon_{n,1})}}{\left\| \mathbf{R}_{\mathbb{R}^2}^t(U_{\text{reg},n}|_{\mathcal{O}}, \bar{U}_{\text{reg},n}|_{\mathcal{O}}, 0) \begin{pmatrix} \mathbf{l}_{\frac{4}{3},n} & 0 \\ 0 & \mathbf{l}_{\frac{4}{3},n} \end{pmatrix} \psi \right\|_{L^4(\mathcal{O} \setminus B(z_1, \varepsilon)) \times L^4(\mathcal{O} \setminus B(z_1, \varepsilon))}}.$$

Here $\mathcal{O}$ is a small open neighbourhood (in order to ensure the existence of the resolvent), which contains a ball $B(z_1, \varepsilon)$, and $U_{\text{reg},n}|_{\mathcal{O}}$ are the restrictions of the sequence of potentials $U_{\text{reg},n}$ to $\mathcal{O}$. We claim that this normalization essentially does not depend on $\mathcal{O}$ and ε . More precisely, for all non-trivial ψ and ϕ and $k \notin \mathcal{F}(U, \bar{U})$ and all appropriate choices of $\mathcal{O}$ and ε the limits

$$\psi(k) = \lim_{n \rightarrow \infty} \mathbf{R}(U_{\text{reg},n}, \bar{U}_{\text{reg},n}, k, 0) \begin{pmatrix} \mathbf{l}_{\frac{4}{3},n} & 0 \\ 0 & \mathbf{l}_{\frac{4}{3},n} \end{pmatrix} \psi_{n,\text{ren}} \\ \phi(k) = \lim_{n \rightarrow \infty} \mathbf{R}^t(U_{\text{reg},n}, \bar{U}_{\text{reg},n}, k, 0) \begin{pmatrix} \mathbf{l}_{\frac{4}{3},n} & 0 \\ 0 & \mathbf{l}_{\frac{4}{3},n} \end{pmatrix} \phi_{n,\text{ren}}$$

converge on Δ to non-trivial functions $\psi(k)$ and $\phi(k)$, which belong on $\Delta \setminus \{z_1\}$ to the kernel and co-kernel of $\mathbf{D}(U, \bar{U}, k)$, respectively, and fulfill on a neighbourhood of z_1 the equations

$$\begin{pmatrix} U(\bar{z} - \bar{z}_1)^{M_1} & \partial(\bar{z} - \bar{z}_1)^{M_1} \\ -\bar{\partial}(z - z_1)^{M_1} & \bar{U}(z - z_1)^{M_1} \end{pmatrix} \psi(k) = 0 \quad \text{and} \quad \begin{pmatrix} U(z - z_1)^{M_1} & \bar{\partial}(z - z_1)^{M_1} \\ \partial(\bar{z} - \bar{z}_1)^{M_1} & \bar{U}(\bar{z} - \bar{z}_1)^{M_1} \end{pmatrix} \phi(k) = 0.$$

In fact, the sequence of resolvents $R(U_{\text{reg},n}, \bar{U}_{\text{reg},n}, k, 0)$ converges as compact operators from $L^{\frac{4}{3}}(\Delta) \times L^{\frac{4}{3}}(\Delta)$ into $L^q(\Delta) \times L^q(\Delta)$ with $q < 4$. If we apply Proposition 3.47 to the integral kernels of these operators with fixed z and z' in the complement of $B(z_1, \varepsilon)$, then the claim follows from the considerations concerning the analogous limits with trivial regular potentials $U_{\text{reg},n}$. These limits yield the entries of the row-vector $\Psi_D(z, k)$ and column-vector $\Phi_D(z, k)$ (compare with *Finite rank Perturbations* 3.22).

Now we show that in the present situation again the sequence of operators A_n does not contribute to the limit of $R(U_n, \bar{U}_n, k, 0)$. If the numbers n_J and m_J are equal to J , then the sequence of projections onto E_{finite} and F_{finite} are again bounded. Due to Lemma 2.6 we may replace the potentials $U_{\text{reg},n}$ by the restrictions $U'_{\text{reg},n}$ of U_n to the complement of $B(z_1, \varepsilon'_{n,1})$ with a sequence of larger radii $0 < \varepsilon_{n,1} < \varepsilon'_{n,1}$ converging to zero, such that $\varepsilon_{n,1}/\varepsilon'_{n,1}$ converges to zero. By inserting in the formula for the perturbations of $R(U'_{\text{reg},n}, \bar{U}'_{\text{reg},n}, k, 0)$ the equations

$$\begin{aligned} R(U'_{\text{reg},n}, \bar{U}'_{\text{reg},n}, k, 0) &= \left(\mathbf{1} - R(0, 0, k, 0) \begin{pmatrix} U'_{\text{reg},n} & 0 \\ 0 & \bar{U}'_{\text{reg},n} \end{pmatrix} \right)^{-1} R(0, 0, k, 0) \\ &= R(0, 0, k, 0) \left(\mathbf{1} - \begin{pmatrix} U'_{\text{reg},n} & 0 \\ 0 & \bar{U}'_{\text{reg},n} \end{pmatrix} R(0, 0, k, 0) \right)^{-1} \end{aligned}$$

we may achieve that the free resolvent $R(0, 0, k, 0)$ occurs at all places on the left hand sides and on the right hand sides of the operators $S_n(k)$. Hence it suffices to estimate the operator $R(0, 0, k, 0)$ as an operator from $L^{\frac{4}{3}}(\Delta \setminus B(0, \varepsilon')) \times L^{\frac{4}{3}}(\Delta \setminus B(0, \varepsilon'))$ into $L^4(B(0, \varepsilon)) \times L^4(B(0, \varepsilon))$. The fourth power of the norm of this operator is smaller than the integral

$$\begin{aligned} \int_{B(0, \varepsilon)} \int_{\Delta \setminus B(0, \varepsilon')} \frac{1}{|z - z'|^4} \frac{d^2 x' d^2 x}{\pi} &\leq \int_{B(0, \varepsilon)} \int_{\mathbb{R}^2 \setminus B(0, \varepsilon')} \frac{1}{|z - z'|^4} \frac{d^2 x' d^2 x}{\pi} \\ &\leq \int_0^\varepsilon \int_{\varepsilon'}^\infty \int_0^\pi \frac{4d\theta R dR r dr}{(R^2 + r^2 + 2Rr \cos(\theta))^2} \\ &\leq \int_0^\varepsilon \int_{\varepsilon'/r}^\infty \int_0^\pi \frac{4d\theta (R/r) d(R/r)}{(1 + (R/r)^2 + 2(R/r) \cos(\theta))^2} \frac{dr}{r} \\ &\leq 2\pi \int_0^\varepsilon \left(\frac{1}{((\varepsilon'/r)^2 - 1)^2} + \frac{1}{(\varepsilon'/r)^2 - 1} \right) \frac{dr}{r} \\ &\leq \pi \int_{(\varepsilon'/\varepsilon)^2}^\infty \frac{(1+x-1)dx}{x(x-1)^2} \leq \frac{\pi}{(\varepsilon'/\varepsilon)^2 - 1}. \end{aligned}$$

Here we used the equalities

$$\begin{aligned} \int_{b_0}^{\infty} \int_{[0,\pi]} \frac{4d\theta b db}{(1+b^2-2b\cos(\theta))^2} &= \int_{b_0}^{\infty} \int_{[0,\pi]} \frac{d\theta b db}{b^2 \left(\frac{1+b^2}{2b} - \cos(\theta)\right)^2} = \int_{b_0}^{\infty} \frac{\pi \frac{1+b^2}{2b} b db}{b^2 \left(\left(\frac{1+b^2}{2b}\right)^2 - 1\right)^{3/2}} \\ &= 2\pi \int_{b_0}^{\infty} \frac{(1+b^2)2bdb}{(b^2-1)^3} = 2\pi \int_{b_0}^{\infty} \left(\frac{2}{(b^2-1)^3} + \frac{1}{(b^2-1)^2} \right) db^2 = 2\pi \left(\frac{1}{(b_0^2-1)^2} + \frac{1}{b_0^2-1} \right). \end{aligned}$$

In fact, due to [Ah, Chapter 4 Section 5.3] the integral $\int_0^\pi \frac{d\theta}{a-\cos(\theta)}$ is for $a > 1$ equal to $\pi(a^2-1)^{-1/2}$. Furthermore, the derivatives with respect to a yields with $a = \frac{1+b^2}{2b}$ and $b > 1$ this formula. Since $\varepsilon_{n,1}/\varepsilon'_{n,1}$ converges to zero, the operators A_n do not contribute to the limit. Moreover, due to the classification of *Finite rank Perturbations* 3.22 in the sixth step of Section 3.4.1 the space of *rank one Perturbations* 3.22 of $R(U, \bar{U}, k, 0)$ is also compact. We remark that due to the considerations of the ninth step in Section 3.4.1, the limits obey the equation in the definition of *rank one Perturbations* 3.22. Hence for $n_J = n_J = J$ the resolvents $R(U_n, \bar{U}_n, k, 0)$ have subsequences, which converge to *Finite rank Perturbations* 3.22 of $R(U, \bar{U}, k, 0)$. The arguments in Lemma 3.35 concerning the general case of values of the sequences $n_1, \dots, n_J$ and $m_1, \dots, m_J$ carry over, and we have proven that all sequences, which allow a decomposition obeying conditions *Decomposition* (i)–(ii) have subsequences, whose corresponding resolvents converge to *Finite rank Perturbations* 3.22 of $R(U, \bar{U}, k, 0)$.

3. The case of iterated decompositions. Finally, we extend our arguments to the case that the decomposition obeys condition *Decomposition* (ii'). In this case for all singular points $z_l \in \{z_1, \dots, z_L\}$ the limits of the sequence of resolvents corresponding to the most singular sequences of potentials at z_1 may contribute to analogous *Finite rank Perturbations* of the resolvents of Dirac operators on $\mathbb{P}^1$. More generally, the *Finite rank Perturbations* of the sequence of resolvents corresponding to those sequences of potentials, whose domains are excluded of one particular sequence of potentials, may contribute to analogous *Finite rank Perturbations* of the sequence of resolvents corresponding to the particular sequence of potentials. More precisely, an easy calculation shows that the resolvents of *Finite rank Perturbations* of the usual Dirac operators do not exist for different values of λ . In case of resolvents on $\mathbb{R}^2/\Lambda$ we overcame this problem by using the family of resolvents $R(U, \bar{U}, k, 0)$. As we already mentioned, the corresponding operators $\tilde{R}(U, \bar{U}, k, 0)$ may indeed be considered as a family of resolvents, if we multiply the corresponding operators with off-diagonal matrices (e. g. with $-J$). On $\mathbb{P}^1$ we may use the same trick and multiply the usual Dirac operator on the left with the matrix $-J$. The corresponding resolvents of the transformed operators are calculated by the usual method of perturbing the operator $(1+z\bar{z}) \begin{pmatrix} U & \partial \\ -\bar{\partial} & \bar{U} \end{pmatrix}$ by the potential $(1+z\bar{z}) \begin{pmatrix} 0 & \lambda/(1+z\bar{z}) \\ -\lambda/(1+z\bar{z}) & 0 \end{pmatrix}$. Since $\lambda/(1+z\bar{z})$ belongs to $L^2(\mathbb{C})$, the methods of Section 2.1 can be used in order to develop the spectral theory of this perturbed Dirac operators. Therefore, the product of the inverse of these operators multiplied on the right with J yields the resolvents of the Dirac operators on $\mathbb{P}^1$ multiplied on the left with $-J$. With this trick the kernel of the *Finite rank Perturbations* of Dirac operators on $\mathbb{P}^1$ analogous to the

Finite rank Perturbations 3.22 are well defined. The whole discussion of the local behaviour of non-trivial elements in the kernel of Dirac operators on $\mathbb{P}^1$ in Section 3.4.4 extends obviously to the elements in the kernels of *Finite rank Perturbations* of Dirac operators on $\mathbb{P}^1$. Whenever we consider *Finite rank Perturbations* of Dirac operators on $\mathbb{P}^1$ in the following paragraphs, then these operators should be considered as implicitly defined by the corresponding *Finite rank Perturbations* of these resolvents of Dirac-like operators.

Remark 3.50. *The generalization of Theorem 3.48 as suggested in Remark 3.30 has an obvious extension to these Finite rank Perturbations of Dirac operators on $\mathbb{P}^1$.*

Let us consider the situation, where at one singular point (i. e. $L = 1$) two sequences (i. e. $L' = 2$) of potentials $U_{\text{sing},n,1}$ and $U_{\text{sing},n,2}$ accumulate in such a way, that the transformed potentials $\pi(g_{n,1}) U_{\text{sing},n,2}$ accumulates at one point of $\mathbb{C} \subset \mathbb{P}^1$. Here $g_{n,1}$ and $g_{n,2}$ denotes the corresponding sequences of Möbius transformations, such that conditions *Decomposition* (i)–(ii) and (i')–(ii) are fulfilled, respectively. In a first step we determine the limit of the transformed of the resolvents $R(U_{\text{sing},n,1} + U_{\text{sing},n,2}, \bar{U}_{\text{sing},n,1} + \bar{U}_{\text{sing},n,2}, k, 0)$ under the Möbius transformations $g_{n,1}$. In doing so we use a variant of the line of arguments concerning the case of a decomposition of potentials obeying condition *Decomposition* (i)–(ii) with one singular point (i. e. $L = L' = 1$). In fact, firstly we determine the sequence of resolvents $R(U_{\text{sing},n,2}, \bar{U}_{\text{sing},n,2}, k, 0)$ transformed under the Möbius transformations $g_{n,1}$. Since the sequence of resolvents on corresponding to the Dirac operators on $\mathbb{P}^1$ with potentials $(1 + z\bar{z})\pi(g_{n,2}) U_{\text{sing},n,2}$ converge, in the calculation of the limit of the former sequence we have to transform the sequences of resolvents $R(0, 0, k, 0)$ under the Möbius transformations $g_{n,1}^{-1}g_{n,2}$. Therefore, the corresponding sequence of scaling parameters t_n should be inserted in the Taylor series of Lemma 3.35. Consequently, the arguments showing the convergence of a subsequence in case conditions *Decomposition* (i)–(ii) are fulfilled, carry over and show firstly that the sequence of resolvents $R(U_{\text{sing},n,2}, \bar{U}_{\text{sing},n,2}, k, 0)$ transformed under the Möbius transformations $g_{n,1}$ converges to an analogous *Finite rank Perturbation* of the resolvents $R_{\mathbb{P}^1}(0, 0, 0)$, and secondly that the sequence of resolvents $R(U_{\text{sing},n,1} + U_{\text{sing},n,2}, \bar{U}_{\text{sing},n,1} + \bar{U}_{\text{sing},n,2}, k, 0)$ transformed under the Möbius transformations $g_{n,1}$ converges to an analogous *Finite rank Perturbation* of $R_{\mathbb{P}^1}(\pi(g_{n,1}) U_{\text{sing},n,1}, \pi(g_{n,1}) \bar{U}_{\text{sing},n,1}, 0)$. In case that these *Finite rank Perturbations* are not defined, since the corresponding operators have non-trivial kernels, we determine the whole family of resolvents parametrized by λ of the Dirac-like operators mentioned above.

In a second step the sequence of resolvents

$$R(U_{\text{reg},n} + U_{\text{sing},n,1} + U_{\text{sing},n,2}, \bar{U}_{\text{reg},n} + \bar{U}_{\text{sing},n,1} + \bar{U}_{\text{sing},n,2}, k, 0)$$

is determined analogously to the case of a sequence obeying conditions *Decomposition* (i)–(ii). However, the *Finite rank perturbations* 3.22 of $R(U, \bar{U}, k, 0)$ are not induced by the kernels and co-kernels of that Dirac operator on $\mathbb{P}^1$, whose potentials are the weak limit of the sequence of potentials $(1 + z\bar{z})\pi(g_{n,1}) U_{\text{sing},n,1}$. Rather they are induced by the kernels and co-kernels of that *Finite rank perturbation* of the former Dirac operator on $\mathbb{P}^1$, which is the limit of the sequence of Dirac operators with potentials $(1 + z\bar{z})\pi(g_{n,1})(U_{\text{sing},n,1} + U_{\text{sing},n,1})$.

These arguments extend to the general case. In fact, we start with the most singular sequences of potentials. Inductively all kernels of the already determined *Finite rank Per-*

turbations on the domains excluded from one sequence of potentials give rise to *Finite rank Perturbations* of the corresponding sequence of Dirac operators on $\mathbb{P}^1$ and finally on $\mathbb{R}^2/\Lambda$. **q.e.d.**

Remark 3.51. Lemma 3.2 extends to these limits of sequences of potentials. More precisely, the restriction of the operator $\tilde{A}_f$ (defined in the proof of this lemma) to the complement of the union of small balls around the singular points $\mathcal{S}_\varepsilon$ (compare with Remark 3.36) is well defined for all *Finite rank Perturbations* 3.22. Since the Dirac operator is a local operator, the commutator in (ii) yields again the restriction of the corresponding variations $(\delta V_f, \delta W_f)$ to the complement of $\mathcal{S}_\varepsilon$. Moreover, for functions, whose support is contained in this complement, the statements (iii) and (iv) are still valid. Since the L^2 -functions, whose support are contained in the complement of some $\mathcal{S}_\varepsilon$ with $\varepsilon > 0$, are dense in the corresponding Hilber space, Proposition 3.5 extends as well to these *Finite rank Perturbations* 3.22.

3.5 The compactified moduli spaces

Due to the previous Section 3.4 all sequences in $\mathcal{M}_{\Lambda, \eta, \mathcal{W}}$ have subsequences, which converge to a *complex Fermi curve* of a *Finite rank Perturbation* 3.22. Since these moduli spaces $\mathcal{M}_{\Lambda, \eta, \mathcal{W}}$ are subspaces of a separable compact metrizable space (compare with Lemma 3.10), the closures $\bar{\mathcal{M}}_{\Lambda, \eta, \mathcal{W}}$ are the sets of all accumulation points of these spaces. Analogously $\bar{\mathcal{M}}_{\Lambda, \eta, \sigma, \mathcal{W}}$ denotes the subspace of $\bar{\mathcal{M}}_{\Lambda, \eta, \mathcal{W}}$, whose *complex Fermi curves* are invariant under the involution σ , and $\bar{\mathcal{M}}_{\Lambda, \eta}$ and $\bar{\mathcal{M}}_{\Lambda, \eta, \sigma}$ denote the unions

$$\bar{\mathcal{M}}_{\Lambda, \eta} = \bigcup_{\mathcal{W} > 0} \bar{\mathcal{M}}_{\Lambda, \eta, \mathcal{W}} \quad \bar{\mathcal{M}}_{\Lambda, \eta, \sigma} = \bigcup_{\mathcal{W} > 0} \bar{\mathcal{M}}_{\Lambda, \eta, \sigma, \mathcal{W}}.$$

In this section we shall investigate these compactified moduli spaces $\bar{\mathcal{M}}_{\Lambda, \eta, \mathcal{W}}$. Due to the *Strong unique continuation property* 2.10 we may replace in Lemma 3.11 the *compact open topology* of the resolvents $R(V, W, k, 0)$ by the *compact open topology* of the restrictions of the resolvents to the complement of $\mathcal{S}_\varepsilon$ (compare with Remark 3.36 and Theorem 3.48). Therefore Theorem 3.48 implies that all elements of $\bar{\mathcal{M}}_{\Lambda, \eta}$ are *complex Fermi curves* of *Finite rank Perturbations* 3.22. In particular, we shall see that the elements of $\bar{\mathcal{M}}_{\Lambda, \eta, \mathcal{W}}$ may be realized by gluing two planes $\mathbb{C}_p^\pm$ (or $\mathbb{C}_p^\pm$) along a combination of possibly infinitely many *horizontal cuts* and finitely many *pairs of parallel cuts* as described in Lemma 3.16.

Proposition 3.52. *The elements $\mathcal{F} \in \bar{\mathcal{M}}_{\Lambda, \eta}$ may be realized by gluing two planes $\mathbb{C}_p^\pm$ along at most infinitely many horizontal cuts and finitely many pairs of parallel cuts as described in Lemma 3.16. More precisely, for all $\mathcal{F} \in \bar{\mathcal{M}}_{\Lambda, \eta}$ and all small $\varepsilon > 0$ (the excluded discs $B(g(\hat{\gamma}, k_{\kappa^\pm}), \varepsilon)$ should be disjoint) there exists a $\delta > 0$, such that $\mathbb{C}_p^\pm$ does not contain any cuts in*

$$\begin{aligned} \{\hat{p} \in \mathbb{C} \mid |\hat{p}| > 1/\delta\} \bigcap_{\kappa \in \Lambda^*} \{\hat{p} \in \mathbb{C} \mid |\hat{p} - g(\hat{\gamma}, k_{\kappa^\pm})| > \varepsilon\} &= \\ = \{\hat{p} \in \mathbb{C} \mid |\hat{p}| > 1/\delta\} \setminus \bigcup_{\kappa \in \Lambda^* \setminus \{0\}} B(g(\hat{\gamma}, k_{\kappa^\pm}), \varepsilon) &, \end{aligned}$$

and at most one horizontal cut in each of the excluded discs $B(g(\hat{\gamma}, k_{\kappa^{\pm}}), \varepsilon)$ indexed by $\kappa \in \Lambda_{\delta}^* = \{\kappa \in \Lambda^* \mid |g(\hat{\gamma}, k_{\kappa^+})| > 1/\delta\}$. Finally, the disc $B(0, 1/\delta)$ contains finitely many horizontal cuts and pairs of parallel cuts. The total number of these cuts is bounded by the number of $\kappa \in \Lambda^*$, whose discs $B(g(\hat{\gamma}, k_{\kappa^{\pm}}), \varepsilon)$ are not disjoint from $B(0, 1/\delta)$.

In spite of being in general different from the similar index set in Theorem 2.12 we denote the index set Λ_{δ}^* by the same symbol. Since $|g(\hat{\gamma}, k_{\kappa^+})| = \frac{\|\hat{\gamma}\|}{2} \|\kappa\| = \frac{\|\hat{\gamma}\|}{\sqrt{2}} \|k_{\kappa^+}\| = \frac{\|\hat{\gamma}\|}{\sqrt{2}} \|k_{\kappa^-}\|$ the transformation $\delta \mapsto \frac{\|\hat{\gamma}\|}{\sqrt{2}} \delta$ transforms the latter index set into the former. Moreover, if the cuts are not disjoint, then the total number of these cuts is given by the half of the degree of the zero divisor of $d\hat{p}$ restricted to the corresponding set of the *complex Fermi curve*.

Proof. In a first step we improve the arguments of Theorem 2.12 and show that all *complex Fermi curves* in $\mathcal{M}_{\Lambda, \eta, W}$ have a representation of the form just described. Due to Theorem 2.12 near infinity the *complex Fermi curve* looks like two open planes, which are connected by infinitely many small handles. We shall show that these handles may be represented by *horizontal cuts*. Due to arguments used in the proof of Theorem 2.12 these handles are small perturbations of the ordinary double point of the free *complex Fermi curve* at $k = 0$. For $k = 0$ the kernel of $\tilde{D}(0, 0, k)$ is the two-dimensional space of the constant functions ($\sim \mathbb{C} \times \mathbb{C} \subset L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$). Therefore, we decompose the resolvent $\tilde{R}(0, 0, k, \lambda)$ into a sum of the reduced resolvent plus the singular matrix $\begin{pmatrix} \lambda & -\pi(k_2 + \sqrt{-1}k_1) \\ -\pi(k_2 - \sqrt{-1}k_1) & \lambda \end{pmatrix}^{-1}$ acting on the constant functions:

$$\tilde{R}(0, 0, k, \lambda) = \tilde{R}_{\text{red}}(0, 0, k, \lambda) + \tilde{S}(0, 0, k, \lambda).$$

For small k the reduced resolvent is holomorphic and bounded and the singular part is a rank-two operator with a pole at $\lambda^2 = \pi^2 g(k, k)$. Consequently the resolvent $\tilde{R}(V, W, k, \lambda)$ has the form

$$\begin{aligned} \tilde{R}(V, W, k, \lambda) &= \tilde{R}_{\text{red}}(0, 0, k, \lambda) \left(\mathbf{1} - \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, \lambda) \right)^{-1} + \\ &+ \left(\mathbf{1} - \tilde{R}_{\text{red}}(0, 0, k, \lambda) \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \right)^{-1} \tilde{S}(V, W, k, \lambda) \left(\mathbf{1} - \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, \lambda) \right)^{-1}. \end{aligned}$$

Here $\tilde{S}(V, W, k, \lambda)$ is the following 2×2 -matrix (acting on the constant functions):

$$\begin{aligned} \tilde{S}(V, W, k, \lambda) &= \\ &= \tilde{S}(0, 0, k, \lambda) \left(\mathbf{1} - \left(\mathbf{1} - \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, \lambda) \right)^{-1} \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \tilde{S}(0, 0, k, \lambda) \right)^{-1} \\ &= \left(\mathbf{1} - \tilde{S}(0, 0, k, \lambda) \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \left(\mathbf{1} - \tilde{R}_{\text{red}}(0, 0, k, \lambda) \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \right)^{-1} \right)^{-1} \tilde{S}(0, 0, k, \lambda). \end{aligned}$$

The arguments in Section 2.1 carry over and show that for small $V, W \in L^2(\mathbb{R}^2/\Lambda)$ the operators $\mathbf{1} - \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, \lambda)$ and $\mathbf{1} - \tilde{R}_{\text{red}}(0, 0, k, \lambda) \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix}$ are invertible. More precisely,

the restriction of the operator $\tilde{R}_{\text{red}}(0, 0, k, \lambda) \left(\mathbf{1} - \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, \lambda) \right)^{-1}$ to the orthogonal complement of the constant functions is the resolvent of the restriction of $\tilde{D}(V, W, k)$ to this orthogonal complement.

Remark 3.53. *Dirac operators with potentials $\tilde{D}(V, W, k)$ do not preserve this orthogonal complement in contrast to the corresponding free Dirac operators $\tilde{D}(0, 0, k)$. But with the help of the natural inclusion of the orthogonal complement of the constant functions into $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ and the natural projection of this Banach space onto the orthogonal complement of the constant functions the restriction to this complement is well defined.*

By the first resolvent formula [R-S-I, Theorem VI.5] we may express the resolvent $R_{\lambda'}$ with argument λ' in terms of the resolvent R_{λ} with argument λ and the resolvent with argument $\frac{1}{\lambda - \lambda'}$ of the resolvent with argument λ :

$$(\lambda - \lambda')R_{\lambda'} = R_{\lambda} \left(\frac{\mathbf{1}}{\lambda - \lambda'} - R_{\lambda} \right)^{-1} = \left(\frac{\mathbf{1}}{\lambda - \lambda'} - R_{\lambda} \right)^{-1} R_{\lambda}.$$

Now the operators $\tilde{D}(0, 0, k)\tilde{R}_{\text{red}}(0, 0, k, \lambda)$ are invertible on $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ with $1 < p < 2$, and the operators $\tilde{R}_{\text{red}}(0, 0, k, \lambda)\tilde{D}(0, 0, k)$ are invertible on $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$ with $2 < q < \infty$. Therefore the arguments of Section 2.1 show that the operators $\mathbf{1} - \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, \lambda)$ on $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ and the operators $\mathbf{1} - \tilde{R}_{\text{red}}(0, 0, k, \lambda) \begin{pmatrix} V & 0 \\ 0 & W \end{pmatrix}$ on $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$ are invertible, if and only if λ does not belong to the spectrum of the restriction of $\tilde{D}(V, W, k)$ to the orthogonal complement of the constant functions. Furthermore, the proof of Lemma 2.6 shows that the inverse of the former operators considered as operators from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ to $L^{p'}(\mathbb{R}^2/\Lambda) \times L^{p'}(\mathbb{R}^2/\Lambda)$ with $1 < p' < p < 2$ and the inverse of the latter operators considered as operators from $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$ to $L^{q'}(\mathbb{R}^2/\Lambda) \times L^{q'}(\mathbb{R}^2/\Lambda)$ with $2 < q' < q < \infty$ depend weakly continuously on the potentials on the subsets described in Lemma 2.6. Hence for all potentials $V, W \in L^2(\mathbb{R}^2/\Lambda)$ and all small open neighbourhoods $\mathcal{O} \subset \mathbb{C}^2$ of 0 and all $\varepsilon > 0$ there exists a $\delta > 0$ such that for all $\kappa \in \Lambda_{\delta}^*$ and all $k \in \mathcal{O}$ the restriction of the operators

$$\begin{aligned} & \left(\mathbf{1} - \begin{pmatrix} \psi_{-\kappa}V & 0 \\ 0 & \psi_{\kappa}W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, 0) \right)^{-1} \begin{pmatrix} \psi_{-\kappa}V & 0 \\ 0 & \psi_{\kappa}W \end{pmatrix} \\ &= \begin{pmatrix} \psi_{-\kappa}V & 0 \\ 0 & \psi_{\kappa}W \end{pmatrix} \left(\mathbf{1} - \tilde{R}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa}V & 0 \\ 0 & \psi_{\kappa}W \end{pmatrix} \right)^{-1} \end{aligned}$$

to the constant functions defines a 2×2 -matrix $A(\psi_{-\kappa}V, \psi_{\kappa}W, k)$, which is bounded by ε . More precisely, these 2×2 -matrices $A(\psi_{-\kappa}V, \psi_{\kappa}W, k)$ are defined as the matrix elements of the given operators with respect to the natural orthonormal basis of the constant functions $\simeq \mathbb{C} \times \mathbb{C} \subset L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$. Due to Lemma 3.21, for all $\kappa \in \Lambda_{\delta}^*$ and all $k \in \mathcal{O}$ the spectrum of the operator $D(\psi_{-\kappa}v, \psi_{\kappa}W, k)$ is given by the polar set of the inverse of the 2×2 -matrices $\mathbf{1} + \begin{pmatrix} 0 & \pi(k_2 + \sqrt{-1}k_1) \\ \pi(k_2 - \sqrt{-1}k_1) & 0 \end{pmatrix}^{-1} A(\psi_{-\kappa}V, \psi_{\kappa}W, k)$. Obviously this polar set is equal to the zero set of the determinant

$$\det \left(\begin{pmatrix} 0 & \pi(k_2 + \sqrt{-1}k_1) \\ \pi(k_2 - \sqrt{-1}k_1) & 0 \end{pmatrix} + A(\psi_{-\kappa}V, \psi_{\kappa}W, k) \right).$$

Since the operator $\tilde{\mathbf{F}}\tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0)\mathbf{F}^{-1}$ acts on $\ell_{\frac{p}{p-1}}(\Lambda^*) \times \ell_{\frac{p}{p-1}}(\Lambda^*)$ as the multiplication with the following sequence of matrices indexed by $\kappa \in \Lambda^*$:

$$\begin{cases} 0 & \text{if } \kappa = 0, \\ \frac{-1}{\pi} \begin{pmatrix} 0 & (k_2 + \kappa_2 - \sqrt{-1}(k_1 + \kappa_1))^{-1} \\ (k_2 + \kappa_2 + \sqrt{-1}(k_1 + \kappa_1))^{-1} & 0 \end{pmatrix} & \text{if } \kappa \neq 0, \end{cases}$$

the partial derivatives of this operator with respect to k_1 and k_2 acts as the multiplication with the partial derivatives of this sequence of matrices, respectively. The arguments in the proof of Lemma 2.6 concerning the operators $\mathbf{F}\mathbf{R}(0, 0, k, \sqrt{-1}\lambda)\mathbf{F}^{-1}$ imply that the partial derivatives of the operator-valued function $k \mapsto \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0)$ define for all $1 < p < q < \infty$ and all small k bounded operators from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ into $L^q(\mathbb{R}^2/\Lambda) \times L^q(\mathbb{R}^2/\Lambda)$. Therefore, the arguments concerning the convergence of the matrices $\mathbf{A}(\psi_{-\kappa}V, \psi_{\kappa}W, k)$ in the limit $g(\kappa, \kappa) \rightarrow \infty$ also imply that the partial derivatives of the function $k \mapsto \mathbf{A}(\psi_{-\kappa}V, \psi_{\kappa}W, k)$ converge uniformly with respect to $k \in \mathcal{O}$ to zero in the limit $g(\kappa, \kappa) \rightarrow \infty$.

Consequently for all $V, W \in L^2(\mathbb{R}^2/\Lambda)$ there exists a $\delta > 0$ and a sequence k_{κ} of elements of $\mathcal{O}$ indexed by Λ_{δ}^* such that the values of the matrix-valued functions $k \mapsto \begin{pmatrix} 0 & \pi(k_2 + \sqrt{-1}k_1) \\ \pi(k_2 - \sqrt{-1}k_1) & 0 \end{pmatrix} + \mathbf{A}(\psi_{-\kappa}V, \psi_{\kappa}W, k)$ at $k = k_{\kappa}$ are diagonal matrices. In fact, the bounds on the partial derivatives of the function $k \mapsto \mathbf{A}(\psi_{-\kappa}V, \psi_{\kappa}W, k)$ imply that the mapping, which maps k to the unique $k' \in \mathcal{O}$ such that $\begin{pmatrix} 0 & \pi(k'_2 + \sqrt{-1}k'_1) \\ \pi(k'_2 - \sqrt{-1}k'_1) & 0 \end{pmatrix} + \mathbf{A}(\psi_{-\kappa}V, \psi_{\kappa}W, k)$ is a diagonal 2×2 -matrix, is a contraction. Therefore, due to the contraction mapping principle [R-S-I, Theorem V.18], this mapping has a unique fixed point, which yields the desired k_{κ} . Let $\hat{v}(V, W, \kappa)$ and $\hat{w}(V, W, -\kappa)$ denote the diagonal elements of these matrices:

$$\mathbf{A}(\psi_{-\kappa}V, \psi_{\kappa}W, k_{\kappa}) = \begin{pmatrix} \hat{v}(V, W, \kappa) & 0 \\ 0 & \hat{w}(V, W, -\kappa) \end{pmatrix}.$$

We conclude that the intersections $\mathcal{F}(\psi_{-\kappa}V, \psi_{\kappa}W) \cap \mathcal{O}$ are small perturbations of the subvarieties $\{k \in \mathcal{O} \mid g(k - k_{\kappa}, k - k_{\kappa}) = \hat{v}(V, W, \kappa)\hat{w}(V, W, -\kappa)\}$. In fact, the bounds of the partial derivatives of the function $k \mapsto \mathbf{A}(\psi_{-\kappa}V, \psi_{\kappa}W, k)$ imply for all $\kappa \in \Lambda_{\delta}^*$ the estimates

$$\left\| \mathbf{A}(\psi_{-\kappa}V, \psi_{\kappa}W, k) - \begin{pmatrix} \hat{v}(V, W, \kappa) & 0 \\ 0 & \hat{w}(V, W, -\kappa) \end{pmatrix} \right\| \leq \varepsilon \|k - k_{\kappa}\|$$

uniformly for all $k \in \mathcal{O}$.

If the mapping $k \mapsto \|\mathbf{A}(\psi_{-\kappa}V, \psi_{\kappa}W, k)\|$ and the corresponding partial derivatives of first order have on the small ball $\mathcal{O} = B(k_{\kappa}, \varepsilon')$ an upper bound ε , which is small compared with ε' , then this zero set defines a two-sheeted covering over a small ball $\hat{p} \in B(g(\hat{\gamma}, k_{\kappa}), \varepsilon'')$ with $0 < \varepsilon \ll \varepsilon'' \ll \varepsilon'$. In particular, on $B(g(\hat{\gamma}, k_{\kappa}), \varepsilon'')$ this two-sheeted covering has either an ordinary double point connecting the two sheets and no other zero of $d\hat{p}$, or two simple branch point connecting the two sheets and no other zero of $d\hat{p}$. Again Lemma 2.11 shows that this yields an asymptotic description of the *complex Fermi curves*

In case of a pair of potentials of the form $(U, \bar{U})$, the map η transforms the matrices $\mathbf{A}(\psi_{-\kappa}U, \psi_{\kappa}\bar{U}, k)$ like $\bar{\mathbf{A}}(\psi_{-\kappa}U, \psi_{\kappa}\bar{U}, k) = \mathbf{J}\mathbf{A}(\psi_{-\kappa}U, \psi_{\kappa}\bar{U}, -\bar{k})\mathbf{J}^{-1}$ (compare with Section 2.3). Consequently the corresponding sequences k_{κ} are fixed points of η , and the sequences $\hat{w}(U, \bar{U}, -\kappa)$ are the complex conjugate of the sequences $\hat{v}(U, \bar{U}, \kappa)$, which we denote

by $\hat{u}(U, \kappa)$. The invariance of a *complex Fermi curve* of potentials of the form $(U, \bar{U})$ under the involution η implies that in case of vanishing $\hat{u}(U, \kappa)$ this part of the *complex Fermi curve* gives rise to an ordinary double point and no cut and in case of non-vanishing $\hat{u}(U, \kappa)$ to one *horizontal cut*. On the other hand, Theorem 2.12 implies that the sets $\mathcal{V}_{\varepsilon, \delta}^{\pm}$ are mapped by $\hat{p}$ biholomorphically onto open subsets of $\mathbb{C}_{\hat{p}}^{\pm}$. Finally, the continuity of the zeroes of $d\hat{p}$ under deformations of $U \in L^2(\mathbb{R}^2/\Lambda)$ implies that the arithmetic genus of $\mathcal{F}$ is preserved. More precisely, if we count the singularities as zeroes of $d\hat{p}$, whose order is equal to twice the *Local contribution to the arithmetic genus* 4.3, then the total order of zeroes of $d\hat{p}$ is locally preserved and equal to twice the number of elements of Λ^* . Therefore, $\mathcal{V}_{\varepsilon, \delta}^{\pm}$ has no zeroes of $d\hat{p}$, the small handles excluded from these sets have two zeroes of $d\hat{p}$ (either as an ordinary double point or as two simple branch points) and the number of zeroes of $d\hat{p}$ on the compact part is twice the number of elements of $\Lambda^* \setminus \Lambda_{\delta}^*$. This proves the theorem for all elements $\mathcal{F} \in \mathcal{M}_{\Lambda, \eta, \mathcal{W}}$.

Due to Theorem 3.48 the accumulation points of sequences in $\bar{\mathcal{M}}_{\Lambda, \eta, \mathcal{W}}$ are *complex Fermi curves* of *Finite rank Perturbations* 3.22 of Dirac operators $D(U, \bar{U}, k)$ with complex potentials in $\{U \in L^2(\mathbb{R}^2/\Lambda) \mid 4\|U\|_2^2 \leq \mathcal{W}\}$. In the eighth step of Section 3.4.1 we extended the asymptotic analysis of Theorem 2.12 to these *Finite rank Perturbations* 3.22. Due to the Strong unique continuation property 2.10 we may use the restrictions (compare with Remark 3.36) of the corresponding resolvents to the complement of small balls around the singular points and the corresponding norms. Therefore, also the *complex Fermi curves* of *Finite rank Perturbations* 3.22 have a representation of the prescribed form. **q.e.d.**

Due to this theorem all *complex Fermi curves* with dividing real parts have finite genus. Moreover, all $\mathcal{F} \in \bar{\mathcal{M}}_{\Lambda, \eta, \sigma, \mathcal{W}}$ with dividing real parts have a small open neighbourhood $\mathcal{U}$ with respect to the subspace topology of the closed subsets of $\bar{\mathbb{C}}^2$ (compare with Lemma 3.10) such that the *geometric genera* of all *complex Fermi curves* in $\mathcal{U}$ with dividing real parts are bounded. In fact, with the exception of finitely many indexes κ the corresponding handles of a *complex Fermi curve* with dividing real part have ordinary double points of the form $(y, \eta(y))$. Furthermore, a small bound on the deformations of the functions $k_1 \pm \sqrt{-1}k_2$ on the compact part guarantees that these ordinary double points are deformed into ordinary double points of the same form, since a deformation into *horizontal cuts* contradicts the dividing property of the real part. On the other hand, the subspace of *complex Fermi curves* with dividing real parts is closed, since a small deformation of a cut connecting the two planes $\mathbb{C}_{\hat{p}}^{\pm}$ disjoint from the real part yields a cut with the same properties. Hence the compactness of $\bar{\mathcal{M}}_{\Lambda, \eta, \mathcal{W}}$ implies

Corollary 3.54. *For all $\mathcal{W} > 0$ there exists a natural number $g_{\max} \in \mathbb{N}$ depending only on the lattice Λ and $\mathcal{W}$ such that all complex Fermi curves $\mathcal{F} \in \bar{\mathcal{M}}_{\Lambda, \eta, \sigma, \mathcal{W}}$ with dividing real parts are contained in $\bar{\mathcal{M}}_{g_{\max}, \Lambda, \eta, \sigma, \mathcal{W}}$.* **q.e.d.**

Remark 3.55. *These numbers $g_{\max}$ are not bounded uniformly for all lattices and become arbitrary large, whenever $\text{vol}(\mathbb{R}^2/\Lambda) \min\{g(\kappa, \kappa) \mid \kappa \in \Lambda^* \setminus \{0\}\}$ becomes arbitrary small. In fact, for all $\kappa \in \Lambda^* \setminus \{0\}$ with $\pi^2 \text{vol}(\mathbb{R}^2/\Lambda)g(\kappa, \kappa) \leq \mathcal{W}$ the complex Fermi curves of the constant potentials have one real double point.*

Proposition 3.52 suggests the following characterization of the elements of $\bar{\mathcal{M}}_{\Lambda, \eta, \mathcal{W}}$. Instead of the data $(Y, \infty^-, \infty^+, k)$ we consider all data (Y, k) of complex spaces Y together with a holomorphic multi-valued function k on Y , which satisfies conditions *Quasi-momenta* (ii)–(iii) of Section 2.5 and the following conditions *Quasi-momenta* (i') and *Quasi-momenta* (vi).

Quasi-momenta (i') For both components $\hat{p}$ and $\check{p}$ of k the Riemann surface Y may be realized by gluing two planes $\mathbb{C}_{\hat{p}}^{\pm}$ (or $\mathbb{C}_{\check{p}}^{\pm}$) along a countable combination of *horizontal cuts* and *pairs of parallel cuts* as described in Lemma 3.16. Moreover, there exists some branch of the function k with the property that the function $k_1 + \sqrt{-1}k_2$ on $\mathbb{C}_{\hat{p}}^+$ converges to zero, whenever the function $\hat{p}$ tends to infinity.

Quasi-momenta (vi) The integral of the two-form

$$2\pi\sqrt{-1} \operatorname{vol}(\mathbb{R}^2/\Lambda) (dk_1 + \sqrt{-1}dk_2) \wedge \overline{(dk_1 + \sqrt{-1}dk_2)}$$

over $\mathbb{C}_{\hat{p}}^+$ is less or equal to $\mathcal{W}$.

Due to Proposition 3.52 the elements of $\bar{\mathcal{M}}_{\Lambda, \eta}$ obey these conditions. We conjecture that these conditions characterize the elements of $\mathcal{M}_{\Lambda, \eta, \mathcal{W}}$ for all $\mathcal{W} > 0$.

Let us remark, that the mapping

$$(V, W) \mapsto (\hat{v}(V, W, \kappa), \hat{w}(V, W, \kappa))_{\kappa \in \Lambda_{\delta}^*}$$

is a nonlinear perturbation of the Fourier transform. For small $\|V\|$ and $\|W\|$ the sequences $\hat{v}(V, W, \kappa)$ and $\hat{w}(V, W, \kappa)$ are defined for all $\kappa \in \Lambda^*$.

Remark 3.56. *In order to define all entries of these sequences, we have to ensure that the handles connecting the parts near k_{κ}^+ with the part near k_{κ}^- of the complex Fermi curve do not overlap. One can show that an upper bound on $4\|V\|_2^2$ and $4\|W\|_2^2$, which prevents such an overlapping essentially is given by $\pi^2 \operatorname{vol}(\mathbb{R}^2/\Lambda) \min\{g(\kappa, \kappa) \mid \kappa \in \Lambda^* \setminus \{0\}\}$. Moreover, there does not exist a non-vanishing upper bound, which prevents this overlapping uniformly for all lattices Λ (compare with Remark 3.55).*

For the zero potentials the functions $A(0, 0, k)$ vanishes identically. Furthermore, the derivative of the mapping $(V, W) \mapsto (\hat{v}(V, W, \kappa), \hat{w}(V, W, \kappa))_{\kappa \in \Lambda^*}$ at the zero potentials is equal to the Fourier transform. An easy calculation shows that the *complex Fermi curves* of pairs of potentials of the form $(u\psi_{\kappa}, \bar{u}\psi_{-\kappa})$ with $u \in \mathbb{C}$ are equal to

$$\mathcal{F}(u\psi_{\kappa}, \bar{u}\psi_{-\kappa}) = \bigcup_{\kappa' \in \Lambda^*} \{k \in \mathbb{C}^2 \mid \pi^2 g(k + k_{\kappa}^+ + \kappa', k + k_{\kappa}^+ + \kappa') = u\bar{u}\}.$$

Moreover, with the help of the arguments of the proof of Lemma 2.26 it is easy to see that if $4u\bar{u} \leq \min\{g(\kappa, \kappa) \mid \kappa \in \Lambda^* \setminus \{0\}\}$ ($\iff \mathcal{W} \leq \pi^2 \operatorname{vol}(\mathbb{R}^2/\Lambda) \min\{g(\kappa, \kappa) \mid \kappa \in \Lambda^* \setminus \{0\}\}$) these pairs of potentials $(u\psi_{\kappa}, \bar{u}\psi_{-\kappa})$ are all potentials of the form $(U, \bar{U})$, whose *complex Fermi curves* have *geometric genus* zero. This leads to a local description of the mapping

$$\{U \in L^2(\mathbb{R}^2/\Lambda) \mid 4\|U\|_2^2 \leq \mathcal{W}\} \rightarrow \mathcal{M}_{\Lambda, \eta, \mathcal{W}}, \quad U \mapsto \mathcal{F}(U, \bar{U}).$$

We shall see that for small W this mapping is a nonlinear perturbation of the mapping

$$L^2(\mathbb{R}^2/\Lambda) \rightarrow \ell_1^+(\Lambda^*) = \{t \in \ell_1(\Lambda^*) \mid t(\kappa) \in \mathbb{R}_0^+ \forall \kappa \in \Lambda^*\}, \quad U \mapsto t, \text{ with } t(\kappa) = |\hat{U}(\kappa)|^2.$$

Here $\hat{U}$ denotes the Fourier transform of U . Moreover, together with the symplectic form, which is equal to the imaginary part of the Hermitian form of $L^2(\mathbb{R}^2/\Lambda)$, this mapping defines an infinite-dimensional completely integrable system. This is a general feature of integrable systems related to Lax operators. The mappings from the potentials to the spectral curves (i.e. a complete sets of integrals) of these integrable systems are nonlinear perturbations of the mapping from the potentials to the absolute values of their Fourier coefficients, which together with the corresponding symplectic form represents simple examples of infinite-dimensional integrable systems. The preimages of the latter map are compact tori, whose dimensions are equal to the number of non-vanishing moduli. The preimages of the former map are the *isospectral sets*, which in general may be identified with open sets of the corresponding Jacobian varieties (compare with Lemma 2.27). Therefore, these dimensions are equal to the genera of the corresponding *complex Fermi curves*. Due to a theorem of Littlewood ([Li, F-R] and [Kis, Chapter 3 §1.6.]) almost all elements of preimages of the latter map have finite L^q -norm for all finite q . This leads to the conjecture that all *complex Fermi curves* $\mathcal{F}(U, \bar{U})$ are equal to the *complex Fermi curve* of some potential $U' \in L^q(\mathbb{R}^2/\Lambda)$, and that the L^q -norm of U' is bounded by some constant depending on q and the maximum of all $\varepsilon > 0$ such that $\|U\|_{\varepsilon, 2}$ is smaller than S_p^{-1} (compare with Lemma 2.6).

A precise version of this perturbation is that the moduli space may be realized by gluing homeomorphically open subsets of $\ell_1^+(\Lambda^*)$. Moreover, these parameterizations of the moduli space has the property, that the genus of the corresponding *complex Fermi curves* is equal to the number of non-vanishing moduli. We remark that these local parameterizations of the moduli space endows it with a finer topology than the topology introduced in Section 3.3. In contrast to the former topology with respect to which the mapping $U \mapsto \mathcal{F}(U, \bar{U})$ is ‘almost’ weakly continuous, the latter topology is the finest topology, such that this mapping is continuous with respect to the usual L^2 -topology. Let us call a topology of the moduli space with the property, that each element of the moduli space has an open neighbourhood, which is homeomorphic to an open subset of $\ell_1^+(\Lambda^*)$, a $\ell_1^+(\Lambda^*)$ -structure.

Theorem 3.57. *The moduli space $\mathcal{M}_\Lambda = \{\mathcal{F}(V, W) \mid V, W \in L^2(\mathbb{R}^2/\Lambda)\}$ is a holomorphic $\ell_1(\Lambda^*)$ -manifold. Furthermore, locally there exists a $\delta > 0$, such that the moduli with indexes $\kappa \in \Lambda_\delta^*$ are unique. Moreover, the compactified moduli space $\bar{\mathcal{M}}_{\Lambda, \eta}$ has an asymptotic $\ell_1^+(\Lambda^*)$ -structure (i.e. it is a closed subset of a $\ell_1(\Lambda^*)$ -manifold, and the restriction to the moduli with indexes $\kappa \in \Lambda_\delta^*$ with a suitable δ yields an open mapping into $\ell_1^+(\Lambda_\delta^*)$). Again these moduli are asymptotically unique. Finally, the complex Fermi curves of finite genus are parameterized by elements of $\ell_1^+(\Lambda^*)$ with finitely many non-vanishing entries. In particular, they are dense with respect to the corresponding topologies.*

Proof. The proof of this theorem is a consequence of the improved asymptotic analysis of Proposition 3.52. Again we first treat the subspace $\mathcal{M}_{\Lambda, \eta}$ and then in a second step we extend our arguments to $\mathcal{M}_{\Lambda, \eta}$.

1. First we define the moduli $t(\kappa)$ for all all *complex Fermi curves* of pairs of potentials of the form $(U, \bar{U})$ and $\kappa \in \Lambda_\delta^*$, with an appropriate δ (depending on U) defined in the proof of Proposition 3.52.

$$t(\kappa) = \pi \oint_{\text{horizontal cut of } \mathbb{C}_p^+ \text{ with index } \kappa.} k_1 dk_2.$$

Due to Lemma 3.16 these *horizontal cuts* are invariant under η . Therefore, the integrals of dk along these cycles vanish. Also the integral in the definition of $t(\kappa)$ does not depend on the branch of k . Moreover, due to the invariance under η , the well defined moduli $t(\kappa)$ are real. Since the form $k_1 dk_2$ is closed, we may integrate it along any closed cycles of an open set of the form $\mathcal{V}_{\varepsilon, \delta}^+$ with an appropriate $\varepsilon > 0$ and $\delta > 0$ around the excluded disc near k_κ^+ . This extends the definition of the moduli to all pairs of potentials with $V, W \in L(\mathbb{R}^2/\Lambda)$.

Due to Lemma 3.12 the contribution of the *horizontal cut* with index κ to the *first integral* is equal to $4 \text{vol}(\mathbb{R}^2/\Lambda) t(\kappa)$. Consequently, in case that all these handles do not overlap and all moduli $t(\kappa)$ are well defined, the *first integral* $4 \int_{\mathbb{R}^2/\Lambda}$ is equal to $4 \text{vol}(\mathbb{R}^2/\Lambda) \sum_{\kappa \in \Lambda^*} t(\kappa)$,

whenever this sum converges.


2. Now we claim that for pairs of potentials of the form $(U, \bar{U})$ the well-defined $t(\kappa)$ are positive, which imply that they belong to $\ell_1^+(\Lambda_\delta^*)$. The real part of k defines a mapping from the $\mathcal{F}(U, \bar{U})$'s into $\mathbb{R}^2$ (which may depend on the branch of k). An easy calculation shows that this mapping is locally a homeomorphism, if and only if $d\check{p}/d\hat{p}$ is not real. Due to Proposition 3.52 asymptotically the function $d\check{p}/d\hat{p}$ is real only on small disjoint cycles, which are homolog to the *horizontal cuts* indexed by Λ_δ^* . Moreover, all these cycles are invariant under η and on each of these cycles the function $d\check{p}/d\hat{p}$ takes all values of $\mathbb{P}_{\mathbb{R}}^1$ exactly twice. Since η does not have fixed points, this implies that these cycles are mapped by the real part of k onto smooth Jordan curves in $\mathbb{R}^2/\Lambda^*$ around some element of the form $\kappa/2$ (which are the images of fixed points of η under the real part of k). If we cut the complement of the compact part of the *complex Fermi curve* described in Theorem 2.12 along these cycles into two pieces, then the real part of the unique branch of k , whose linear combinations $k_1 \pm \sqrt{-1}k_2$ become at $\infty^\pm$ arbitrary small, respectively, maps these two components homeomorphically onto the complement of some compact part of $\mathbb{R}^2$ and those unique discs around $\kappa/2$ and $-\kappa/2$, whose boundary is the Jordan curve with index κ , respectively. Finally, they are glued along these Jordan curves by the map $k \mapsto k - \kappa$. The arguments of the proof Lemma 3.16 show that $t(\kappa)$ is equal to π times the volume of the corresponding disc in $\mathbb{R}^2$, whose boundary is the image of the cycle with index κ under the real part of k , plus or minus π times the volume of the analogous disc, whose boundary is the images of this cycle under the imaginary part of k . Furthermore, these arguments also show that $t(\kappa)$ is equal to π times the volume of the unique disc in $\mathbb{R}^2$, whose boundary is the image of the *horizontal cut* with index κ under the real part of k . But locally in a small tubular neighbourhood of the unique cycle with index κ , on which $d\check{p}/d\hat{p}$ takes real values, the real part of k takes values in the complement of the disc, whose boundary is the image of this cycle. We conclude that $t(\kappa)$ is larger or equal to the volume of this disc and therefore non-negative. Finally, $t(\kappa)$ vanishes if and only if the corresponding part of the *complex Fermi curve* has an ordinary double point and does not

have a *horizontal cut*. To sum up, we have proven that the mapping $U \mapsto t$ maps $L^2(\mathbb{R}^2/\Lambda)$ into $\ell_1^+(\Lambda_\delta^*)$ (with some suitable δ depending on U), and $4 \operatorname{vol}(\mathbb{R}^2/\Lambda) \|t\|_1 \leq 4 \|U\|_2^2$.

These arguments show that image under the imaginary part of k of those unique cycles, on which $d\check{p}/d\hat{p}$ takes real values, have negative ‘volume’. Since the imaginary part of k is preserved under η , the imaginary part of k maps these cycles not injectively into $\mathbb{R}^2$. In fact, for the constant potentials the unique non-trivial cycle is real and the imaginary part of k maps this cycle onto the origin. In general the preimages contain at least two points of the *complex Fermi curves*, which are interchanged by η . Therefore, the images of these cycles under the imaginary part of k look like smooth curves

$$S^1 \rightarrow \mathbb{R}^2 \quad \exp(2\pi\sqrt{-1}\varphi) \mapsto (x, y),$$

with the property that the quotient $dy/dx = (dy/d\varphi)/(dx/d\varphi)$ takes each value in $\mathbb{P}_{\mathbb{R}}^1$ exactly once. Such curves cannot be immersed (i. e. they must have common zeroes of $dy/d\varphi$ and $dx/d\varphi$), and the exterior parts of these curves are locally convex in the complement of the non-immersed points. This yields another explanation why the corresponding contribution to the *first integral* is negative. The figure on the right gives the image of a non-trivial example.



3. Now we shall show that the mapping $(V, W) \mapsto t$ is an open mapping from $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ into $\ell_1(\Lambda_\delta^*)$ with suitable δ depending on V and W . For this purpose we shall first prove that the mapping $(V, W) \mapsto (\hat{v}(V, W, \cdot), \hat{w}(V, W, \cdot))$ is an open mapping from $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ into $\ell_2(\Lambda_\delta^*) \times \ell_2(\Lambda_\delta^*)$ with suitable δ depending on V and W .

Lemma 3.58. *The derivatives of the mapping $(V, W) \mapsto (\hat{v}(V, W, \cdot), \hat{w}(V, W, \cdot))$ are bounded operators from $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ into $\ell_2(\Lambda_\delta^*) \times \ell_2(\Lambda_\delta^*)$ (with suitable δ depending on (V, W))*

Proof. First we investigate for the same $k \in \mathcal{O}$ as in the proof of Proposition 3.52 in three steps the derivative of the mapping

$$(V, W) \mapsto A_{V,W,k} \quad \text{with } A_{V,W,k}(x) = \sum_{\kappa \in \Lambda_\delta^*} \psi_\kappa(x) A(\psi_{-\kappa} V, \psi_\kappa W, k) \quad \forall x \in \mathbb{R}^2/\Lambda$$

In doing so we have to estimate the Fourier transform $A_{V,W,k}$ of the sequence of matrices $A(\psi_\kappa V, \psi_\kappa W, k)$ indexed by $\kappa \in \Lambda_\delta^*$. For this purpose we make use of the Bochner q -integrable functions from $\mathbb{R}^2/\Lambda$ into some Banach space E . They are denoted by $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} E$ [DF, Section 7.]. More precisely, we shall consider sequences of elements of $L^q(\mathbb{R}^2/\Lambda)$ indexed by $\kappa \in \Lambda^*$ (or Λ_δ^*) as elements of $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} \ell.(\Lambda^*)$. Consequently, the Fourier transform $\mathbb{1} \otimes F$ maps such tensor products into the tensor products $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L(\mathbb{R}^2/\Lambda)$. The derivative

$\frac{\delta A(\psi_{-\kappa}V, \psi_{\kappa}W, k)}{(\delta V, \delta W)}$ is the restriction of the following operator to the constant functions:

$$\left(\mathbf{1} - \begin{pmatrix} \psi_{-\kappa}V & 0 \\ 0 & \psi_{\kappa}W \end{pmatrix} \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0) \right)^{-1} \begin{pmatrix} \psi_{-\kappa}\delta V & 0 \\ 0 & \psi_{\kappa}\delta W \end{pmatrix} \left(\mathbf{1} - \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa}V & 0 \\ 0 & \psi_{\kappa}W \end{pmatrix} \right)^{-1}.$$

1. In a first step we show that for all $k \in \mathcal{O}$ and for all $1 < p < 2$ and $1/p + 1/q = 1$ both entries of the functions

$$(x, x') \mapsto \sum_{\kappa \in \Lambda^*} \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa}V & 0 \\ 0 & \psi_{\kappa}W \end{pmatrix} \begin{pmatrix} \psi_0 \\ \psi_0 \end{pmatrix}(x) \psi_{\kappa}(x')$$

belong to the Banach space $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L^p(\mathbb{R}^2/\Lambda)$. We view the elements of this Banach space as measurable functions f on $\mathbb{R}^2/\Lambda \times \mathbb{R}^2/\Lambda$ with norm

$$\|f\|_{q,p} = \left(\int_{\mathbb{R}^2/\Lambda} \left(\int_{\mathbb{R}^2/\Lambda} |f(x, x')|^p d^2x' \right)^{\frac{q}{p}} d^2x \right)^{1/q}.$$

The off-diagonal entries of the resolvents $\tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0)$ are convolutions with functions in the *Lorentz spaces* $L^{2,\infty}(\mathbb{R}^2/\Lambda)$ (compare with Lemma 3.20) and the diagonal entries vanish. Moreover, the sum $\sum_{\kappa \in \Lambda^*} \psi_{\kappa}(x)$ is equal to $\text{vol}^{1/2}(\mathbb{R}^2/\Lambda)$ times the Dirac's δ -function on $\mathbb{R}^2/\Lambda$.

Therefore, we have to show that for $f \in L^{2,\infty}(\mathbb{R}^2/\Lambda)$ and $g \in L^2(\mathbb{R}^2/\Lambda)$ the function $(x, x') \mapsto f(x - x')g(x')$ belongs to $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L^p(\mathbb{R}^2/\Lambda)$. The distribution function $\mu_{|f|^p}$ and the non-increasing rearrangement $(|f|^p)^*$ of the function $|f|^p$ may be expressed in terms of the distribution function μ_f and the non-increasing rearrangement f^* of f :

$$\mu_{|f|^p}(\lambda) = \mu_f(\lambda^p) \quad (|f|^p)^*(t) = (f^*(t))^p.$$

Hence the functions $|f|^p$ and $|g|^p$ belong to the *Lorentz space* $L^{\frac{2}{p},\infty}(\mathbb{R}^2/\Lambda)$ and $L^{\frac{2}{p}}(\mathbb{R}^2/\Lambda)$ and the corresponding norms are equal to

$$\| |f|^p \|_{\frac{2}{p},\infty} = \|f\|_{2,\infty}^p \quad \text{and} \quad \| |g|^p \|_{\frac{2}{p}} = \|g\|_2^p, \text{ respectively.}$$

Due to the *Generalized Young's inequality* 3.40 the $L^{\frac{q}{p}}$ -norm of the convolution of $|f|^p$ with $|g|^p$ is bounded by some constant times $\| |f|^p \|_{\frac{2}{p},\infty} \cdot \| |g|^p \|_{\frac{2}{p}}$. We remark that the relation $1/q + 1/p = 1$ implies $p/2 + p/2 = p/q + 1$. To sum up, the $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L^p(\mathbb{R}^2/\Lambda)$ -norm of $(x, x') \mapsto f(x - x')g(x')$ is bounded by the corresponding constant of the *Generalized Young's inequality* 3.40 times $\|f\|_{2,\infty} \cdot \|g\|_2$.

2. Secondly we show that for the same p, q and the same $k \in \mathcal{O}$ as before all entries of the Fourier transform of the sequence of operators

$$\tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \text{ and } \left(\mathbf{1} - \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1}$$

are bounded operators on $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L^p(\mathbb{R}^2/\Lambda)$. Due to the Hahn Decomposition Theorem [Ro-2, Chapter 11 Section 5] all L^p -functions may be decomposed into a finite complex linear combination of positive L^p -functions. Therefore, due to [DF, Section 7.3. Theorem] the multiplication with V and W define bounded operators from $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L^p(\mathbb{R}^2/\Lambda)$ to $L^{\frac{2q}{q+2}}(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_{\frac{2q}{q+2}}} L^p(\mathbb{R}^2/\Lambda)$. Multiplication by $\psi_{\pm\kappa}$ induces on these tensor products the operators $f(x, x') \mapsto g(x, x')$ with $g(x, x') = f(x, x' \pm x)$, which are obviously isometries. Finally, the entries of $\tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0)$ are convolution operators with $L^{2,\infty}$ -functions and therefore, due to the Hahn Decomposition Theorem [Ro-2, Chapter 11 Section 5], regular in the sense of [DF, Section 7.3.], which again due to [DF, Section 7.3. Theorem] induce bounded operators from $L^{\frac{2q}{q+2}}(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_{\frac{2q}{q+2}}} L^p(\mathbb{R}^2/\Lambda)$ into $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L^p(\mathbb{R}^2/\Lambda)$. This shows the claim for the first operator.

In the proof of Proposition 3.52 we mentioned that the restrictions of the sequence of operators $\left(\mathbf{1} - \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, \lambda) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1} \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, \lambda)$ to the orthogonal complement of the constant functions are the resolvents of the restrictions of $\tilde{\mathbf{D}}(\psi_{-\kappa} V, \psi_{\kappa} W, k)$ to this orthogonal complement (compare with Remark 3.53). This generalizes to the Fourier transform of these sequences of operators indexed by $\kappa \in \Lambda_{\delta}^*$ acting on the L^p -function on $\mathbb{R}^2/\Lambda$ with values in $L^q(\mathbb{R}^2/\Lambda)$. Hence the restriction of the Fourier transform of the former sequence to the orthogonal complement of $1 \otimes L^p(\mathbb{R}^2/\Lambda) \subset L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L^p(\mathbb{R}^2/\Lambda)$ is the resolvent of the restriction of the Fourier transform of the latter sequence. Now the methods of Section 2.1 carry over to these resolvents and imply that for all $V, W \in L^2(\mathbb{R}^2/\Lambda)$ there exists a $\delta > 0$, such that all entries of the Fourier transform of the sequence of operators

$$\left(\mathbf{1} - \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1} \text{ and } \left(\mathbf{1} - \tilde{\mathbf{R}}_{\text{red}}^t(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1}$$

are bounded operator on $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L^p(\mathbb{R}^2/\Lambda)$.

3. In the third step we prove that for the same $k \in \mathcal{O}$ as before the L^2 -norms of the diagonal entries and for all $1 < p < 2$ the L^p -norms of the off-diagonal entries of $\delta \mathbf{A}_{V,W,k}$ may be estimated in terms of $\|\delta V\|_2$ and $\|\delta W\|_2$. These entries of $\delta \mathbf{A}_{V,W,k}$ are integrals of pointwise products of the sequences $\psi_{-\kappa} \delta V$ and $\psi_{\kappa} \delta W$ with the entries of the sequence of functions

$$\begin{aligned} & \left(\mathbf{1} - \tilde{\mathbf{R}}_{\text{red}}^t(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1} \begin{pmatrix} \psi_0 \\ \psi_0 \end{pmatrix} = \begin{pmatrix} \psi_0 \\ \psi_0 \end{pmatrix} \\ & + \left(\mathbf{1} - \tilde{\mathbf{R}}_{\text{red}}^t(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1} \tilde{\mathbf{R}}_{\text{red}}^t(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \begin{pmatrix} \psi_0 \\ \psi_0 \end{pmatrix} \end{aligned}$$

and the entries of the sequence of functions

$$\begin{aligned} & \left(\mathbf{1} - \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1} \begin{pmatrix} \psi_0 \\ \psi_0 \end{pmatrix} = \begin{pmatrix} \psi_0 \\ \psi_0 \end{pmatrix} \\ & + \left(\mathbf{1} - \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1} \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \begin{pmatrix} \psi_0 \\ \psi_0 \end{pmatrix}, \end{aligned}$$

respectively. Due to the first two steps the Fourier transforms of the second terms of the latter two sequences of functions belong to the tensor products $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L^p(\mathbb{R}^2/\Lambda)$. Since the Fourier transform of pointwise multiplication is the convolution [R-S-II, Theorem IX.3], we shall estimate the mapping

$$\begin{aligned} L^{q_1}(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_{q_1}} L^{p_1}(\mathbb{R}^2/\Lambda) \times L^{q_2}(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_{q_2}} L^{p_2}(\mathbb{R}^2/\Lambda) & \rightarrow L^{q_3}(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_{q_3}} L^{p_3}(\mathbb{R}^2/\Lambda), \\ (f, g) & \mapsto h \text{ with } h(x, x') = \int_{\mathbb{R}^2/\Lambda} f(x, x' - x'') g(x, x'') d^2 x''. \end{aligned}$$

Due to Young's inequality [R-S-II, Section IX.4] for $1/p_1 + 1/p_2 = 1/p_3 + 1$ the convolution is a bounded bilinear mapping from the values of the L^{q_1} -function f and the values of the L^{q_2} -function g into the values of the L^{q_3} -function h . Therefore, Hölder's inequality [R-S-I, Theorem III.1 (c)] yields the estimate

$$\|h\|_{q_3, p_3} \leq \|f\|_{q_1, p_1} \|g\|_{q_2, p_2}$$

for $1/q_1 + 1/q_2 = 1/q_3$. We remark that the relations $1/p_1 + 1/p_2 = 1/p_3 + 1$ and $1/q_1 + 1/q_2 = 1/q_3$ are compatible with the relations $1/p_1 + 1/q_1 = 1/p_2 + 1/q_2 = 1/p_3 + 1/q_3 = 1$. Hence for all $1 < p < \infty$ and $1/p + 1/q = 1$ the Fourier transform of the pointwise multiplication of the entries of

$$\left(\mathbf{1} - \tilde{\mathbf{R}}_{\text{red}}^t(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1} \tilde{\mathbf{R}}_{\text{red}}^t(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \begin{pmatrix} \psi_0 \\ \psi_0 \end{pmatrix}$$

with the entries of

$$\left(\mathbf{1} - \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1} \tilde{\mathbf{R}}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \begin{pmatrix} \psi_0 \\ \psi_0 \end{pmatrix}$$

belong to $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L^p(\mathbb{R}^2/\Lambda)$. If $\delta V, \delta W \in L^p(\mathbb{R}^2/\Lambda)$, then the integrals of the products with δV and δW are bounded regular (in the sense of [DF, Section 7.3.]) maps from $L^q(\mathbb{R}^2/\Lambda)$ into $\mathbb{C}$. Since the Fourier transform of the pointwise multiplication with the sequences $\psi_{\pm\kappa}$ indexed by $\kappa \in \Lambda^*$ is the bounded operator $f \mapsto g$ with $g(x, x') = f(x, x' \pm x)$ on $L^q(\mathbb{R}^2/\Lambda) \tilde{\otimes}_{\Delta_q} L^p(\mathbb{R}^2/\Lambda)$, we finally obtain, again due to [DF, Section 7.3. Theorem], that the operator, which maps $(\delta V, \delta W)$ to the Fourier transform of the restrictions of the sequence

of operators

$$\begin{aligned} & \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, 0) \left(\mathbf{1} - \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, 0) \right)^{-1} \\ & \quad \begin{pmatrix} \psi_{-\kappa} \delta V & 0 \\ 0 & \psi_{\kappa} \delta W \end{pmatrix} \\ & \quad \left(\mathbf{1} - \tilde{R}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1} \tilde{R}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \end{aligned}$$

to the constant functions, is a bounded operator from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ into the L^p -functions on $\mathbb{R}^2/\Lambda$ with values in the 2×2 -matrices.

It remains to estimate integrals over pointwise products of the sequences $\psi_{-\kappa} V$ and $\psi_{\kappa} W$ with the entries of the sequence of functions

$$\begin{pmatrix} \psi_{-\kappa} \delta V & 0 \\ 0 & \psi_{\kappa} \delta W \end{pmatrix} \tilde{R}_{\text{red}}^t(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} \delta V \\ \psi_{\kappa} \delta W \end{pmatrix} \text{ and } \begin{pmatrix} \psi_{-\kappa} \delta V & 0 \\ 0 & \psi_{\kappa} \delta W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} \delta V \\ \psi_{\kappa} \delta W \end{pmatrix}$$

and the entries of the sequence of functions

$$\left(\mathbf{1} - \tilde{R}_{\text{red}}^t(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1} \begin{pmatrix} \psi_0 \\ \psi_0 \end{pmatrix} \text{ and } \left(\mathbf{1} - \tilde{R}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \right)^{-1} \begin{pmatrix} \psi_0 \\ \psi_0 \end{pmatrix},$$

respectively. With similar arguments (some V, W are replaced by $\delta V, \delta W$) we decompose the corresponding operators again into sums of two operators, whose second terms yield bounded operators from $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ into the L^2 -functions on $\mathbb{R}^2/\Lambda$ with values in the 2×2 -matrices.

The images of the remaining terms are the Fourier transforms of the restrictions of the sequences

$$\begin{aligned} & \begin{pmatrix} \psi_{-\kappa} \delta V & 0 \\ 0 & \psi_{\kappa} \delta W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \\ & \begin{pmatrix} \psi_{-\kappa} V & 0 \\ 0 & \psi_{\kappa} W \end{pmatrix} \tilde{R}_{\text{red}}(0, 0, k, 0) \begin{pmatrix} \psi_{-\kappa} \delta V & 0 \\ 0 & \psi_{\kappa} \delta W \end{pmatrix} \end{aligned}$$

to the constant functions, respectively. These correspond to off-diagonal matrices and induce bounded operators from $L^p(\mathbb{R}^2/\Lambda) \times L^p(\mathbb{R}^2/\Lambda)$ into the L^p -functions on $\mathbb{R}^2/\Lambda$ with values in the 2×2 -off-diagonal matrices for all $1 < p < 2$.

4. By definition of $\hat{v}(V, W, \cdot)$, $\hat{w}(V, W, \cdot)$ the variation of the off-diagonal entries of $\mathbf{A}_{V, W, k}$ yield variations of the sequences k_{κ} and contribute indirectly through the partial derivatives of the diagonal parts of $\mathbf{A}_{V, W, k}$ with respect to k to the variations of

$$v = \sum_{\kappa \in \Lambda_{\delta}^*} \hat{v}(V, W, \kappa) \quad w = \sum_{\kappa \in \Lambda_{\delta}^*} \hat{w}(V, W, \kappa)$$

The considerations concerning these partial derivatives in the proof of Proposition 3.52 combined with the methods of the first three steps imply that for suitable $k \in \mathcal{O}$ all L^q -norms

with $1 < q < \infty$ of these partial derivatives are bounded (in terms of $\|V\|_2$ and $\|W\|_2$). Since the Fourier transform of pointwise multiplication is the convolution [R-S-II, Theorem IX.3], the variations of the off-diagonal terms of $A_{V,W,k}$ contribute to the variations of v, w over the convolutions with the partial derivatives of the diagonal parts of $A_{V,W,k}$ with respect to k . Due to Young's inequality [R-S-II, Section IX.4], for all $q < \infty$ the L^q -norms of these contributions are finite and the lemma is proven. **q.e.d.**

Continuation of the proof of Theorem 3.57. The proof of this lemma implies that for all $V, W \in L^2(\mathbb{R}^2/\Lambda)$ there exists a $\delta > 0$ such that the mapping

$$(V, W) \mapsto (v, w) \quad \text{with} \quad \begin{pmatrix} v \\ w \end{pmatrix} = \sum_{\kappa \in \Lambda_\delta^*} \begin{pmatrix} \psi_\kappa \hat{v}(V, W, \kappa) \\ \psi_\kappa \hat{w}(V, W, \kappa) \end{pmatrix}$$

is locally a holomorphic mapping from an open neighbourhood of $(V, W) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ onto an open neighbourhood of $(v, w) \in L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$. In fact, in order to estimate the L^2 -norms of v and w we integrate the derivative along a path to the zero potentials with $v = w = 0$. Furthermore, for small δ locally the restrictions of the derivatives of this mapping to the inverse Fourier transform of $\ell_2(\Lambda_\delta^*) \times \ell_2(\Lambda_\delta^*)$ are bounded invertible operators. Hence the implicit function theorem [R-S-I, Supplementary material V.5 Theorem S.11] implies that this mapping is open. In particular, the *finite type potentials* are dense in $L^2(\mathbb{R}^2/\Lambda)$. We remark that for potentials of the form $(U, \bar{U})$ the positivity of the moduli combined with the asymptotic analysis of Proposition 3.52 implies that the L^2 -norms of $u = \sum_{\kappa \in \Lambda_\delta^*} \psi_\kappa \hat{u}(U, \kappa)$ are bounded in terms of $\|U\|_2$. Moreover, the methods of Lemma 3.58

can be used in order to estimate directly the L^2 -norms of v and w in terms of $\|V\|_2$ and $\|W\|_2$.

Due to the estimate $\left\| A(\psi_{-\kappa} V, \psi_\kappa W, k) - \begin{pmatrix} \hat{v}(V, W, \kappa) & 0 \\ 0 & \hat{w}(V, W, -\kappa) \end{pmatrix} \right\| \leq \varepsilon \|k - k_\kappa\|$ in the proof of Proposition 3.52 these arguments extend to the moduli t and show that the mapping $(V, W) \mapsto t$ is a holomorphic open mapping from $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ into $\ell_1(\Lambda_\delta^*)$ with suitable δ depending on V and W .

4. Now we shall define locally all moduli $t(\kappa)$. We shall complete the cycles of the handles index by $\kappa \in \Lambda_\delta^*$ to a set of generators of a Lagrangian subgroup of the first homology group of the *complex Fermi curves* with respect to the intersection form. In addition we assume that the integrals of dk along these cycles vanish. Moreover, in neighbourhoods of *complex Fermi curves* with singularities, we assume that a small neighbourhood of all deformed singularities contain as many cycles, as the *Local contribution to the arithmetic genus* 4.3. The cycles of the handles described in Theorem 2.12 and Proposition 3.52 have these properties. Due to Proposition 3.52 the elements of this basis may be indexed by $\kappa \in \Lambda^*$. For *complex Fermi curves* of the form $\mathcal{F}(U, \bar{U})$ we complete the *horizontal cuts* indexed by $\kappa \in \Lambda_\delta^*$ to a set of generators of those cycles in the first homology group, which are invariant under the corresponding action of η . Consequently we define all moduli as

$$t(\kappa) = \pi \oint_{\text{cycle with index } \kappa} k_1 dk_2.$$

Since the integral of dk along these cycles vanish, the corresponding moduli do not depend on the choice of the branch. Moreover, the moduli of *complex Fermi curves* of the form $\mathcal{F}(U, \bar{U})$ are real. Due to Proposition 3.7, these moduli define locally coordinates of the manifold $\mathcal{M}_{g,\Lambda}$.

We shall now show that locally these moduli completely determine the *complex Fermi curves*; i. e. if two potentials have locally the same moduli, then they have the same *complex Fermi curve*. Let $\mathcal{U}$ be a small open subset of $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$, on which a complete choice of moduli is defined. Two pairs of potentials in $\mathcal{U}$ whose moduli coincide, are limits of two convergent sequences in $\mathcal{U}$, whose *complex Fermi curves* have finite *geometric genus*, and whose sequences of moduli coincide. Therefore, due to the continuity of the map $(V, W) \mapsto \mathcal{F}(V, W)$, it suffices to show that the moduli completely determine the *complex Fermi curves* of finite *geometric genus*. This follows from Proposition 3.7.

To sum up, these moduli define locally holomorphic coordinates of $\mathcal{M}_\Lambda$, which take values in $\ell_1(\Lambda^*)$. In particular, $\mathcal{M}_\Lambda$ is a $\ell_1(\Lambda^*)$ -manifold.

5. We conclude that all *complex Fermi curves* in $\mathcal{M}_\Lambda$, whose normalization is invariant under η without fixed points, belongs to $\bar{\mathcal{M}}_{\Lambda,\eta,\mathcal{W}}$ with $\mathcal{W} = 4 \int_{\mathbb{R}^2/\Lambda} V(x)W(x)d^2x$. This follows

from Lemma 2.27 and the fact that the potentials, whose *complex Fermi curves* have finite *geometric genus*, are dense in $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$. Consequently the moduli space $\mathcal{M}_{\Lambda,\eta}$ is a closed real subspace of $\mathcal{M}_\Lambda$ (with respect to the $\ell_1(\Lambda^*)$ -topology). Moreover, the moduli with indexes in Λ_δ^* are non-negative. Therefore, this moduli space has an asymptotic $\ell_1^+(\Lambda_\delta^*)$ -structure.

We may extend this local description of the moduli space to neighbourhoods of *complex Fermi curves* of *Finite rank Perturbations* 3.22 of a limit of sequences of potentials in $L^2(\mathbb{R}^2/\Lambda)$. In fact, to these sequences we may add potentials, whose support is contained in the complement of the unions $\mathcal{S}_{D,\varepsilon}$ of small balls around the singular points in $\mathbb{R}^2/\Lambda$ (compare with Theorem 3.48 and Remark 3.51). Since the union

$$\bigcup_{\varepsilon>0} \{ \delta U \in L^2(\mathbb{R}^2/\Lambda) \mid \text{support}(\delta U) \cap \mathcal{S}_{D,\varepsilon} = \emptyset \}$$

is dense in $L^2(\mathbb{R}^2/\Lambda)$ all elements of $\bar{\mathcal{M}}_{\Lambda,\eta}$ are contained in an open set, which is mapped by the restricted moduli onto an open set of $\ell_1^+(\Lambda_\delta^*)$. **q.e.d.**

In particular, we proved

Corollary 3.59. All complex Fermi curves $\mathcal{F}(V, W)$ with potentials $V, W \in L^2(\mathbb{R}^2/\Lambda)$, whose normalizations are invariant under the involution η but without fixed points, belong to $\bar{\mathcal{M}}_{\Lambda,\eta,\mathcal{W}}$ with $\mathcal{W} = 4 \int_{\mathbb{R}^2/\Lambda} V(x)W(x)d^2x$. **q.e.d.**

Remark 3.60. The boundedness of the first integral of the complex Fermi curves yields additional restrictions of *Finite rank Perturbations* 3.22 of $R(U, \bar{U}, k, 0)$. In fact, it is quite easy to see that the complex Fermi curves of the *Finite rank Perturbations* 3.22 of the free resolvent have disconnected normalization, if the first integral is bounded. However, there

exists such rank one Perturbations, whose normalizations are connected. The corresponding complex Fermi curves might correspond to potentials $U \in L^{2,\infty}(\mathbb{R}^2/\Lambda)$.

Due to the estimates established in the proof of Proposition 3.52, the derivatives $\delta t(\kappa)/\delta U$ of the moduli are locally analytic vector fields on $L^2(\mathbb{R}^2/\Lambda)$. For all compact Riemann surfaces Y the long exact cohomology sequence

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{C} \xrightarrow{\exp} \mathbb{C}^* \rightarrow H^1(Y, \mathbb{Z}) \rightarrow H^1(Y, \mathcal{O}) \xrightarrow{\exp} H^1(Y, \mathcal{O}^*) \rightarrow H^2(Y, \mathbb{Z}) \rightarrow 0$$

corresponding to the exact sequence

$$0 \rightarrow \mathbb{Z} \hookrightarrow \mathcal{O} \xrightarrow{\exp} \mathcal{O}^* \rightarrow 1$$

of sheaves shows that the first cohomology group $H^1(Y, \mathbb{Z})$ considered as a subspace of the Lie algebra of the Picard group is exactly the kernel of the exponential mapping of the Picard group. By definition of the moduli $t(\kappa)$ and the one-forms $\Omega_{V,W}$ (compare with Lemma 3.1) the variations $\delta t(\kappa)$ with respect to $(\delta V, \delta W)$ is equal to the integral $\Omega_{V,W}(\delta V, \delta W)$ over the unique cycle of the handle with index κ . For *finite type potentials* Lemma 3.2 implies therefore that the Hamiltonian vector fields of the moduli $t(\kappa)$ induce periodic flows with period 1. Since the *finite type potentials* are dense, the same is true for all potentials. Since in the limit $g(\kappa, \kappa) \rightarrow \infty$ the norm of these vector fields converges to zero, for all $U \in L^2(\mathbb{R}^2/\Lambda)$ and all $\varepsilon > 0$ there exists some $\delta > 0$ such that the flows corresponding to the Hamiltonian vector fields of all moduli $t(\kappa)$ with $\kappa \in \Lambda_\delta^*$ maps U for all times into $B(U, \varepsilon)$. In particular, for all $U \in L^2(\mathbb{R}^2/\Lambda)$ and all small $\varepsilon > 0$ the isospectral set of U contains an infinite-dimensional torus in $B(U, \varepsilon)$, which is homeomorphic to $(\mathbb{R}/\mathbb{Z})^{\{\kappa \in \Lambda_\delta^* | t(\kappa) > 0\}}$. Also the corresponding action is induced by the Hamiltonian vector fields of the moduli with indexes $\kappa \in \Lambda_\delta^*$.

4 The Generalized Willmore functional

4.1 The Weak Singularity condition

Not all real potentials U correspond to some immersion of a torus into $\mathbb{R}^3$. In this section we want to find some characteristic properties of all *complex Fermi curves*, which correspond to immersions. All *complex Fermi curves*, which correspond to immersions have, roughly speaking, some higher order singularity:

Lemma 4.1. *Assume that the Dirac operator $\left(\begin{smallmatrix} U & \partial \\ -\bar{\partial} & U \end{smallmatrix} \right)$ with real potential U has an eigenfunction $\chi(x) = \begin{pmatrix} \chi_1(x) \\ \chi_2(x) \end{pmatrix}$ with eigenvalue zero, which obeys $\chi(x + \gamma) = \pm \chi(x)$ for all $x \in \mathbb{R}^2$ and all periods $\gamma \in \Lambda$ and the Equivalent form of the Periodicity condition 1.2. Then there exists some $\kappa \in \Lambda^*$, whose equivalence class $[\kappa/2]$ belongs to the complex Fermi curve $\mathcal{F}(U, U)/\Lambda^*$. Moreover, the preimage of this point in the normalization of $\mathcal{F}(U, U)/\Lambda^*$ contains either at least four elements or two elements, which are zeroes of both differentials $d\hat{p}$ and $d\check{p}$.*

Proof. The transformation of the eigenfunction under shifts of the periods implies that there exists some element κ of the dual lattice, such that $[\kappa/2]$ belongs to $\mathcal{F}(U, U)/\Lambda^*$. Due to (ii) of Corollary 2.17 the preimage of this element $[\kappa/2]$ in the normalization of $\mathcal{F}(U, U)/\Lambda^*$ has an even number of elements. Now let us assume that the preimage has two elements y and y' . Due to Corollary 2.17 y' is equal to $\rho(\sigma(y)) = \sigma(\rho(y))$. Thus the differentials dk_1 and dk_2 have zeroes either at both points y and y' or at none of them. Let us assume on the contrary to the statement of the lemma that there exists some period $\hat{\gamma}$, such that the differential $dg(\hat{\gamma}, k)$ is not equal to zero at y and y' . Thus the projection $\check{P}_{\hat{\gamma}}([\kappa/2])$ defined in (iii) of Lemma 2.23 has rank two. Due to (v) of Lemma 2.23 the eigenfunction χ , which fulfills the conditions of the theorem, is invariant under $\check{P}_{\hat{\gamma}}([\kappa/2])$. Hence, this eigenfunction χ and $J\bar{\chi}$ span the eigenspace of this projection. The normalization of $\mathcal{F}(U, U)/\Lambda^*$ is a smooth Riemann surface. Hence the pullback of the eigenfunctions ψ and ϕ to the normalization can be chosen to be locally holomorphic without any zeroes. Since one of the involutions σ and ρ permutes y and y' and the other involution leaves these points invariant, the span of $\phi(y)$ and $\phi(y')$ is equal to the span of $J\psi(y)$ and $J\psi(y')$ and to the span of $\bar{\psi}(y)$ and $\bar{\psi}(y')$. Thus the *Equivalent form of the Periodicity condition* 1.2 implies that $P_{\hat{\gamma}}([\kappa/2])\psi(y)$ and $P_{\hat{\gamma}}([\kappa/2])\psi(y')$ are equal to zero. This is a contradiction to (vi) of Lemma 2.23. Thus both differentials dk_1 and dk_2 are zero at these two elements y and y' . **q.e.d.**

For any singularity of the *complex Fermi curve* we may distinguish between multiple points and cusps:

Multiple point: A point of the normalization of the *complex Fermi curve* in the preimage of some singularity is called a *multiple point*, if there exists some $\gamma \in \Lambda$, whose differential $dg(\gamma, k)$ does not vanish at this point.

Cusp: A point of the normalization of the *complex Fermi curve* in the preimage of some singularity is called *cusp*, if it is a zero of both differentials dk_1 and dk_2 .

Now we may paraphrase the condition of Lemma 4.1.

Weak Singularity condition 4.2. (See also 1.5.) *There exists some $\kappa \in \Lambda^*$, such that $[\kappa/2]$ belongs to the complex Fermi curve. Moreover, the preimage of this point in the normalization of $\mathcal{F}/\Lambda^*$ contains either at least four multiple points, or at least two cusps.*

If we decompose the preimage of $[\kappa/2]$ in the normalization of $\mathcal{F}(U, U)/\Lambda^*$ into orbits of the two involutions σ and ρ three possible orbits can occur.

Orbit of type 1: Two fixed points of ρ , which are permuted by σ .

Orbit of type 2: Two fixed points of σ , which are permuted by ρ .

Orbit of type 3: Four points and both involutions σ and ρ have no fixed point.

In particular, the *Weak Singularity condition* 4.2 permits the following situations:

Multiple points O1/O1: Two *Orbits of type 1* containing *multiple points* (8).

Multiple points O1/O2: One *Orbit of type 1* containing *multiple points* and one *Orbit of type 2* containing *multiple points* (7).

Multiple points O2/O2: Two *Orbits of type 2* containing *multiple points* (6).

Multiple points O3: One *Orbit of type 3* containing *multiple points* (8).

Cusps O1: One *Orbit of type 1* containing *cusps* (8).

Cusps O2: One *Orbit of type 2* containing *cusps* (23).

Cusps O3: One *Orbit of type 3* containing *cusps* (37).

These situations differ in the possible orders of the singularity. The numbers in the round brackets are the lowest *Local contribution to the arithmetic genus* of the *complex Fermi curves*. The last two cases yield very high contributions and they are exceptional situations.

Local contribution to the arithmetic genus 4.3. *If $\mathcal{O}_{Y,\text{normal}}$ is the normalization sheaf (i. e. the direct image of the structure sheaf of the normalization under the normalization map [Gr-Re, Chapter 6. §4.] of a compact pure one-dimensional complex space Y , then the long exact cohomology sequence [Fo, §15.] corresponding to the exact sequence of sheaves*

$$0 \rightarrow \mathcal{O}_Y \rightarrow \mathcal{O}_{Y,\text{normal}} \rightarrow \mathcal{O}_{Y,\text{normal}}/\mathcal{O}_Y \rightarrow 0$$

shows that the arithmetic genus of Y , which is equal to $\dim H^1(Y, \mathcal{O}_Y)$, is equal to the geometric genus, which is equal to $\dim H^1(Y, \mathcal{O}_{Y,\text{normal}})$, plus $\dim H^0(Y, \mathcal{O}_{Y,\text{normal}}/\mathcal{O}_Y)$. The support of this sheaf is the singular locus of Y [Gr-Re, Chapter 6. §5.3.], and the space of global sections is equal to the direct sum of all non-trivial stalks. The dimension of these stalks is the local contribution to the arithmetic genus at the corresponding singularities [Se, Chapter IV §1.4. and §2.7.].

The lemma shows that the *complex Fermi curve* of some potential, which corresponds to some immersion, obeys some condition. In Section 5.1 we will see that this condition on the *complex Fermi curve* of some real potential implies neither that this potential is a *Weierstraß potential* 1.3 nor that the *complex Fermi curve* is a *Weierstraß curve* 1.4. In fact, in Section 5.1 we shall formulate the *Singularity condition* 5.3, which is equivalent to the condition that the *complex Fermi curve* is a *Weierstraß curve* 1.4. Moreover, we will describe the subset of the corresponding *isospectral sets*, which contain the *Weierstraß potentials* 1.3. For *Complex Fermi curves*, which are invariant under the involution σ , however, the *Weak Singularity condition* 4.2, is equivalent to the *Quaternionic Singularity condition* 1.6 combined with the condition on the singularity being invariant under σ . In particular, the *Weak Singularity condition* 4.2 is stronger than the *Quaternionic Singularity condition* 1.6. In some sense the *Weak Singularity condition* 4.2 seems to be the most convenient combination of the advantages of immersion into $\mathbb{R}^4$ (corresponding to the *Quaternionic Singularity condition* 1.6), which yield stronger conditions on relative minimizers, and the advantages of immersion into $\mathbb{R}^3$, whose *complex Fermi curves* have an additional involution σ . In fact, in

contrast to the *Singularity condition* 5.3 the *Weak Singularity condition* 4.2 allows to deform *Orbits of type 1* and *Orbits of type 3* into *Orbits of type 2*. This deformation is essential for the proof that the relative minimizers have dividing real parts. Moreover, in Section 4.3 we shall see that this property is almost as restrictive as the *real- σ -hyperellipticity*, which characterizes minimal tori in S^3 [Hi].

In Section 3.5 we have seen that the moduli space has a compactification containing *complex Fermi curves* of *Finite rank perturbations* 3.22. Hence it is natural to extend the Willmore functional not only to all *Generalized Weierstraß potentials* 1.7, but also to all *Generalized Weierstraß curves*.

Generalized Weierstraß curves 4.4. Complex Fermi curves in $\bar{\mathcal{M}}_{\Lambda, \eta, \sigma}$, which obey the Weak Singularity condition 4.2 are called Generalized Weierstraß curves.

Due to Lemma 3.10, for all $\mathcal{W} > 0$ the spaces $\bar{\mathcal{M}}_{\Lambda, \eta, \sigma, \mathcal{W}}$ are compact with respect to the subspace topology of the space of compact subsets of $\mathbb{C}^2$. Since the subset consisting of *Generalized Weierstraß curves* 4.4 is closed the restriction of the *first integral* to this subspace is lower semi-continuous. This restriction is called

Generalized Willmore functional 4.5.

In Theorem 3.57 we have seen that the compactified moduli space has an $\ell_1^+(\Lambda^*)$ -structure. Consequently, this compactified moduli space is very similar to a manifold and some specific regular one-forms correspond to tangent vectors.

4.2 Relative minimizers of the *Generalized Willmore functional*

In this section we investigate relative minimizers of the *Generalized Willmore functional* 4.5, i. e. minimizers of the restrictions to open sets [Str]. The main result is that all relative minimizers have

Dividing real parts. A complex Fermi curve $\mathcal{F} \in \bar{\mathcal{M}}_{\Lambda, \eta, \sigma}$ has a dividing real part, if the relative complement of the real part (i. e. the set of fixed points of ρ) in the normalization of $\mathcal{F}/\Lambda^*$ has two connected components [B-C-R, 11.6]

Remark 4.6. Due to Proposition 3.52 (compare with Corollary 3.54), all complex Fermi curves with dividing real parts have finite geometric genus. In Corollary 5.10 we shall show with the arguments of Proposition 3.5 that all Constrained Willmore tori 5.8 have finite geometric genus. Unfortunately these arguments do not apply to the Generalized Willmore functional 4.5. More precisely, in order to apply Proposition 3.5 to the Generalized Willmore functional 4.5, we have to prove that all variations of Generalized Weierstraß potentials 1.7 may be characterized by the finite-order vanishing of the corresponding one-form at the preimage in the normalization of the singularity described in the Weak Singularity condition 4.2 analogous to Proposition 5.5. But the subvariety of Generalized Weierstraß potentials 1.7 may have singularities, whose complex Fermi curves may be of infinite geometric genus. More

precisely, Lemma 3.2 implies that the corresponding complex Fermi curves have global meromorphic functions, but they may take the same values at $\infty^\pm$. Therefore, the corresponding complex Fermi curves can have infinite geometric genus. Varieties with singularities have different notions of tangent spaces [Wh, Chapter 7]. In particular, differentiable paths contained in the variety correspond to elements of the tangent cone [Wh, Chapter 7 Section 2 and Theorem 3C]. Hence in order to carry over the methods of Corollary 5.10 it would suffice to characterize the elements of the tangent cone by a finite-order vanishing of the corresponding one-forms at the preimage in the normalization of the singularity mentioned in the Weak Singularity condition 4.2. We did not succeed in proving this. For this reason the characterization of relative minimizers of the Generalized Willmore functional 4.5 in this Section has to apply to Generalized Weierstraß curves 4.4 of infinite geometric genus. For this purpose we proved Theorem 3.57, which will turn out to be the essential tool of the extension of the deformation theory devolved in Section 3.2 to complex Fermi curves of infinite geometric genus.

Of special importance is the subclass of

Real- σ -hyperelliptic complex Fermi curves. In this case the normalization of the quotient of $\mathcal{F}/\Lambda^*$ modulo σ is biholomorphic to $\mathbb{P}^1$ with two points removed. Moreover, on this quotient the involutions η and ρ induce the unique involution of $\mathbb{P}^1$ with one real cycle.

Remark 4.7. A real hyperelliptic Riemann surface is determined by a collection of branch points on $\mathbb{P}^1$, which are invariant under some anti-holomorphic involution of $\mathbb{P}^1$. Since $\mathbb{P}^1$ has two equivalence classes of anti-holomorphic involutions, these real hyperelliptic Riemann surfaces also splits into two classes. In the first case the anti-holomorphic involution on $\mathbb{P}^1$ has one real cycle, and in the second case the real part is empty.

Due to Corollary 2.13 the normalization is not necessarily connected. If the real part shrinks to one or several singular points we obtain as a another subclass the *complex Fermi curves* with

Disconnected normalization.

Since the fixed points of σ are the branch points of the natural two-sheeted covering of the complex Fermi curve over the quotient modulo σ , due the Riemann–Hurwitz formula [Fo, 17.14] their number has to be even. Therefore, each connected component of a disconnected normalization has besides the fixed point at $\infty^\pm$ in addition an odd number of fixed points of σ . Since these fixed points of σ may be deformed into real cycles, all complex Fermi curves in $\mathcal{M}_{\Lambda, \eta, \sigma}$ with disconnected normalization are indeed limits of complex Fermi curves with dividing real parts.

Remark 4.8. Both anti-holomorphic involutions ρ and η induce on the quotient modulo σ the same anti-holomorphic involution, whose fixed points are the real part of the quotient modulo σ . Since η does not have any fixed points on the normalization, the preimage of the real part of the normalization of the quotient modulo σ in normalization is the real part of the normalization. Therefore a complex Fermi curve has a dividing real part if and only if the normalization of the quotient modulo the involution σ has a dividing real part.

Theorem 4.9. *The relative minimizers of the Generalized Willmore functional 4.5 belong to one of the following classes:*

Minimizers with dividing real parts: *The normalization of the quotient of $\mathcal{F}/\Lambda^*$ modulo the involution σ is a real hyperelliptic Riemann surface, whose hyperelliptic involution does not have any real fixed points. More precisely, the real part is a two-sheeted covering of the real cycle of $\mathbb{P}^1$, which is the real part of the quotient modulo the hyperelliptic involution. Moreover, the preimage in the normalization of the singularity described in the Weak Singularity condition 4.2 contains Multiple points O1/O1, and no other orbit. Finally, the function $d\check{p}/d\hat{p}$ takes different values on the two Orbits of type 1, and the exterior derivative of this function does not vanish on the two Orbits of type 1.*

Real- σ -hyperelliptic minimizers: *The minimizers are real- σ -hyperelliptic and the preimage in the normalization of the singularity described in the Weak Singularity condition 4.2 contains either Multiple points O1/O1, or Multiple points O1/O2, or Cusps O1, and no other orbit.*

Minimizers with disconnected normalization: *The minimizers have disconnected normalizations and the preimage in the normalization of the singularity described in the Weak Singularity condition 4.2 contains either Multiple points O2/O2, or Multiple points O3, or Cusps O2 or Cusps O3.*

Proof. In a first step we show that all relative minimizers in $\bigcup_{g \in \mathbb{N}} \bar{\mathcal{M}}_{g,\Lambda,\eta,\sigma}$ belong to one of the three cases described in the Theorem. In a second step we extend our arguments to all relative minimizers in $\bar{\mathcal{M}}_{\Lambda,\eta,\sigma}$.

1. Local minimizers of finite *geometric genus*. Let us describe the general construction of local flows on the space $\bigcup_{g \in \mathbb{N}} \mathcal{M}_{g,\Lambda,\eta,\sigma}$. Due to Proposition 3.7 the complex moduli space

$\mathcal{M}_{g,\Lambda}$ of algebraic curves with arithmetic genus equal to g is a $(g+1)$ -dimensional complex manifold. Moreover, at each element Y of this manifold the tangent space is isomorphic to the space of regular forms of the algebraic curve $\tilde{Y}$, which is obtained from Y by identifying the two marked points ∞^- and ∞^+ to an ordinary double point. The *Generalized Weierstraß curves* 4.4 are elements of the moduli space $\bar{\mathcal{M}}_{g,\Lambda,\eta,\sigma}$, and may be characterized as those elements of the complex moduli space, which are invariant under the involutions $\eta : k \mapsto -\bar{k}$ and $\sigma : k \mapsto -k$, and on whose normalizations the first involution does not have fixed points. Due to Lemma 2.15 these involutions induce involutions on the moduli space $\mathcal{M}_{g,\Lambda}$, and those *complex Fermi curves*, which are invariant under the former involutions, are the fixed points of the latter involutions. For all *complex Fermi curves* in $\bar{\mathcal{M}}_{g,\Lambda,\eta,\sigma}$ we may find some open neighbourhood of the corresponding element of $\mathcal{M}_{g,\Lambda}$, which is invariant under these involutions of the complex moduli space. Due to the proof of Proposition 3.7 the regular forms of $\tilde{Y}$ form a holomorphic vector bundle on the complex moduli space, which is isomorphic to the tangent bundle. Hence any regular form ω fulfilling the equations $\eta^*\omega = \bar{\omega}$ and $\sigma^*\omega = \omega$ on Y may be extended to a holomorphic vector field on some neighbourhood of Y in the complex moduli space, which fulfills the corresponding extensions of these equations to this

invariant neighbourhood. The flows corresponding to these vector fields leave invariant the fixed point set of the two involutions on the moduli space. With this method we construct local flows on the fixed point set of both involutions of the complex moduli space. In a second step we investigate the fixed points of the involutions η on the corresponding invariant *complex Fermi curves*. It may happen that in spite of η on Y having no fixed point, this is not true for some neighbourhood of Y in the fixed point set of both involutions on the complex moduli space. We will see that for some one-dimensional families $\mathcal{F}_t$ of *complex Fermi curves* being invariant under η , this involution has fixed points on the normalizations of $\mathcal{F}_t$ for $t < 0$ and no fixed points for $t \geq 0$. In these situations we call those regular forms fulfilling $\eta^*\omega = \bar{\omega}$ and $\sigma^*\omega = \omega$ *admissible*, whose flows corresponding to positive times do not leave the subset of *complex Fermi curves* with no fixed points of η on the corresponding normalizations. The set of these *admissible* regular forms considered as elements of the tangent space of the moduli space (Proposition 3.7 and Theorem 3.57) is called real tangent semi cone along the subvariety of *Generalized Weierstraß curves* 4.4 [Wh, Chapter 7 Section 2 and Theorem 3C].

Let us apply these considerations to those singularities of Y , which are different from the singularity described in the *Weak Singularity condition* 4.2. The former singularities may be removed without leaving the subset of *Generalized Weierstraß curves* 4.4. We shall distinguish between the following cases:

Ordinary double points of the form $(y, \eta(y))$. Without loss of generality we may assume that both components of k are zero at these two points, and that the differential $d\hat{p}$ does not vanish at these double points. The Taylor series of the function $\check{p}(\hat{p})$ at these two points has to look like $\check{p} = a_1\hat{p} + a_2\hat{p}^2 + \dots$ and $\check{p} = \bar{a}_1\hat{p} - \bar{a}_2\hat{p}^2 + \dots$, respectively. Therefore, the *complex Fermi curve* is locally given by the equation $R(\hat{p}, \check{p}) = 0$. More precisely, $R(\hat{p}, \check{p}) =$

$$(\check{p})^2 - \check{p}((a_1 + \bar{a}_1)\hat{p} + (a_2 - \bar{a}_2)\hat{p}^2 + \dots) + a_1\bar{a}_1\hat{p}^2 + (a_2\bar{a}_1 - a_1\bar{a}_2)\hat{p}^3 + \dots = 0.$$

Let us assume in addition that the imaginary part of a_1 does not vanish. Otherwise the function $d\check{p}/d\hat{p}$ takes the same values at both elements of the double point, and we have some higher order singularity. The form

$$\omega = \frac{1 + \mathbf{O}(\hat{p}) + \mathbf{O}(\check{p})}{\partial R(\hat{p}, \check{p})/\partial \check{p}} d\hat{p}$$

is regular and the deformation of any extension of this regular form to a holomorphic vector field of the moduli space is locally given by the equation

$$R(\hat{p}, \check{p}) + t(1 + \mathbf{O}(\hat{p}) + \mathbf{O}(\check{p})) + \mathbf{O}(t^2) = 0.$$

The vector field may be chosen to be invariant under the involutions σ and ρ . In this case the whole family corresponding to real t is invariant under these involutions. Fixed points of η correspond in our coordinates to elements, whose coordinates $\hat{p}$ and $\check{p}$ are imaginary. In general the real part of k has to take values in $\Lambda^*/2$ at these double points of the form $(y, \eta(y))$. For simplicity we assumed that k vanishes at the double

point. Now it is easy to see that the involution η does not have fixed points on the part of the family corresponding to $t \leq 0$. We conclude that any *admissible* form may have poles of first-order at ordinary double points of the form $(y, \eta(y))$, but the residue has to be purely imaginary and the sign of the imaginary part has to be the same as the sign of the imaginary part of $d\check{p}/d\hat{p}$ at these points.

Higher order singularities of the form $(y, \eta(y))$. There are two possibilities: either the function $d\check{p}/d\hat{p}$ takes the same value at these two points, or both differentials $d\hat{p}$ and $d\check{p}$ vanish at these two points. Let us first consider the case that for some choice of generators of Λ the differential $d\hat{p}$ does not vanish at both points of $(y, \eta(y))$. In this case the first Taylor coefficient a_1 introduced above is real. If we add to $\check{p}(\hat{p})$ some polynomial of $\hat{p}$, which takes for imaginary-valued $\hat{p}$ imaginary values, we may achieve that the two Taylor series are given by $\check{p} = a_l \hat{p}^l + a_{l+1} \hat{p}^{l+1} + \dots$ and $\check{p} = -\bar{a}_l (-\hat{p})^l - a_{l+1} (-\hat{p})^{l+1} + \dots$, where l is the lowest index with $a_l + (-1)^l \bar{a}_l \neq 0$. Now it is easy to see that the singular part of an *admissible* regular form at this singularity has to fulfill some positivity condition. The same is true in the other cases of higher singularities.

Singularities, which are disjoint from their image under η . In these cases any regular form ω fulfilling $\sigma^* \omega = \omega$ and $\eta^* \omega = \bar{\omega}$ is *admissible*.

If a *Generalized Weierstraß curve* 4.4 admits a local flow decreasing the *first integral*, it cannot correspond to a relative minimizer of the *Generalized Willmore functional* 4.5. Therefore, we shall try to characterize all those situations, where such flows do not exist. The main task is to determine all vector fields, whose flows preserves the *Weak singularity condition* 4.2. Again we use the local description of some one-sheeted coverings of the *complex Fermi curves* by an equation $R(\hat{p}, \check{p}) = 0$. In order to simplify notations, we will always assume that the function k (and therefore also the components $\hat{p}$ and $\check{p}$) is zero at the singularity described in the *Weak singularity condition* 4.2. Since the elements of the form $[\kappa/2]$ are fixed points of the two involutions η and σ , this simplification is compatible with these two involutions. Moreover, in some situations it will be more convenient to choose the second component $\check{p}$ to be a complex component $\check{p} = g(\check{\mu}, k)$ of k , like at the end of Section 2.4. For the construction of the complex manifold of deformations of *complex Fermi curves* in Proposition 3.7 we covered Y by some open smooth set and small neighbourhoods of the singularities. In order to apply this construction to deformations of *complex Fermi curves* fulfilling the *Weak singularity condition* 4.2 we modify the representation of Y in a small neighbourhood of this singularity. The function $\hat{p}$ depends on the choice of the generators of the lattice. Obviously there exists some choice, such that $d\hat{p}$ has no zero at all *multiple points* in the preimage of $[\kappa/2]$ described in the *Weak singularity condition* 4.2. At each *cusp* of this preimage, the differentials of almost all components of k have a zero of some order $d - 1 \in \mathbb{N}$. The differential of a specific complex linear combination of k_1 and k_2 has a zero of higher-order $m - 1$. Moreover, the natural numbers $d < m$ have to be co-prime, because otherwise the preimage of the normalization of this point contains several elements. Obviously we may also assume that for all *cusps* in the preimage of $[\kappa/2]$ the differential $d\hat{p}$ has a zero of minimal-order, compared with all other components of k . This implies that the

function $\check{p}/\hat{p}$ extends to a holomorphic function on some neighbourhood of the preimage of $[\kappa/2]$. Now we describe Y in some small neighbourhood of this preimage, by some equation of the form $R_\sigma(\hat{p}^2, \check{p}/\hat{p}) = 0$. Again we may choose this function to be a polynomial with respect to $\check{p}/\hat{p}$, whose degree is equal to the local number of sheets of the covering map $\hat{p}$, and whose highest coefficient is equal to 1. Due to this choice the other coefficients are unique holomorphic functions depending on $\hat{p}$. If the small neighbourhoods are invariant under σ , this equation depends only on $\hat{p}^2$. In fact, in this case both functions $\check{p}/\hat{p}$ and $\hat{p}^2$ are invariant under σ , and this equation yields a local description of Y_σ . If we consider $\check{p}$ as a function of $\hat{p}$ and the moduli (as we did in Proposition 3.7) the regular form ω describing a local deformation is equal to

$$\omega = \frac{\partial \check{p}}{\partial t} d\hat{p} = 1/2 \frac{\partial(\check{p}/\hat{p})}{\partial t} d(\hat{p}^2) = -\frac{\partial R_\sigma(\hat{p}^2, \check{p}/\hat{p}, t)/\partial t}{2\partial R_\sigma(\hat{p}^2, \check{p}/\hat{p}, t)/\partial(\check{p}/\hat{p})} d(\hat{p}^2).$$

Since the regular forms ω of $\tilde{Y}$ fulfilling the equation $\sigma^*\omega = \omega$ are the pullbacks of the regular forms of the quotient $\tilde{Y}_\sigma$ of $\tilde{Y}$ modulo σ under the canonical map $\tilde{Y} \rightarrow \tilde{Y}_\sigma$, we conclude with the help of Lemma 2.20 that if the form ω is locally regular with respect to the description of $\tilde{Y}_\sigma$ by the equation $R_\sigma(\hat{p}^2, \check{p}/\hat{p}) = 0$, then it describes a deformation of *complex Fermi curves*, on which the function $\check{p}/\hat{p}$ extends locally near the preimage of $[\kappa/2]$ to a holomorphic function. In particular, all these *complex Fermi curves* obey the *Weak singularity condition* 4.2. Hence we may modify the construction of the complex manifold of deformations in Proposition 3.7 and obtain a complex submanifold of deformations of *complex Fermi curves*, which are invariant under the involution σ , and obey the *Weak singularity condition* 4.2.

Therefore, due to the considerations above, all *admissible* regular forms of this modified description of $\tilde{Y}_\sigma$ correspond to deformations of *Generalized Weierstraß curves* 4.4. As a consequence of this modification of the description of $\tilde{Y}_\sigma$, the *complex Fermi curves* have to be described in some neighbourhood of $[\kappa/2]$ as complete intersections in $\mathbb{C}^2$, which are invariant under σ . Therefore, over *Orbits of type 1* the involution η has fixed points on $\tilde{Y}$. But over *Orbits of type 2* and *Orbits of type 3* we may choose Y in such a way that η does not have fixed points. In particular, in the situations *Multiple points O2/O2*, *Multiple points O3*, *Cusps O2* and *Cusps O3* all regular forms of this choice of $\tilde{Y}_\sigma$ correspond to deformations of *Generalized Weierstraß curves* 4.4. If the corresponding Y_σ is not disconnected (compare with Remark 3.8), then there exists such deformations with decreasing *Generalized Willmore functional* 4.5.

Before we consider the remaining situations let us explain the structure of these deformations. Since the pullbacks of all local holomorphic forms of the normalization of the quotient modulo σ under the canonical two-sheeted covering from the normalization onto the normalization of the quotient modulo σ have a zero at all fixed points of σ , the deformations corresponding to such forms preserve fixed points of σ and therefore *Orbits of type 2*. An easy calculation shows that in general these four situations are deformed into two (and sometimes even more) *Orbits of type 2*. In fact, a double point of the form $(y, \sigma(y))$ should be considered as the limit of two colliding branch points (which are fixed points of σ) of this two-sheeted covering. At these singularities the functions $\hat{p}$ and $\check{p}$ and their first derivatives take the

same value. Therefore, the *Local contributions to the arithmetic genus* 4.3 this singularity is at least equal to 2. The deformation into a double point containing two fixed points of σ changes the *geometric genus* of the *complex Fermi curve*, but does not change the *geometric genus* of the quotient modulo σ . Consequently, the local contribution of these double points containing two fixed points of σ to the arithmetic genus is at least equal to 1.

In the remaining cases *Multiple points O1/O1*, *Multiple points O1/O2* and *Cusps O1* it suffices to determine the subspace of *admissible* regular forms of this modified quotient $\tilde{Y}_\sigma$ modulo σ . We observe that in these situations the normalization is connected. In fact, if the normalization is disconnected, then the involution ρ interchanges both infinities $\infty^\pm$ and therefore also both connected components. Therefore, in this case the real part is empty, which contradicts the existence of an *Orbit of type 1*. In the following discussion of these situations we shall derive sufficient conditions on the real regular forms of $\tilde{Y}_\sigma$ to be *admissible*. We will see that if a regular form takes special values at the preimage of $[\kappa/2]$ in the real part of the normalization, then it is *admissible*.

More precisely, let us use the description of the *complex Fermi curves* by equations of the form $R(\hat{p}, \check{p}) = 0$. Here R may be chosen to be a polynomial with respect to $\check{p}$. If its degree is equal to the number of those sheets of the covering map $\hat{p}$, which contain one of the elements of Y under consideration, and if the highest coefficient is equal to 1, then R is unique. Consequently the other coefficients are holomorphic functions depending on $\hat{p}$. Due to Lemma 2.20 the regular forms are locally of the form

$$\frac{f(\hat{p}, \check{p})}{\partial R(\hat{p}, \check{p})/\partial \check{p}} d\hat{p}.$$

Here again f may be chosen to be a polynomial with respect to $\check{p}$, whose degree is smaller than the degree of R . Due to the definition of the regular form corresponding to a deformation, $-f$ is equal to the derivative of R with respect to the deformation parameter of the corresponding family. The Taylor coefficients of R define locally holomorphic functions on the moduli space $\mathcal{M}_{g,\Lambda}$ constructed in Proposition 3.7. We will obtain some conditions on the derivatives of these Taylor coefficients with respect to the moduli, which are equal to the corresponding Taylor coefficients of f . Then we apply the following Lemma, which is a consequence of Serre duality ([Fo, §18.2] and [Se, Chapter IV §3.10.]).

Lemma 4.10. *Let $\tilde{Y}_\sigma$ be the quotient modulo σ corresponding to some $\mathcal{F}(Y, \infty^-, \infty^+, k) \in \bar{\mathcal{M}}_{g,\Lambda,\eta,\sigma}$, and let $y_1, \dots, y_l$ be finitely many points of $\tilde{Y}_\sigma$ in the complement of the double point (∞^-, ∞^+) . The values of all real regular forms of $\tilde{Y}_\sigma$, whose residue at ∞^+ is equal to $-\sqrt{-1}$ is the orthogonal complement of all meromorphic functions of $\tilde{Y}_\sigma$, which are holomorphic on $\tilde{Y}_\sigma \setminus \{(\infty^-, \infty^+), y_1, \dots, y_l\}$ and bounded on some small neighbourhood of (∞^-, ∞^+) with respect to the pairing given by the total residue.* **q.e.d.**

In particular, if Y corresponds to a relative minimizer, then it must not contain an *admissible* regular form, whose flow decreases the *Generalized Willmore functional* 4.5. Hence, due to the lemma, it has a meromorphic function with poles contained in the preimage of $[\kappa]/2$ described in the *Weak Singularity condition* 4.2 under the mapping k . In Proposition 3.5 and

Theorem 5.9 we construct this function with the help of some special tools of the integrable systems having Lax operators.

Corollary 4.11. *The exterior derivative of a linear combination of Taylor coefficients of R at a singularity corresponding to a regular form, which has poles at all points of the preimage in the normalization of this singularity cannot vanish.*

Proof. The only relation on the singular parts of meromorphic differentials on smooth Riemann surfaces Y is that the sum of the residues vanishes. Due to Lemma 2.20 this relation is even locally satisfied for all regular forms of Y , considered as meromorphic differentials on the normalization of Y . Hence the exterior derivative of such a Taylor coefficient considered as a holomorphic function on the moduli space cannot vanish. **q.e.d.**

This corollary implies that in the remaining situations *Multiple points O1/O2*, *Multiple points O1/O1* and *Cusps O1* the preimage of $[\kappa/2]$ contains no other *Orbit of type 1*. In fact, otherwise we choose a one-sheeted covering Y of the *complex Fermi curve* with no fixed points of η , and two points, which contain all *Orbits of type 1*. Now we may apply the corollary to all coefficients of order less than 2 in the situation of *Multiple points O1/O1/O2*, and to all coefficients of order less than 3 in the situations *Multiple points O1/O1/O1*, *Cusps O1/Multiple points O1* and *Cusps O1/O1*.

Let us now consider these situations.

Multiple points O1/O2. Due to our general consideration made above we have only to consider the *Orbit of type 1*. The Taylor expansions of the function $\check{p}$ in terms of $\hat{p}$ at the two points of the *Orbit of type 1* has to be of the form $\check{p} = a_1\hat{p} + a_2\hat{p}^2 + a_3\hat{p}^3 + \dots$ and $\check{p} = a_1\hat{p} - a_2\hat{p}^2 + a_3\hat{p}^3 - \dots$, respectively. Let $2m$ be the smallest even index, whose coefficient a_{2m} does not vanish. After adding to the function $\check{p}(\hat{p})$ some odd function with respect to $\hat{p}$, we may achieve that all coefficient with odd index vanish. We conclude that the quotient modulo σ may be described locally by some equation of the form $(\check{p}/\hat{p})^2 = a_{2m}^2(\hat{p}^2)^{2m-1} + 2a_{2m}a_{2m+2}(\hat{p}^2)^{2m} + \dots$. If we choose a regular form ω , which is locally given by

$$\omega = (2\check{p}/\hat{p})^{-1}(1 + \mathbf{O}(\hat{p}^2))d\hat{p}^2 = \pm a_{2m}^{-1}(\hat{p}^2)^{1-m}(1 + \mathbf{O}(\hat{p}^2))d\hat{p},$$

and extend this form to some holomorphic section of the tangent bundle of the moduli space on some neighbourhood of Y , then the corresponding family of *complex Fermi curves* may be described by the equation

$$(\check{p}/\hat{p})^2 = a_{2m}^2(\hat{p}^2)^{2m-1} + 2a_{2m}a_{2m+2}(\hat{p}^2)^{2m} + \dots + t(1 + \mathbf{O}(\hat{p}^2)) + \mathbf{O}(t^2).$$

This shows that for small non-zero t these *complex Fermi curves* have double points of the form (y, y') , where both points are fixed points of σ . The corresponding form has a pole of order $m - 1$ on the normalization of Y_σ over the *Orbit of type 1*.

If we include the involution η into our considerations, we see that on the normalizations of the corresponding *complex Fermi curves* the involution η may have fixed points. More

precisely, all the coefficients $a_1, a_2, \dots$ are real, if the corresponding *multiple points* belong to the real part. Then the last formula shows that at least for small t on the part of the deformation family corresponding to $t < 0$ the involution η does not have fixed points. The corresponding form on the normalization of the quotient modulo σ has a pole of order $l - 1$. With the help of Lemma 4.10 we conclude that for a relative minimizer m is equal to 1, and the *complex Fermi curve* is *real- σ -hyperelliptic*.

Cusps O1. We consider first a double point of the form $(y, \sigma(y))$ containing two *cusps* identified to one higher order singularity (the *Local contributions to the arithmetic genus* 4.3 of these singularities is at least equal to 8). In this case for almost all lattice vectors γ the function $g(\gamma, k)$ has a zero of order d , and for some element $\mu \in \mathbb{C}^2$ the function $g(\mu, k)$ has a zero of order $m > d$. These two numbers has to be co-prime. Otherwise the normalization would contain several points over these *cusps*. Moreover, due to Corollary 4.11 for a relative minimizer d has to be equal to 2. In fact, otherwise we choose a one-sheeted covering Y , on which η does not have fixed points. An application of this corollary to all coefficients of $R(\hat{p}, \check{p})$ of order less than 3 yields a deformation with decreasing *first integral*. In order to simplify notation we may assume that $\hat{p}$ has a zero of order 2 and $\check{p}$ has a zero of order m at the points $(y, \sigma(y))$ of the normalization. This implies that the normalization is locally nearby both points of $(y, \sigma(y))$ a two-sheeted covering with respect to $\hat{p}$. They obey locally some equation $\check{p}^2 = a_1 \hat{p}^m + \text{higher-order terms}$, and $\hat{p}^2 = -a_1 \check{p}^m + \text{higher-order terms}$, respectively. This implies

$$R_\sigma(\hat{p}^2, \check{p}/\hat{p}) = (\check{p}/\hat{p})^4 - a_1^2 (\hat{p}^2)^{m-2} + \text{higher-order terms} = 0.$$

If we choose a regular form ω , which is locally given by

$$\omega = \frac{1 + \mathbf{O}(\hat{p}^2) + \mathbf{O}(\check{p}/\hat{p})}{\partial R_\sigma(\hat{p}^2, \check{p}/\hat{p}) / \partial (\check{p}/\hat{p})} d(\hat{p}^2),$$

the flow corresponding to an appropriate extension of this form to a holomorphic vector field of the moduli space, will deform the equation above locally to an equation of the form

$$R_\sigma(\hat{p}^2, \check{p}/\hat{p}) + t(1 + \mathbf{O}(\hat{p}^2) + \mathbf{O}(\check{p}/\hat{p})) + \mathbf{O}(t^2) = 0.$$

We conclude that the multiple point $(y, \sigma(y))$ is deformed into four fixed points of σ . An easy calculation shows that this regular form ω may have a pole at the multiple points of order $3(m - 3)$. This order is zero, if and only if $m = 3$.

If we include η into our considerations, then we see again that only on half of the deformation family the involution η does not have fixed points. Therefore, Lemma 4.10 implies again that m is equal to 3 and that the *complex Fermi curve* is *real- σ -hyperelliptic*.

Multiple points O1/O1. If we choose $\tilde{Y}_\sigma$ in such a way that it contains two different singular points over the two *Orbits of type 1*, we may apply the arguments used in *Multiple points O1/O2* concerning the *Orbit of type 1*. This implies that one of the following cases occurs:

Lowest order singularity. The m 's corresponding to both *Orbits of type 1* are equal to 1 and the quotient Y_σ of Y modulo σ is hyperelliptic. Moreover, if f denotes the real function on Y_σ , which is invariant under the hyperelliptic involution, then the values of the function $d(d\check{p}/d\hat{p})/df$ at the two *Orbits of type 1* have the same sign.


Higher order singularity. One of the m 's is equal to 1 and Y is *real- σ -hyperelliptic*.

In order to complete the proof it suffices to show that the case of the *Lowest singularity* satisfies the conditions of *Minimizers with dividing real parts*.

We claim that, if the function $d\check{p}/d\hat{p}$ takes the same values at both *Orbits of type 1*, then Y cannot correspond to a local minimizer. In this case we choose $\tilde{Y}_\sigma$ to have one singular point over both *Orbits of type 1*. The analogous calculation of $R_\sigma(\hat{p}^2, \check{p}/\hat{p})$ and the corresponding deformations of lowest-order show that in this case there exists *admissible* regular forms of the modification of $\tilde{Y}_\sigma$ with prescribed singularity at the two *Orbits of type 1*. Hence there exists deformations decreasing the *Generalized Willmore functional* 4.5. We remark that in case of the *Lowest singularity* the *Local contributions to the arithmetic genus* 4.3 is not smaller than 8. If $d\check{p}/d\hat{p}$ takes the same value at both *Orbits of type 1*, then the *Local contributions to the arithmetic genus* 4.3 is not smaller than 12. In Lemma 5.1 (ii) we will see that in this case the *isospectral set* contains only *Weierstraß potentials* 1.3.

Now we show that in case of the *Lowest singularity* the hyperelliptic involution of Y_σ does not have real fixed points. Let us assume on the contrary that the hyperelliptic involution of the quotient modulo σ has real fixed points. This implies that on all connected components of the real part of the quotient modulo σ , the form df has exactly two zeroes, and furthermore f maps each connected component of the real part onto some real interval and besides the two endpoints of the interval (which have to be fixed points of the hyperelliptic involution and therefore zeroes of df) the preimage of each element of this interval contains exactly two points, at which the form df has different signs (with respect to any orientation of the real part). This implies that one real cycle of the quotient modulo σ contains the two points corresponding to the two *Orbits of type 1*, and that the form $d(d\check{p}/d\hat{p})$ has different signs at these two points (with respect to any orientation of this real cycle). The preimage of this real cycle under the natural map from the *complex Fermi curve* onto the quotient modulo σ consists of either one or two real cycles.

Let us now consider the images of these real cycles under the map k . These images are one or two closed curves in $\mathbb{R}^2/\Lambda^*$, which are invariant under the involution $k \mapsto -k$. Moreover, if we identify each pair of points, which are interchanged by this involution, then we obtain one connected component. Finally, nearby the four points corresponding to the two *Orbits of type 1* the images has to look like four paths, which all intersect in one point in two different tangent lines. More precisely, to each path there exists another path, which touches the path in this point. In the neighbouring figure we indicate theses images nearby the two *Orbits of type 1* and denote the open ends in the anti-clockwise-order by A, B, C, D, E, F, G and H .



We conclude that there are exactly four possibilities to connect the open ends. We denote these possibilities by combining those capitals into common brackets, which correspond to connected open ends. In fact, an open end of any path has to be connected to an open end of another path, which does not have the same tangent. Also, due to the condition that $d(d\check{p}/d\hat{p})$ has different signs at the two *Orbits of type 1*, each open end has to be connected with some other open end, which in our schematic picture may be obtained from the first open end by some rotation through ± 90 degrees. Therefore, we have the following possibilities:

1. $(A, C), (E, G), (B, D), (F, H)$
2. $(A, C), (E, G), (B, H), (F, D)$
3. $(A, G), (E, C), (B, D), (F, H)$
4. $(A, G), (E, C), (B, H), (F, D)$

We should remark that the integral of the form dk along a way connecting these open ends does not have to vanish, but due to the invariance under the involution $k \mapsto -k$, the integral has to be minus the integral of the corresponding image under this involution. Let 0 denote the image of the two *Orbits of type 1* under the map k . We conclude that in the first two cases the sequence of open ends denoted by $\dots - 0 - A - C - 0 - E - G - 0 - \dots$ and in the third and in the fourth case the sequence of open ends denoted by $\dots - 0 - A - G - 0 - E - C - 0 - \dots$ describes a closed curve in $\mathbb{R}^2$, which is defined up to translation by some element of Λ^* . On the other hand, in the first and the third case the sequence of open ends denoted by $\dots - 0 - B - D - 0 - F - H - 0 - \dots$ and in the second and in the fourth case the sequence of open ends denoted by $\dots - 0 - B - H - 0 - F - D - 0 - \dots$ also describes a closed curve in $\mathbb{R}^2$, which again is defined up to translation by some element of Λ^* . Moreover, since at all real double points the function f has to take the same value, these two closed curves may have selfintersections, but they must not intersect each other and the corresponding shifted curves outside of the points corresponding to the two *Orbits of type 1*. In fact, if

these points are identified with some other point of the corresponding curves, then the preimage of the corresponding element $[\kappa/2]$ under the normalization map contains at least three *Orbits of type 1*. However, this case was excluded in the above considerations. If we chose any orientation of these cycles, they represent some chains of the singular homology of $\mathbb{R}^2$. Since the first homology groups of $\mathbb{R}^2$ are trivial, the intersection numbers of these two cycles have to be zero ([S-T, §73] and [Do, Chapter VII §4]). An easy calculation of the local contributions to the intersection numbers of the two cycles shows, that those, which are not zero, have the same sign (depending on the choice of the orientation). Moreover, for some shifted representatives a local contribution is not zero. Hence these cycles have to intersect in some other points, which is a contradiction. Therefore, the hyperelliptic involution of the quotient modulo σ does not have real fixed points, and the real part divides the quotient into two connected components. Moreover, the real part of the quotient modulo σ has one connected component, if the *geometric genus* is even, and two connected components, if the *geometric genus* is odd. Finally, the real part of the *complex Fermi curve* divides the normalization into two connected components.

2. Relative minimizers of infinite *geometric genus*. If $\mathcal{F}(V, W)$ is a *complex Fermi curve* of potentials $V, W \in L^2(\mathbb{R}^2/\Lambda)$, which is invariant under η without fixed points, then due to Corollary 3.59 all forms of the form $\Omega_{V,W}(\delta V, \delta W)$, whose complex conjugate are equal to the pullback under η , correspond to infinitesimal deformations of *complex Fermi curves* in $\bar{\mathcal{M}}_{\Lambda,\eta}$. More precisely, the arguments of the proof of Theorem 3.57 extend to all *complex Fermi curves*, which are


- (i) invariant under η without fixed points,
- (ii) and have a finite value of the *first integral*.

Consequently, the correspondence between regular one-forms and one-dimensional families of deformations of *complex Fermi curves* obeying the *Weak Singularity condition* 4.2 extends from $\bigcup_{g \in \mathbb{N}} \bar{\mathcal{M}}_{g,\Lambda,\eta,\sigma}$ to $\bigcup_{\mathcal{W} > 0} \bar{\mathcal{M}}_{\Lambda,\eta,\sigma,\mathcal{W}}$. Therefore, Proposition 3.5 shows that the same conclusion holds in the general case (compare with Remark 3.51). In particular, all relative minimizers of the *Generalized Willmore functional* 4.5, whose Willmore functional is smaller than $\mathcal{W}$, belong to $\bar{\mathcal{M}}_{g_{\max},\Lambda,\eta,\sigma,\mathcal{W}}$, with the corresponding integer $g_{\max}$ introduced in Corollary 3.54.
q.e.d.

At the end of this section we want to determine all possible real double points of a relative minimizer. In the situations *Multiple points O1/O2*, *Cusps O1* and a *Higher singularity* of *Multiple points O1/O1* the function f on Y_σ of degree one has to take the same values at the double points. Therefore, they are double points of the form $(y, \sigma(y))$.

In the situation of two *Lowest singularities* of *Multiple points O1/O1* we know that in the notation introduced above we have the following possibilities to connect the open ends nearby the two *Orbits of type 1*:

1. $(A, B), (E, F), (C, D), (G, H)$
2. $(A, B), (E, F), (C, H), (G, D)$
3. $(A, F), (E, B), (C, D), (G, H)$
4. $(A, F), (E, B), (C, H), (G, D)$
5. $(A, D), (E, H), (B, C), (F, G)$
6. $(A, D), (E, H), (B, G), (F, C)$
7. $(A, H), (E, D), (B, C), (F, G)$
8. $(A, H), (E, D), (B, G), (F, C)$



Again we use the cycles corresponding to the first two pairs and the last two pairs (for example in the first case the cycles denoted by $\dots - 0 - A - B - 0 - E - F - 0 - \dots$ and $\dots - 0 - C - D - 0 - G - H - 0 - \dots$). Again they correspond to closed cycles in $\mathbb{R}^2$ and their intersection numbers has to vanish. A direct calculation shows that the local contribution to the intersection numbers at the two *Orbits of type 2* cancels. Therefore, the number of points with non-zero contribution to these intersection numbers has to be even. Furthermore, due to the invariance under σ , these points have to occur in pairs with equal contribution to the intersection numbers. Finally, due to Lemma 4.10, the function f has to take the same value at any two points of the normalization, which belong to one singularity. Hence the pairs of such double points with non-zero contribution to the intersection number occur in different pairs, where f takes different values, or four double points, where f takes only one value. Now we claim that the first case is impossible. In fact, if we choose two of these pairs with opposite local contribution to the intersection number, we may use these double points in order to connect the cycles nearby the two *Orbits of type 1* in such a way, that the differential df has different sign at the two *Orbits of type 1* with respect to any orientation of these cycles. The considerations concerning this case in the proof of Theorem 4.9 implies that then we have a double point containing two real points of the normalization, where f takes different values. This contradicts Lemma 4.10. In the second case we have again two *Orbits of type 1* over some other element of $\Lambda^*/2\Lambda^*$. In this case the same considerations apply to this singularity.

We conclude that we may deform all these *Orbits of type 1* by some arbitrary small deformation increasing the *first integral* into *Orbits of type 2*. After this deformation the real part will not have any double point. We claim that the function $k_1 + \sqrt{-1}k_2$ maps the connected component Y^+ of the complement of the real part in the normalization, which contains ∞^+ , onto the relative complement in a finite-sheeted covering of the torus $\mathbb{C}/\Lambda_{\mathbb{C}}^*$ of finitely many disjoint sets of this torus. Here $\Lambda_{\mathbb{C}}^* = \{\kappa_1 + \sqrt{-1}\kappa_2 \in \mathbb{C} \mid \kappa \in \Lambda^*\}$ denotes the complex version of the dual lattice. We shall prove the analogous stronger statement for the corresponding quotients modulo the involution σ . The Weierstraß function $\wp$ corresponding to the complex torus $\mathbb{C}/\Lambda_{\mathbb{C}}^*$ (see e.g. [Bat, 13.12]) represents this complex torus as a two-sheeted covering over $\mathbb{P}^1$, whose hyperelliptic involution is induced by the involution $\sigma : k \mapsto$

$-k$. The four elements of $\Lambda^*/2\Lambda^*$ are the four fixed points of this involution. Consequently the composition of the function $k_1 + \sqrt{-1}k_2$ with $\wp$ yields a function on the quotient Y_σ^+ of Y^+ modulo the involution σ . Nearby the two *Orbits of type 2* (the deformed *Orbits of type 1*) this function maps Y_σ^+ onto the complement of finitely many (either one or two) disjoint regions of $\mathbb{P}^1$. Since the real part has no double points, and therefore does not intersect itself, the same is true for the whole of Y_σ^+ . Consequently we obtain

Corollary 4.12. *All relative minima may be deformed by some arbitrary small deformation increasing the first integral into a Generalized Weierstraß curve 4.4 obeying the following conditions:*

- (i) *The real part divides the normalization into two connected components Y_{normal}^- and Y_{normal}^+ containing ∞^- and ∞^+ , respectively.*
- (ii) *The function $k_1 + \sqrt{-1}k_2$ maps Y_{normal}^+ into the relative complement in a finite-sheeted covering of $\mathbb{C}/\Lambda_{\mathbb{C}}^*$ of finitely many disjoint closed subsets of this complex torus.*
- (iii) *All multiple points, which connect the two components Y_{normal}^+ and Y_{normal}^- are of the form $(y, \eta(y))$, where the imaginary part of the function $d\bar{p}/d\bar{p}$ has the same sign as at the corresponding points $\infty^\pm$, respectively.*
- (iv) *For some $[\kappa/2] \in \Lambda^*/2\Lambda^*$ the normalization of the complex Fermi curve contains either two Orbits of type 2 or one Orbit of type 3 over this element.* **q.e.d.**

Corollary 3.54 suggests, that for large g all relative minimizers in $\bar{\mathcal{M}}_{g,\Lambda,\eta,\mathcal{W}}$ fulfill the conditions of Theorem 4.9. A proof of such a statement could make it possible to prove the Willmore conjecture without making use of the results of Sections 3.4–3.5. At an earlier state of this project, we thought to have a proof of such a statement. However, we overlooked the possibility of *real cusps* (compare with Proposition 3.9 and Remark 4.22). We did not succeed to overcome this problem. Afterwards we succeed to prove the compactness of the moduli spaces without any bound on the *geometric genus* of the *complex Fermi curves*.

4.3 Absolute minimizers of the *Generalized Willmore functional*

In Section 4.2 we obtained a classification of relative minimizers of the *Generalized Willmore functional* 4.5. In this section we will determine all relative minimizers of the *Generalized Willmore functional* 4.5, whose *first integral* is not larger than 8π . The *Weak Singularity condition* 4.2 is, roughly speaking, a condition on four points of the *complex Fermi curve*. As a preparation we shall first determine the minimum of the restriction of the *first integral* to the space of all *complex Fermi curves*, which contain a fixed element $k' \in \mathbb{R}^2$.

Lemma 4.13. *For all elements $[k'] \in \mathbb{R}^2/\Lambda^*$ there exists a unique absolute minimizer of the restriction of the first integral to the space of all complex Fermi curves in $\bigcup_{\mathcal{W}>0} \bar{\mathcal{M}}_{\Lambda,\eta,\sigma,\mathcal{W}}$, which contains $[k']$. These minimizers fit together to a continuous map*

$$\mathcal{F}_{\min} : \mathbb{R}^2/\Lambda^* \rightarrow \bar{\mathcal{M}}_{3,\Lambda,\eta,\sigma,4\pi}, [k] \mapsto \mathcal{F}_{\min}([k])$$

into the set of real- σ -hyperelliptic complex Fermi curves in $\bar{\mathcal{M}}_{3,\Lambda,\eta,\sigma,4\pi}$. The restriction of the first integral to the image of this mapping yields a homeomorphism onto $[0, 4\pi]$. Therefore, we may parameterize this family as well by $\mathcal{W}$:

$$\mathcal{F}_{\min} : [0, 4\pi] \rightarrow \bar{\mathcal{M}}_{3,\Lambda,\eta,\sigma,4\pi}, \mathcal{W} \mapsto \mathcal{F}_{\min}(\mathcal{W}).$$

Proof. The space of all $\mathcal{F} \in \bar{\mathcal{M}}_{\Lambda,\eta,\sigma}$, which contain an element $[k'] \in \mathbb{R}^2/\Lambda^*$ is a subvariety, which is locally given by the zero set of at most two analytic functions. Due to the compactness this subvariety contains a minimizer of the *first integral*.

The preimage of $[k']$ (and $[-k']$) in the normalization of $\mathcal{F}$ contains either fixed points of ρ (i. e. real points) or pairs, which are interchanged by ρ . If in the latter case the *Structure sheaf* 2.24 contains functions, which take different values at a pair, then a *Bäcklund transformation* with the corresponding values of the eigenfunctions ψ transforms the potential into a potential, whose *Structure sheaf* 2.24 contains only functions taking the same value at these pairs. We remark that the formulas of Lemma 2.29 also apply to *Finite rank perturbations* 3.22. Nearby this transformed potential the subvariety described in the lemma is the zero set of one analytic function. If the preimage of $[k']$ contains real points, then the subvariety described in the Lemma is the zero set of one analytic function. Therefore, we may assume that nearby the minimizer the subvariety described in the lemma is equal to the zero set of one analytic function.

Then Proposition 3.5 shows that that if $\mathcal{F} \in \mathcal{M}_{\Lambda,\eta,\sigma}$ is a relative minimum of the restriction of the *first integral* to this subvariety, then this *complex Fermi curve* is σ -hyperelliptic (i. e. the quotient modulo the involution σ is isomorphic to $\mathbb{P}^1$ with two points removed). We remark that these arguments lead to the same conclusion, if the derivative of that single analytic function vanishes, whose zero set is the subvariety described in the Lemma. Moreover, in case that the preimage of $[k']$ in the normalization contains pairs, which are interchanged by ρ , then the corresponding function constructed in Proposition 3.5 would have no poles, and the corresponding *complex Fermi curve* has a disconnected normalization. (We shall see that this is the case for $[k'] = 0$ and those $[k']$, which are the unique real elements of the unique *complex Fermi curves* with disconnected normalization, whose *first integral* is equal to 4π). If the preimage of $[k']$ in the normalization contains a fixed point of ρ , then the *complex Fermi curve* is *real σ -hyperelliptic*. To sum up, the minima in $\mathcal{M}_{\Lambda,\eta,\sigma}$ have dividing real parts.

These conclusions are also true for minima in $\bigcup_{\mathcal{W}>0} \bar{\mathcal{M}}_{\Lambda,\eta,\sigma,\mathcal{W}}$. In fact the construction of $(\delta V_f, \delta W_f)$ in Lemma 3.2 extends to the corresponding variations on the complement of the union $\mathcal{S}_{D,\varepsilon}$ of the union of arbitrary small balls around the singular set described in Theorem 3.48 (compare with Remark 3.51). Since the union over ε of all subspaces of $L^2(\mathbb{R}^2/\Lambda) \times L^2(\mathbb{R}^2/\Lambda)$ of elements, which vanish on $\mathcal{S}_{D,\varepsilon}$ is dense, the arguments given above extends to all minima of the *first integral* restricted to the subvariety described in the Lemma. Therefore, due to Corollary 3.54, these minima have finite genus.

We shall see in Lemma 4.16 that the *first integral* of any *complex Fermi curve* with disconnected normalization is a multiple of 4π . The unique *complex Fermi curve*, whose *first integral* vanishes (i. e. the free *complex Fermi curve* introduced in Section 2.2) contains

only the real element $[k'] = 0$. Obviously, the corresponding *complex Fermi curves* depend continuously on $[k']$. Therefore, for small non-zero k' these *real- σ -hyperelliptic* minimizers have genus zero. These *complex Fermi curves* are given by the equation $g(k, k) = g(k', k')$. More precisely, if $g(k', k')$ is smaller than $1/4$ times the square of the length of the shortest non-zero dual lattice vector, this *complex Fermi curve* is the minimizer and all double points of this *complex Fermi curve* are of the form $(y, \eta(y))$. If $g(k', k')$ is equal to $1/4$ times the square of the length of the shortest non-zero dual lattice element, then the *complex Fermi curve* contains *Orbits of type 1* over a finite number of elements of $\Lambda^*/2\Lambda^*$. If the length of k' becomes larger, the *complex Fermi curve* will be deformed into a *complex Fermi curve*, with as much *Orbits of type 2*, as the former *complex Fermi curve* has *Orbits of type 1*. We will see that in general there is only one *Orbit of type 2*. In this case the real part has two connected components, and the integral of dk over any component is non-zero. The map $k_1 + \sqrt{-1}k_2$ maps the closure of Y^+ onto some part of the elliptic curve $\mathbb{C}/\Lambda_{\mathbb{C}}^*$. Moreover, this map has no branch point, and therefore may be considered as a subset of this elliptic curve. Obviously, the image of the closure of Y^+ in the elliptic curve covers all points $[k_1]$, whose *complex Fermi curve* $\mathcal{F}_{\min}(k_1)$ belongs to the family of the deformation of $\mathcal{F}_{\min}(k')$ into $\mathcal{F}_{\min}(0)$. If we move k' into the complement of the image of the closure of Y^+ under the map $k_1 + \sqrt{-1}k_2$, then this subset will become larger until the boundaries meet each other. In this case the corresponding *complex Fermi curve* has again some *Orbits of type 1* over some elements of $\Lambda^*/2\Lambda^*$. In fact, since it is defined as a relative minimizer, all double points of the whole family have to be either of the form $(y, \eta(y))$ or the function introduced in Proposition 3.5 has to take the same values at these points, which implies that they are of the form $(y, \sigma(y))$. Since both points are invariant under ρ they fit together to an *Orbit of type 1* over some element of $\Lambda^*/2\Lambda^*$. If we enlarge the length of $g(k', k')$ further, these *Orbits of type 1* will again be deformed into *Orbits of type 2*. The maps of the closures of the corresponding Y^+ will still define some one-sheeted covering of the corresponding elliptic curve. But the boundary will look like one or two circle, which are homologous to zero. If we move k' inside the part, which corresponds to the complement of the image of Y^+ under the map $k_1 + \sqrt{-1}k_2$ in the elliptic curve, the *first integral* will be enlarged and any *Orbit of type 1* of the boundary will be deformed into an *Orbit of type 2*. Finally, the whole elliptic curve is covered, and the *first integral* is equal to 4π . This shows that for all k' the *first integral* is not larger than 4π . Moreover, by construction this family is periodic with respect to the dual lattice and therefore defines some continuous map from $\mathbb{R}^2/\Lambda^*$ into the set of σ -hyperelliptic *complex Fermi curves*, which contains the preimage of this map. We conclude that for all $\mathcal{W} \in [0, 4\pi]$ there exists one *real- σ -hyperelliptic complex Fermi curve*, all whose double points are of the form $(y, \eta(y))$. Obviously $\mathcal{F}_{\min}(k_1)$ coincide with $\mathcal{F}_{\min}(k_2)$, if and only if k_2 belongs to the real part of $\mathcal{F}_{\min}(k_1)$ which is equivalent that k_1 belongs to the real part of $\mathcal{F}_{\min}(k_2)$.

We now claim that this family contains all *real- σ -hyperelliptic complex Fermi curves*, whose *first integral* is not larger than 4π , and which are relative minimizers of the restrictions of the *first integral* to the *complex Fermi curves* containing any element of the real part of the given *complex Fermi curve*. In fact, all these elements have to be *real- σ -hyperelliptic*, and they can have only double points of the form $(y, \eta(y))$. Moreover, we may deform any

such *complex Fermi curve* within this class into the unique *complex Fermi curve*, whose *first integral* is zero. Hence this family has to meet the other family at some point. But the space of *real- σ -hyperelliptic complex Fermi curves* is locally a one-dimensional space with some multiple points, where it has *Orbits of type 1*. Any such *Orbit of type 1* may be either deformed into double points of the form $(y, \eta(y))$ not belonging to the real part, or by changing the *first integral* in the other direction either into *Orbits of type 2* or into real double points not of the form $(y, \eta(y))$. In particular, the space of *real- σ -hyperelliptic complex Fermi curves*, all whose double points are of the form $(y, \eta(y))$, is locally a one-dimensional manifold. Also the *first integral* defines an injective function to $\mathbb{R}^+$. This proves the claim.

We conclude that this family contains all *real- σ -hyperelliptic complex Fermi curves*, whose *first integral* is smaller than 4π , and which contain only double points of the form $(y, \eta(y))$. **q.e.d.**

In particular, if the absolute minimizers of the restriction of the *first integral* to the space of all *complex Fermi curves*, which contain an element of $\Lambda^*/2\Lambda^*$, are of the following form:

Corollary 4.14. *For all elements $[\kappa/2] \in \Lambda^*/2\Lambda^*$ there exists a unique complex Fermi curve $\mathcal{F}_{\min}(\kappa/2)$, which minimizes the restriction of the first integral to the space of all complex Fermi curves, which contain the element $[\kappa/2]$ ($\mathcal{F}_{\min}(0)$ is the unique complex Fermi curve with vanishing first integral). All these minimizers are σ -hyperelliptic and the geometric genus is at most equal to two. More precisely, the genus is equal to the number of Orbits of type 2, which are contained in the normalization of Y . Each Orbit of type 2 is mapped onto one non-zero element of $\Lambda^*/2\Lambda^*$, and over all non-zero elements of $\Lambda^*/2\Lambda^*$ there exists at most one Orbit of type 2. Finally, the first integral of $\mathcal{F}_{\min}(\kappa/2)$ is larger or equal to the first integral of $\mathcal{F}_{\min}(\kappa'/2)$, if and only if $[\kappa'/2]$ is contained in $\mathcal{F}_{\min}(\kappa/2)$. **q.e.d.***

Remark 4.15. *In the proof of Lemma 4.13 we have actually shown, that the first integral of $\mathcal{F}_{\min}(\kappa/2)$ is larger than the first integral of $\mathcal{F}_{\min}(\kappa'/2)$, if and only if $\mathcal{F}_{\min}(\kappa/2)$ contains an Orbit of type 2 over $[\kappa'/2]$. Moreover, the first integrals are equal, if and only if both complex Fermi curves coincide and contain Orbits of type 1 over both elements.*

The proof of Lemma 4.13 shows that there is either one element in $\Lambda^*/2\Lambda^*$ or two different elements of $(\mathbb{R}^2 \setminus \Lambda^*/2) / \Lambda^*$, whose corresponding minimum of the *first integral* is equal to 4π . Moreover, the proof of this lemma shows that the real part of this *complex Fermi curve* is empty. In fact, these points are the limits of the real parts of the family of zero energy limits with lower values of the *first integral* as described in this Lemma. Since the real part of any *real- σ -hyperelliptic complex Fermi curve* divides the normalization into two connected components, we conclude that this *complex Fermi curve* has disconnected normalization. Let us consider these *complex Fermi curves* with disconnected normalization in some more detail. Due to Theorem 2.12 they have finite *geometric genus*.

Lemma 4.16. *If $\mathcal{F} \in \bigcup_{g \in \mathbb{N}} \bar{\mathcal{M}}_{g, \Lambda, \eta}$ has a disconnected normalization, then the component $\bar{\mathcal{F}}_{\text{normal}}^+$ of the normalization of $\bar{\mathcal{F}}$, which contains ∞^+ , is a finite-sheeted covering of the unique complex torus, which is conformally equivalent to $\mathbb{R}^2 / \Lambda$. Moreover, the first integral*

of $\mathcal{F}$ is 4π times the number of sheets of this covering. Furthermore, for all conformal classes there exists a real two-dimensional family of such complex Fermi curves with disconnected normalization in $\bigcup_{g \in \mathbb{N}} \bar{\mathcal{M}}_{g, \Lambda, \eta, 4\pi}$, and only one such complex Fermi curve in $\bigcup_{g \in \mathbb{N}} \bar{\mathcal{M}}_{g, \Lambda, \eta, \sigma, 4\pi}$.

Proof. If $\mathcal{F} \in \bigcup_{g \in \mathbb{N}} \bar{\mathcal{M}}_{g, \Lambda, \eta}$ has disconnected normalization, then the functions $k_1 \pm \sqrt{-1}k_2$ extend to holomorphic multi-valued functions on the connected component $Y_{\text{normal}}^{\pm}$ of the normalization of Y , which contain $\infty^{\pm}$, respectively. More precisely, they define holomorphic functions into the complex torus $\mathbb{C}/\Lambda_{\mathbb{C}}^*$. Due to the Open Mapping Theorem [Co-I, Chapter IV §7. 7.5] these mappings are either constant or open. Due to Lemma 3.16 in the first case the *complex Fermi curve* is the unique *complex Fermi curve*, whose *first integral* vanishes. Due to the compactness of $Y_{\text{normal}}^{\pm}$ in the second case these mappings are finitely-sheeted coverings. Moreover, due to Lemma 3.16, the *first integral* is equal to 4π times the number of sheets of these coverings. With the help of the formula of Lemma 3.12 it can be shown that in these cases the *first integral* is equal to 4π times the intersection number of those cycles in $H_1(Y_{\text{normal}}^+, \mathbb{Z})$, whose intersection numbers with any cycle in $H_1(Y_{\text{normal}}^+, \mathbb{Z})$ is equal to the integral of $d\hat{p}$ and $d\check{p}$ over the cycle, respectively. Due to Poincaré duality this intersection number is again equal to the number of sheets of these coverings.

If the *first integral* is equal to 4π , the normalization $\bar{\mathcal{F}}_{\text{normal}}^+$ has to be biholomorphic to the complex torus $\mathbb{C}/\Lambda_{\mathbb{C}}^*$ with one marked point 0 corresponding to $\infty^{\pm}$. Furthermore, the one-forms $d\hat{p}$ and $d\check{p}$ on the complex torus are uniquely determined by their singular part at ∞^+ (compare with condition *Quasi-momenta* (i) in Section 2.5) and the condition that for all $\kappa \in \Lambda^*$ the integral along the corresponding cycle of $\mathbb{C}/\Lambda_{\mathbb{C}}^*$ is equal to $g(\hat{\gamma}, \kappa)$ and $g(\check{\gamma}, \kappa)$, respectively. We conclude that the function k on $\bar{\mathcal{F}}_{\text{normal}}^{\pm}$ are determined up to some additive constants k^+ and k^- . Since on $\bar{\mathcal{F}}_{\text{normal}}^{\pm}$ the function $k_1 \pm \sqrt{-1}k_2$ maps $\infty^{\pm}$ onto 0, and the involution $k \mapsto -\bar{k}$ interchanges $\bar{\mathcal{F}}_{\text{normal}}^{\pm}$, there exists only a real two-dimensional family of such *complex Fermi curves* (compare with Proposition 3.7 and Remark 3.8). The invariance under the involution $k \mapsto -\bar{k}$ determines the function k uniquely up to shifts by dual lattice elements. **q.e.d.**

We shall construct the unique *complex Fermi curve* in $\bar{\mathcal{M}}_{3, \eta, \sigma, 4\pi}$ with disconnected normalization explicitly. Obviously it is enough to determine the meromorphic multi-valued functions $\hat{p}$ and $\check{p}$ on the component $\bar{\mathcal{F}}_{\text{normal}}^+$. In the foregoing lemma we have shown that this Riemann surface is biholomorphic to the elliptic curve $\mathbb{C}/\Lambda_{\mathbb{C}}^*$. We realize this elliptic curve as the quotient $\mathbb{C}/(2\omega\mathbb{Z} + 2\omega'\mathbb{Z})$, where $2\omega = \hat{\kappa}_1 + \sqrt{-1}\hat{\kappa}_2$ and $2\omega' = \check{\kappa}_1 + \sqrt{-1}\check{\kappa}_2$. After some translation we may identify the point ∞^+ with the point $z = 0$. Therefore, the involution σ is induced by the map $z \mapsto -z$. The functions $\hat{p}$ and $\check{p}$ satisfy $\hat{p}(z+2\omega) = \hat{p}(z)+1$, $\hat{p}(z+2\omega') = \hat{p}(z)$ and $\check{p}(z+2\omega) = \check{p}(z)$, $\check{p}(z+2\omega') = \check{p}(z)+1$. Since $\hat{p}$ and $\check{p}$ have poles of first-order at $z = 0$ this uniquely determines these functions, and they may be expressed in terms of the Weierstraß functions (see e.g. [Bat, 13.12]):

$$\hat{p}(z) = \sqrt{-1} \frac{\omega' \zeta(z) - \eta' z}{\pi} \qquad \check{p}(z) = -\sqrt{-1} \frac{\omega \zeta(z) - \eta z}{\pi}.$$

This implies that $(\hat{p}, \check{p})$ is at ω equal to $(-1/2, 0)$, at ω' equal to $(0, -1/2)$ and at $\omega + \omega'$ equal to $(-1/2, -1/2)$. Together with 0 these are all fixed points of σ , therefore the normalization of this *complex Fermi curve* has over all non-zero elements of $\Lambda^*/2\Lambda^*$ exactly one *Orbit of type 2* and no other *Orbit of type 2*. This follows also from the proof of Lemma 4.13, since the unique *complex Fermi curve* with disconnected normalization whose *first integral* is equal to 4π , is the same *complex Fermi curve* as the unique element of the family described in Lemma 4.13 whose *first integral* is equal to 4π . Moreover, since $d\check{p} - \tau d\hat{p}$ is equal to $-dz/\omega$ the integral of the form dz from any element of $\Lambda^*/2\Lambda^*$ to any other element has to be an element of $\omega\mathbb{Z} + \omega'\mathbb{Z}$. This shows that these *Orbits of type 2* are all elements of $\Lambda^*/2\Lambda^*$ of this *complex Fermi curve*. Therefore, there does not exist a *Generalized Weierstraß curves* 4.4 with disconnected normalization, whose *first integral* is smaller than 8π . However, we may use this *complex Fermi curve* in order to construct *Generalized Weierstraß curves* 4.4 with disconnected normalization, whose *first integral* is equal to 8π .

In Lemma 3.16 we cut the *complex Fermi curves* into two parts, on both of which the function k is single-valued. Each part contains one of the two marked points $\infty^\pm$, and the condition that $k_1 \pm \sqrt{-1}k_2$ should vanish at $\infty^\pm$ uniquely determines the branch of k on both parts. The differences between these functions along the cuts are elements of Λ^* . Also it might happen, that these elements generate a sublattice. Let us denote this sublattice by $\Lambda_{\text{effective}}^* \subset \Lambda^*$ and the corresponding dual lattice by $\Lambda_{\text{effective}} \supset \Lambda$. More precisely, we define the effective period lattice $\Lambda_{\text{effective}}$ as the dual lattice of $\Lambda_{\text{effective}}^*$:

$$\Lambda_{\text{effective}} = \{ \gamma \in \mathbb{R}^2 \mid g(\gamma, \kappa) \in \mathbb{Z} \ \forall \kappa \in \Lambda_{\text{effective}}^* \}.$$

If $\Lambda_{\text{effective}}^*$ is contained in a one-dimensional subspace of $\mathbb{R}^2$, there exists a lattice vector $\gamma \in \Lambda$, which is orthogonal to the whole lattice $\Lambda_{\text{effective}}^*$. This implies that $g(\gamma, k)$ extends to a single-valued meromorphic function on the normalization of the *complex Fermi curve*, and the corresponding flow acts trivially on all line bundles of the normalization of this *complex Fermi curves*. In this case $\Lambda_{\text{effective}}$ is by definition no longer any lattice. However, the intersection of this group with the one-dimensional subspace generated by $\Lambda_{\text{effective}}^*$, is a one-dimensional lattice. In the most degenerated case $\Lambda_{\text{effective}}^*$ is equal to $\{0\}$, and $\Lambda_{\text{effective}}$ is equal to $\mathbb{R}^2$. This implies that both components of k extend to single-valued meromorphic functions on the normalization of the *complex Fermi curve*, and both flows act trivially on the line bundles of the normalization of the *complex Fermi curve*. In this case the *geometric genus* is zero and the corresponding potentials of the form $(U, \bar{U})$ have only one non-vanishing Fourier coefficient (compare with the motivation of Theorem 3.57). In particular, the corresponding real potentials are constant. More generally, if the normalization of a given *complex Fermi curve* is connected, then due to Lemma 2.27, there exists a complex potential U , which is invariant under translations of all elements of $\Lambda_{\text{effective}}$, and whose *complex Fermi curve* is equal to the given one.

Remark 4.17. *The integral of dk over cycles defines a map from $H_1(Y_{\text{normal}}, \mathbb{Z})$ into Λ^* . The image of this map is contained in the sublattice $\Lambda_{\text{effective}}^* \subset \Lambda^*$. But due to condition Quasi-momenta (i) in Section 2.5 the image might be a proper sublattice of this sublattice. However, in these cases the transformation of Lemma 2.11 transforms the *complex Fermi curve* into*

other isomorphic complex Fermi curve without changing the image of this map, and such that this image coincides with the sublattice corresponding to the transformed complex Fermi curves.

Now let us assume that $\mathcal{F} \in \bigcup_{\mathcal{W} > 0} \bar{\mathcal{M}}_{\Lambda, \eta, \mathcal{W}}$ is a complex Fermi curve, whose effective dual lattice $\Lambda_{\text{effective}}^*$ is a proper sublattice of Λ^* . Obviously, there exists a proper sublattice $\check{\Lambda}^*$ with the following properties:

Dual sublattice (i) The proper sublattice $\check{\Lambda}^* \subset \Lambda^*$ contains $\Lambda_{\text{effective}}^*$.

Dual sublattice (ii) The order of $\Lambda^*/\check{\Lambda}^*$ is finite.

The dual lattice denoted by $\check{\Lambda}$ contains Λ , and the order of $\check{\Lambda}/\Lambda$ is the same as the order of the quotient in *Dual sublattice (ii)*. For all complex Fermi curves $\check{\mathcal{F}} \in \bigcup_{g \in \mathbb{N}} \bar{\mathcal{M}}_{g, \check{\Lambda}, \eta}$, the finite union of the subvarieties $\check{\mathcal{F}} \subset \mathbb{C}^2$ shifted by representatives contained in all elements of $\Lambda^*/\check{\Lambda}^*$ yields a complex Fermi curve $\mathcal{F} \in \bigcup_{g \in \mathbb{N}} \bar{\mathcal{M}}_{g, \Lambda, \eta}$, whose effective dual lattice is contained

in $\check{\Lambda}^*$. Moreover, this relation provides a one-to-one correspondence between complex Fermi curves with lattice $\check{\Lambda}$ and complex Fermi curves with lattice Λ , whose effective dual lattice is contained in $\check{\Lambda}^*$. The quotient $\check{\mathcal{F}}/\check{\Lambda}^*$ is a one-sheeted covering over the quotient $\mathcal{F}/\Lambda^*$. In particular, the normalizations of these quotients are isomorphic. The first integral of the complex Fermi curve $\mathcal{F}$ with lattice Λ is $|\check{\Lambda}/\Lambda|$ times the first integral of the complex Fermi curve $\check{\mathcal{F}}$ with lattice $\check{\Lambda}$.

Furthermore, if the complex Fermi curve with lattice $\check{\Lambda}$ is a Generalized Weierstraß curve 4.4, then the complex Fermi curve with lattice Λ is also a Generalized Weierstraß curve 4.4. However, there are some situations, where the complex Fermi curve with lattice Λ is a Generalized Weierstraß curve 4.4, whereas the complex Fermi curve with lattice $\check{\Lambda}$ is not a Generalized Weierstraß curve 4.4. Of special importance is the following situation: Let $\mathcal{F}$ be a complex Fermi curve with lattice $\check{\Lambda}$, containing two different elements of $\check{\Lambda}^*/2\check{\Lambda}^*$, whose difference is mapped under the canonical map

$$\check{\Lambda}^*/2\check{\Lambda}^* \rightarrow \Lambda^*/2\Lambda^*$$

onto zero. Then the complex Fermi curve with lattice Λ is a Generalized Weierstraß curve 4.4 in contrast to the complex Fermi curve with lattice $\check{\Lambda}$ not necessarily being a Generalized Weierstraß curve 4.4. We will see that this is true for the minimizers of the Generalized Willmore functional 4.5.

For any lattice Λ^* there exist exactly three sublattices $\check{\Lambda}^*$, with the property that the quotient lattice $\Lambda^*/\check{\Lambda}^*$ is isomorphic to $\mathbb{Z}_2$. In fact, all these lattices have to contain the lattice $2\Lambda^*$, and exactly one non-zero element of $\Lambda^*/2\Lambda^*$ is contained in the image of the natural map $\check{\Lambda}^*/2\check{\Lambda}^* \rightarrow \Lambda^*/2\Lambda^*$. Hence we may parameterize these sublattices by the non-zero elements of this group. For all non-zero $[\kappa/2] \in \Lambda^*/2\Lambda^*$ let $\Lambda_{[\kappa/2]}^*$ denote the unique sublattice with the following properties:

$\Lambda_{[\kappa/2]}^*$ (i) The group $\Lambda^*/\Lambda_{[\kappa/2]}^*$ is isomorphic to $\mathbb{Z}_2$.

$\Lambda_{[\kappa/2]}^*$ (ii) The image of the natural map $\Lambda_{[\kappa/2]}^*/2\Lambda_{[\kappa/2]}^* \rightarrow \Lambda^*/2\Lambda^*$ is equal to $\{0, [\kappa/2]\}$.

The above considerations have the following consequence: If $\check{\mathcal{F}}$ is the unique *complex Fermi curve*, with dual lattice $\Lambda_{[\kappa/2]}^*$ with disconnected normalization, and whose *first integral* is equal to 4π , then the normalization of the corresponding *complex Fermi curve* $\mathcal{F}$ with dual lattice Λ^* has two *Orbits of type 2* over the element $[\kappa/2]$ and one *Orbit of type 2* over the zero element of $\Lambda^*/2\Lambda^*$. One can prove that these are all elements of the form $\Lambda^*/2\Lambda^*$. Moreover, the *first integral* is equal to 8π . In particular, we have three *Generalized Weierstraß curves* 4.4, whose *first integrals* are equal to 8π . We will now show that these are all *Generalized Weierstraß curves* 4.4 with disconnected normalization, whose *first integrals* are equal to 8π .

Lemma 4.18. *The set of Generalized Weierstraß curves 4.4 with disconnected normalizations, whose first integrals are equal to 8π , consists of exactly three elements. More precisely, for each non-zero element $[\kappa/2] \in \Lambda^*/2\Lambda^*$ there exists exactly one such complex Fermi curve fulfilling the Weak Singularity condition 4.2 with this element $[\kappa/2]$ and no such complex Fermi curve fulfilling the Weak Singularity condition 4.2 with the zero element of $\Lambda^*/2\Lambda^*$. The effective dual lattice of these Generalized Weierstraß curves 4.4 is equal to $\Lambda_{[\kappa/2]}^*$.*

Proof. Given a *Generalized Weierstraß curve* 4.4 $\mathcal{F}$ with disconnected normalization, whose *first integral* is equal to 8π , let $\tilde{\mathcal{F}}_{\text{normal}}^\pm$ denote the components of the normalization, which contain $\infty^\pm$. Due to the properties of the restriction of the multi-valued function k to $\tilde{\mathcal{F}}_{\text{normal}}^+$, this compact Riemann surface is the spectral curve of an elliptic solution of the Kadomtsev–Petviashvili Equation. More precisely, on this compact Riemann surface there exists a *Baker–Akhiezer* function corresponding to these solutions, and the corresponding elliptic curve is the unique elliptic curve, which is conformally equivalent to the flat torus $\mathbb{R}^2/\Lambda$. In [Kr-2, Theorem 4.] these spectral curves are constructed as N -sheeted coverings of the corresponding elliptic curves. Due to the general construction of the algebraic-geometric solutions [Kr-1, D-K-N] the space, on which these solutions are defined, is mapped linearly into the Jacobian variety of the corresponding spectral curve. Hence for elliptic solutions the elliptic curve is mapped linearly into the corresponding Jacobian variety. The dual map [L-B, Chapter 2 §4] is a map from the Jacobian variety onto the elliptic curve. The composition of the canonical map from the spectral curve into the Jacobian variety with this map makes the spectral curve of an elliptic solutions in a canonical way to a finite-sheeted covering of the corresponding elliptic curve. In the present situation the function $k_1 + \sqrt{-1}k_2$ yields this canonical covering (compare with Lemma 4.16). Due to Lemma 4.16 the *first integrals* of the corresponding *complex Fermi curves* are equal to $4N\pi$. Hence in the present situation N is equal to 2. Due to [Kr-2, Lemma 2.] the genus of $\tilde{\mathcal{F}}_{\text{normal}}^+$ is therefore not larger than 2 (in [Kr-2, Lemma 2.] the *arithmetic genus* of the spectral curve is shown to be equal to N).

Since the *complex Fermi curve* is assumed to obey the *Weak Singularity condition* 4.2, the connected component $\tilde{\mathcal{F}}_{\text{normal}}^+$ has a singularity. Due to Proposition 3.7 there exist families of deformations of these *complex Fermi curves* within the set of *complex Fermi curves* with

disconnected normalization, whose *first integral* is equal to 8π , such that this singularity is removed. The components of the normalizations of these families, which contain ∞^+ , correspond to these elliptic solutions of the Kadomtsev–Petviashvili Equation. We conclude that the genus of $\bar{\mathcal{F}}_{\text{normal}}^+$ is at most equal to 1. On the other hand, the *first integral* of the unique *complex Fermi curve*, whose normalization has two connected components of genus zero, is zero. Therefore, $\bar{\mathcal{F}}_{\text{normal}}^+$ has genus one and due to the Riemann–Hurwitz formula [Fo, 17.14] it is an unbranched two-sheeted covering of the unique elliptic curve, which is conformally equivalent to $\mathbb{R}/\Lambda$. This implies that the effective dual lattice has to be a proper sublattice and that $\Lambda^*/\Lambda_{\text{effective}}^*$ is isomorphic to $\mathbb{Z}_2$. We conclude that the *complex Fermi curve* is one of the three *Generalized Weierstraß curves* 4.4 constructed above. **q.e.d.**

With the help of Lemma 4.13 we may construct a whole family of *Generalized Weierstraß curves* 4.4, whose *first integrals* are not larger than 8π . In this lemma we constructed a family of *complex Fermi curves* parameterized by the corresponding values of the *first integral* $\mathcal{W} \in [0, 4\pi]$. For all non-zero elements $[\kappa/2] \in \Lambda^*/2\Lambda^*$ let

$$[0, 8\pi] \rightarrow \bar{\mathcal{M}}_{3,\Lambda,\eta,\sigma,8\pi}, \mathcal{W} \mapsto \mathcal{F}_{\min,[\kappa/2]}(\mathcal{W}/2)$$

denote this family of *complex Fermi curves* with lattice $\Lambda_{[\kappa/2]} \subset \Lambda$ considered as a family of *complex Fermi curves* with lattice Λ . Since the *first integrals* of these *complex Fermi curves*, considered as a *complex Fermi curve* with lattice Λ , are twice the *first integrals* of these *complex Fermi curves*, considered as *complex Fermi curves* with lattice $\Lambda_{[\kappa/2]}$, the parameter $\mathcal{W}$ is equal to the *first integral* of the corresponding *complex Fermi curves*. From the proof of Lemma 4.13 we conclude that there exists a unique $\mathcal{W}_{[\kappa/2]}$, such that the *complex Fermi curve* $\mathcal{F}_{\min,[\kappa/2]}(\mathcal{W}/2)$ considered as a *complex Fermi curve* with lattice $\Lambda_{[\kappa/2]}$ contains both elements in the preimage of $[\kappa/2]$ under the natural map $\Lambda_{[\kappa/2]}^*/2\Lambda_{[\kappa/2]}^* \rightarrow \Lambda^*/2\Lambda^*$, if and only if $\mathcal{W} \geq \mathcal{W}_{[\kappa/2]}$. For all non-zero elements $[\kappa/2] \in \Lambda^*/2\Lambda^*$ this yields a family of *complex Fermi curves* fulfilling the *Weak Singularity condition* 4.2 with the element $[\kappa/2]$ parameterized by the corresponding *first integrals* $\mathcal{W} \in [\mathcal{W}_{[\kappa/2]}, 8\pi]$. In the proof of Lemma 4.13 we described the singularities of the family $\mathcal{F}_{\min,[\kappa/2]}(\mathcal{W})$ with $\mathcal{W} \in [0, 4\pi]$, considered as a family of *complex Fermi curves* with lattice $\Lambda_{[\kappa/2]}$. But besides these singularities this family also contains singularities, which are not singularities of the same family considered as a family of *complex Fermi curves* with lattice $\Lambda_{[\kappa/2]}$. More precisely, we have seen above that the subvarieties in $\mathbb{C}^2/\Lambda^*$ of this family considered as a family of *complex Fermi curves* with lattice Λ are one-sheeted coverings of the subvarieties in $\mathbb{C}^2/\Lambda_{[\kappa/2]}^*$ of this family considered as a family of *complex Fermi curves* with lattice $\Lambda_{[\kappa/2]}$. Those elements of the corresponding *complex Fermi curves* with lattice Λ , where this map is not biholomorphic, are exactly the singularities of these *complex Fermi curves*, which are not singularities of the corresponding *complex Fermi curves* with lattice $\Lambda_{[\kappa/2]}$.

Lemma 4.19. *For all $\mathcal{W} \in [\mathcal{W}_{[\kappa/2]}, 8\pi]$ the complex Fermi curve $\mathcal{F}_{\min,[\kappa/2]}(\mathcal{W}/2)$ has besides the singularity of the Weak Singularity condition 4.2 and the singularities of this complex Fermi curve considered as a complex Fermi curve with lattice $\Lambda_{[\kappa/2]}$ only ordinary double points of the form $(y, \eta(y))$, where the function dk_2/dk_1 has positive imaginary part at those points of these double points, which belong to $\bar{\mathcal{F}}_{\text{normal}}^+$, and no other singularities.*

Proof. Let $\mathcal{F}(t)$ be a continuous family of *complex Fermi curves* parameterized by $t \in (-\varepsilon, \varepsilon)$, which for $t < 0$ has a continuous family of double points of the form $(y(t), \eta(y(t)))$ with the property that the imaginary part of dk_2/dk_1 at $y(t)$ is positive. Moreover, let us assume in addition that for $t = 0$ the imaginary part of this function at $y(0)$ is zero, and that in some neighbourhood of $(y(0), \eta(y(0)))$ the *Local contributions to the arithmetic genus* 4.3 does not depend on t . Then at least for small $t < 0$ there exists another continuous family of double points $(y'(t), \eta(y'(t)))$, with the property that the imaginary part of dk_2/dk_1 at $y'(t)$ is negative, and the limit $t \rightarrow 0$ of $y(t)$ and $y'(t)$ coincide. In fact, all elements y , at which the real part of the function k takes some value in $\Lambda^*/2\Lambda^*$, belong to double points of the form $(y, \eta(y))$. The two form $d\Re(k_1) \wedge d\Re(k_2)$ vanishes on the normalization, if the function dk_2/dk_1 is real. Besides these zeroes all other zeroes of $d\Re(k_1) \wedge d\Re(k_2)$ on the normalization are *multiple points*. The former have real co-dimension one and the latter real co-dimension two. Hence the image under the map $\Re(k)$ in $\mathbb{R}^2$ is locally a finite-sheeted covering. Furthermore, the lines, where the function dk_2/dk_1 takes real values, are boundaries of these coverings, where two different sheets with opposite orientation meet each other.

Now let us consider some continuous family $\mathcal{F}(t)$ of *complex Fermi curves*, whose real part divides the normalization into two connected components. Moreover, the *complex Fermi curve* $\mathcal{F}(0)$ should have the property, that all singularities are double points of the form $(y, \eta(y))$, where dk_2/dk_1 has positive imaginary part at the point of $(y, \eta(y))$ contained in the component of ∞^+ . Then this property is conserved for all $0 \leq t$ as long as the real part of $\mathcal{F}(t)$ does not contain an element of $\Lambda^*/2\Lambda^*$. In fact, due to the assumption these double points can loose this property only if they meet the boundary between $\bar{\mathcal{F}}_{\text{normal}}^+$ and $\bar{\mathcal{F}}_{\text{normal}}^-$, which is the real part.

We distinguished between singularities of these *complex Fermi curves* considered as *complex Fermi curves* with lattice $\Lambda_{[\kappa/2]}$ and singularities, which are not singularities of these *complex Fermi curves* considered as *complex Fermi curves* with lattice $\Lambda_{[\kappa/2]}$. Ordinary double points of the form $(y, \eta(y))$ belong to the first class, if at these points the real part of the function k takes values in the image of the canonical map $\Lambda_{[\kappa/2]}^*/2\Lambda_{[\kappa/2]}^* \rightarrow \Lambda^*/2\Lambda^*$, and otherwise to the second class. From the above considerations it follows, that it suffices to show that for one *complex Fermi curve* of this family the statement of the lemma is true and that the real part of this family does not contain an element in $\Lambda^*/2\Lambda^*$, which does not belong to the image of the canonical map $\Lambda_{[\kappa/2]}^*/2\Lambda_{[\kappa/2]}^* \rightarrow \Lambda^*/2\Lambda^*$.

We first consider the rectangular lattice $\Lambda = \mathbb{Z}^2 \subset \mathbb{R}^2$, and the family corresponding to the non-zero element $[(1/2, 1/2)] \in \Lambda^*/2\Lambda^*$. The corresponding lattice $\Lambda_{[(1/2, 1/2)]}$ is generated by the elements $(1/2, 1/2)$ and $(1/2, -1/2)$, and the corresponding dual lattice $\Lambda_{[(1/2, 1/2)]}^*$ by the elements $(1, 1)$ and $(1, -1)$. It is quite easy to see (and we shall do this at the end of this section), that in this case $\mathcal{W}_{[(1/2, 1/2)]}$ is equal to $2\pi^2$, and that the corresponding *complex Fermi curve* may be described by the equation $g(k, k) = 1/2$. An easy calculation shows that besides the singularity of the *Weak Singularity condition* 4.2 all singularities of this *complex Fermi curve* are ordinary double points of the form $(y, \eta(y))$, where the imaginary part of the function dk_2/dk_1 is positive at those elements of these double points, which belong to $\bar{\mathcal{F}}_{\text{normal}}^+$. The real part of this *complex Fermi curve* is a circle of radius $1/\sqrt{2}$. From Lemma 4.13 we conclude that all real parts of the family $\mathcal{F}_{\min, [(1/2, 1/2)]}(\mathcal{W}/2)$ with $\mathcal{W} \in (2\pi^2, 8\pi]$ cover

the complement in the $\mathbb{R}^2/\Lambda_{[(1/2,1/2)]}^*$ of the closed disc around the origin with radius $1/\sqrt{2}$. Obviously, this complement contains only one element of $\Lambda^*/2\Lambda_{[(1/2,1/2)]}^*$, which belongs to the kernel of the canonical map $\Lambda^*/2\Lambda_{[(1/2,1/2)]}^* \rightarrow \Lambda^*/2\Lambda^*$. This shows that for the rectangular class and the family corresponding to the element $[(1/2,1/2)] \in \Lambda^*/2\Lambda^*$ the statement of the lemma is true.

Now we consider all these families as one family parameterized by the *first integral* and the conformal classes. It is quite easy to see, that the groups $\Lambda^*/2\Lambda^*$ build a non-trivial bundle over the moduli space of conformal classes (compare with $\mathcal{M}_1$ at the end of this section). More precisely, the non-zero elements of this group fit to a connected three-sheeted covering over this moduli space, and the zero element to a one-sheeted covering. This follows from the explicit calculation of the conformal equivalence classes of the lattices $\Lambda_{[\kappa/2]}$ in dependence of the conformal equivalence classes of Λ given at the end of this section. We conclude that this family is connected. With the above considerations it suffices to show that the real part of all elements of this family does not contain any element of $\Lambda^*/2\Lambda^*$, which is not contained in the image of the canonical map $\Lambda^*/2\Lambda_{[(1/2,1/2)]}^* \rightarrow \Lambda^*/2\Lambda^*$. In order to prove this we restrict this family again to one conformal class and some non-zero element $[\kappa/2] \in \Lambda^*/2\Lambda^*$. Due to the proof of Lemma 4.13 the real parts of all these families $\mathcal{F}_{\min, [\kappa/2]}(\mathcal{W})$, with $\mathcal{W} \in [0, 4\pi]$ meet every element of $\mathbb{R}^2/\Lambda^*$ exactly twice. In particular, each of the two elements of $\Lambda^*/2\Lambda^*$, which are not contained in the image of the canonical map $\Lambda^*/2\Lambda_{[\kappa/2]}^* \rightarrow \Lambda^*/2\Lambda^*$, is met twice. However, the singularity of the *Weak Singularity condition* 4.2 always includes four double points, which are not double points of any *Generalized Weierstraß curve* 4.4 of this family considered as a *complex Fermi curve* with lattice $\Lambda_{[\kappa/2]}$. We conclude that the real parts of the family $\mathcal{F}_{\min, [\kappa/2]}(\mathcal{W}/2)$ meet all those elements of $\Lambda^*/2\Lambda^*$, which are not contained in the image of the canonical map $\Lambda^*/2\Lambda_{[\kappa/2]}^* \rightarrow \Lambda^*/2\Lambda^*$ for values of $\mathcal{W}$ in $[0, \mathcal{W}_{[\kappa/2]})$, and not for values in $[\mathcal{W}_{[\kappa/2]}, 8\pi]$. This completes the proof. **q.e.d.**

Now we can classify all relative minimizers of the *Generalized Willmore functional* 4.5, whose *first integrals* are not larger than 8π .

Proposition 4.20. *The restriction of the first integral to the space of all complex Fermi curves, which obey the Weak Singularity condition 4.2 for some element $[\kappa/2] \in \Lambda^*/2\Lambda^*$ has exactly one relative minimizer, whose first integral is not larger than 8π , if $[\kappa/2]$ is not zero, and otherwise no such relative minimizer. The complex Fermi curves of these relative minimizers are $\mathcal{F}_{\min, [\kappa/2]}(\mathcal{W}_{[\kappa/2]}/2)$. In particular, the corresponding values of the first integral are equal to $\mathcal{W}_{[\kappa/2]}$.*

As a preparation for the proof of this Proposition we first give the following

Lemma 4.21. *Let Y be a compact connected Riemann surface with some anti-holomorphic involution ρ , such that the real part $Y_{\mathbb{R}}$ (= the set of fixed points of ρ) divides Y into two connected components Y^+ and Y^- with $\partial Y^+ = \partial Y^- = Y_{\mathbb{R}}$. Let $y^+ \in Y^+$ and $y^- \in Y^-$ be two different points, which are interchanged by the involution ρ . Then there exists a meromorphic real (i. e. $\rho^*\omega = \bar{\omega}$) differential form ω , which has poles of first-order at y^+ and y^- and no other poles, and is positive on the real part, with respect to the orientation induced from the orientation of Y^+ on the boundary $Y_{\mathbb{R}} = \partial Y^+$.*

Proof. Let $\Omega_{y^++y^-}$ be the space of real meromorphic forms with a pole of at most first-order at y^+ and y^- and with no other poles. We shall first prove that for any choice of finitely many points $y_1, \dots, y_n$ of $Y_{\mathbb{R}}$ there exists some $\omega \in \Omega_{y^++y^-}$, which is positive at these points with respect to the given orientation. The values of all elements of $\Omega_{y^++y^-}$ at these points in terms of local real positive holomorphic forms $dz_1, \dots, dz_n$ form a linear subspace V of $\mathbb{R}^n$. If this subspace contains no element v , all whose components are positive, then the orthogonal complement (with respect to the Euclidean scalar product) of this subspace contains some non zero element $c \in \mathbb{R}^n$, all whose components are non-negative. In fact, due to the Separating hyperplane theorem [R-S-I, Theorem V.4] there exists a linear functional on $\mathbb{R}^n$, whose values on the open convex cone of all elements, all whose components are positive, are larger or equal to the values on V . This functional may be written as the Euclidean scalar product with some element $c \in \mathbb{R}^n$, which has the desired properties. This element c in the orthogonal complement of V defines some Mittag-Leffler distribution, and due to [Fo, Theorem 18.2] this Mittag-Leffler distribution has a global solution f . Obviously $\rho^* \bar{f}$ solves the same Mittag-Leffler distribution. So let us assume that f is real (i. e. $f = \rho^* \bar{f}$). Since all entries of c are non-negative the restriction of the function f to any connected component of $Y_{\mathbb{R}}$ takes all real values with the same multiplicity. But $\Omega_{y^++y^-}$ contains besides the real holomorphic forms also meromorphic forms with poles of at most first-order at y^+ and y^- . The same arguments as in the proof of [Fo, Theorem 18.1 and 18.2] show, that in this case f has a zero at y^+ and y^- , which is impossible. Hence there exists some element $\omega \in \Omega_{y^++y^-}$, which is positive at $y_1, \dots, y_n$.

Now we choose any norm on $\Omega_{y^++y^-}$ (they are all equivalent, since $\Omega_{y^++y^-}$ is a finite-dimensional vector space). The subspace of $\Omega_{y^++y^-}$ consisting of forms, whose norm is equal to 1 is compact. All subsets of this compact space of forms, which are non-negative at finitely many points of $Y_{\mathbb{R}}$, are closed. Moreover, due to our first claim the intersection of finitely many of such subsets is non-empty. Hence the intersection of all these subsets is also non-empty. Hence there exists some form $\omega \in \Omega_{y^++y^-} \setminus \{0\}$, which is non-negative on $Y_{\mathbb{R}}$. This form can have only finitely many zeroes. If we add some small multiple of an element of $\Omega_{y^++y^-}$, which is positive at all the zeroes of ω contained in $Y_{\mathbb{R}}$, we obtain an element of $\Omega_{y^++y^-}$, which is positive on $Y_{\mathbb{R}}$. **q.e.d.**

Proof of Proposition 4.20. Let us first prove that we may deform all relative minimizers of the restriction of the *first integral* to the space of *Generalized Weierstraß curves* 4.4 into a *Generalized Weierstraß curve* 4.4, whose normalization has two connected components. Let us consider the class of *complex Fermi curves*, which obey conditions (i)–(iv) of Corollary 4.12. Since the complement of the image of the map described in (ii) are disjoint subsets, the total area of this complement is smaller than the volume of $\mathbb{R}^2/\Lambda^*$. Therefore, due to Lemma 3.16, on these *complex Fermi curves* the *first integral* takes values in the interval $(4(d-1)\pi, 4d\pi]$, where d denotes the number of sheets in (ii). Since we are interested in *complex Fermi curves*, whose *first integral* belongs to $(4\pi, 8\pi]$, in our case this number of sheets is equal to two. Due to Corollary 4.12 we may deform all relative minimizers of the restriction of the *first integral* to the space of *Generalized Weierstraß curves* 4.4 by an arbitrary small deformation into a *complex Fermi curve* within this class.

Now we claim that we may deform any element of this class within this class into a *complex Fermi curve*, with disconnected normalization. Due to Proposition 3.7 and Lemma 4.21 we may deform the *complex Fermi curves* within this class in such a way that the disjoint closed subsets mentioned in (ii) become smaller and smaller, if the dividing real parts of the *complex Fermi curves* do not have *cusps*. Due to Proposition 3.9 and the remarks following this proposition this condition is necessary in order to preserve condition (i). Hence we may shrink these pieces mentioned in (ii) until the boundary of these pieces intersect each other. In this case the real part of the deformed *complex Fermi curve* has some multiple points. Either they are double points of the form $(y, \sigma(y))$, or they are pairs of double points of the form (y, y') and $\sigma(y), \sigma(y')$. In the second case the deformations of Proposition 3.7 allow to split these pieces into several pieces and shrink them further. In the first case the deformations of *Orbits of type 1* into *Orbits of type 2* constructed in the proof of Proposition 3.7 concerning *Multiple points O1/O2* and *Multiple points O1/O1* have the same effect. Due to Corollary 3.54 the space of all *complex Fermi curves* with dividing real part, whose *first integral* is not larger than 8π is compact and contained in $\bar{\mathcal{M}}_{g_{\max}, \Lambda, \eta, \sigma, 8\pi}$. Furthermore, the proof of this lemma actually shows that at all zeroes of $d\hat{p}$ and $d\check{p}$ the imaginary part of k is bounded. Hence the formula for the *first integral* in Lemma 3.16 shows that the *first integral* is continuous on this space, in contrast to the lower-continuity on $\bar{\mathcal{M}}_{g, \Lambda, \eta}$. Hence we may shrink the complement of the image of the map described in (ii) until it is empty. The normalization of the corresponding *complex Fermi curve* has two connected components. Since we have started with a *complex Fermi curve*, whose *first integral* is smaller than 8π the final *complex Fermi curve* of this deformation is one of the three *Generalized Weierstraß curves* 4.4 described in Lemma 4.18. Obviously properties (i)–(ii) are preserved under these deformations.

It remains to show that also conditions (iii)–(iv) are preserved, and that the *complex Fermi curves* within this class do not have real cusps. The considerations in the proof of Lemma 4.19 concerning ordinary double points of the form $(y, \eta(y))$ imply that condition (iii) is preserved under these deformations. Due to deformations of *Multiple points O2/O2* and *Multiple points O3* constructed in the proof of Theorem 4.9 we may always deform by some arbitrary small deformation an *Orbit of type 3* into two *Orbits of type 2*, and this deformation may be chosen within this class. Due to Lemma 4.21 this deformation may be chosen in such a way, that these pieces never contain the element corresponding to $[\kappa/2]$. Hence the two *Orbits of type 2* in the normalizations of this family can only be deformed into an *Orbit of type 3* over $[\kappa/2]$. In this case again the deformations of *Multiple points O2/O2* and *Multiple points O3* constructed in the proof of Theorem 4.9 show that the deformation can be continued by deforming this *Orbit of type 3* back into two *Orbits of type 2*. Therefore, condition (iv) is preserved.

Finally, a real *cusp* contradicts conditions (i)–(iii) with $\mathcal{W} \in (4\pi, 8\pi]$. In fact, if both differentials $d\hat{p}$ and $d\check{p}$ have zeroes of at least second order, then at least three sheets of the function $k_1 + \sqrt{-1}k_2$ have a branch point at the real *cusp*. Obviously this contradicts condition (ii) for two-sheeted coverings ($\iff \mathcal{W} \in (4\pi, 8\pi]$). Furthermore, all other real *cusps* are locally of the form $\check{p}^2 = \hat{p}^{2m+1}$. All deformations of such *cusps* contain either no singularities nearby the real *cusp*, or double points of the form $(y, \rho(y))$, or ordinary real double points, or real *cusps* of the same form of less order (corresponding to smaller m).

In the first case due to condition (i) all zeroes of $d\hat{p}$ nearby the deformed real *cusp* have to be real, and condition (ii) is violated. The second case contradicts condition (iii), and the third case condition (ii). Finally, the fourth case is excluded by induction. This shows that we may deform all relative minima within the class of *complex Fermi curves* obeying conditions (i)–(iv) into a *Generalized Weierstraß curve* 4.4 with disconnected normalization.

In a second step we claim that, if the number of sheets in condition (ii) is chosen to be equal to two, (in this case the *first integral* takes values in the interval $(4\pi, 8\pi]$), these classes are equal to the families $\mathcal{F}_{\min, [\kappa/2]}(\mathcal{W}/2)$, with $\mathcal{W} \in (W_{[\kappa/2]}, 8\pi]$. Due to the first claim it suffices to show that this family is the only possible deformation of the *Generalized Weierstraß curves* 4.4 described in Lemma 4.18 within this class. Due to Lemma 4.19 all elements of these family may not be deformed by a small deformation within the class of *complex Fermi curves* with dividing real parts into a *complex Fermi curve*, whose effective dual lattice is not contained in $\Lambda_{[\kappa/2]}^*$. Lemma 4.13 implies that all double points of these *complex Fermi curves* considered as *complex Fermi curves* with lattice $\Lambda_{[\kappa/2]}$ have only two types of singularities:

Ordinary double points. Ordinary double points of the form $(y, \eta(y))$, where the imaginary part of the function dk_2/dk_1 takes positive values at those points of these double points, which belong to $\bar{\mathcal{F}}_{\text{normal}}^+$.


Real double points. Double points of the form $(y, \eta(y))$, which belong to the real part. In this case the function dk_2/dk_1 takes the same real value at both elements of these double points.

More precisely, the *real double points* occur only for those *complex Fermi curves* of this family, which are described in Corollary 4.14. Moreover, due to the proof of Lemma 4.13 the only possible deformations of these *complex Fermi curves* within the class of *complex Fermi curves* obeying conditions (i)–(ii) is the family described in Lemma 4.13. This proves the second claim.

We conclude that the *complex Fermi curves* $\mathcal{F}_{\min, [\kappa/2]}(\mathcal{W}_{[\kappa/2]}/2)$ are the unique relative minimizers of the restriction of the *first integral* to the subset of all *complex Fermi curves* fulfilling the *Weak Singularity condition* 4.2 with the element $[\kappa/2] \in \Lambda^*/2\Lambda^*$. **q.e.d.**

Remark 4.22. For special conformal classes there exist real- σ -hyperelliptic Generalized Weierstraß curves 4.4 with dividing real parts and real cups, whose first integral belong to $(8\pi, 8\pi + \varepsilon)$. In fact, those conformal classes, for which the two unique real elements of $\mathcal{F}_{\min}(4\pi)$ belong to a small neighbourhood of an element in $\Lambda^*/2\Lambda^*$, have a complex Fermi curve with dividing real part, two Orbits of type 2 over two different non-trivial elements of $\Lambda^*/2\Lambda^*$ and a real cusp. The corresponding first integral belongs to $(4\pi, 4\pi + \varepsilon)$. This can be seen by continuation of the family $\mathcal{F}_{\min}(\mathcal{W})$ constructed in Lemma 4.13 along another path.

At the unique element of the family, where the real part has one Orbit of type 1 and two Orbits of type 2 over all non-trivial elements of $\Lambda^*/2\Lambda^*$ we increase the first integral along the unique path of real- σ -hyperelliptic complex Fermi curves, which have real multiple points (the deformed Orbits of type 1). Since the real part of these complex Fermi curves is a self-intersecting circle in $\mathbb{R}^2$, this family has to end at a real- σ -hyperelliptic complex Fermi curve with two real cusps. This is indicated in the neighbouring figure.



Remark 4.23. It is not difficult to see, that there exists a Generalized Weierstraß curve 4.4 with disconnected normalization, which corresponds to the trivial spin structure, and whose first integral is equal to 12π . Therefore, it seems to be possible to determine with similar methods the minimum of the first integral on the space of all Generalized Weierstraß curves 4.4 corresponding to the trivial spin structure.

Due to this proposition the absolute minimum of the *Generalized Willmore functional* 4.5 is equal to the minimum of $\mathcal{W}_{[\kappa/2]}$, where $[\kappa/2]$ runs over the non-zero element of $\Lambda^*/2\Lambda^*$. In order to determine these minima in dependence of the conformal classes, we have to combine the mappings $\Lambda \mapsto \Lambda_{[\kappa/2]}$ and the *complex Fermi curves* $\mathcal{F}_{\min}(\kappa/2)$ described in Corollary 4.14. More precisely, for any *complex Fermi curve* $\check{\mathcal{F}}_{\min}(\check{\kappa}/2)$ described in Corollary 4.14 with lattice $\check{\Lambda}$, and which contains another non-zero $[\check{\kappa}'/2] \in \check{\Lambda}^*/2\check{\Lambda}^*$, there exists a unique sublattice $\Lambda \subset \check{\Lambda}$ and some unique non-zero $[\kappa/2] \in \Lambda^*/2\Lambda^*$, which contains $[\check{\kappa}/2]$ and $[\check{\kappa}'/2]$ in the preimage under the canonical map $\check{\Lambda}^*/2\check{\Lambda}^* \rightarrow \Lambda^*/2\Lambda^*$. Despite some degenerate situations, this other element of $\Lambda^*/2\Lambda^*$ belongs to an *Orbit of type 2*. Consequently, for genus one this other element is unique, and for genus two there are two choices. The lattice $\check{\Lambda}$ is equal to $\Lambda_{[\kappa/2]}$. Moreover, the *complex Fermi curve* $\check{\mathcal{F}}_{\min}(\check{\kappa}/2)$ with lattice $\Lambda_{[\kappa/2]}$, considered as a *complex Fermi curve* with lattice Λ , minimizes the restriction of the *first integral* to the space of *complex Fermi curves* fulfilling the *Weak Singularity condition* 4.2 with the element $[\kappa/2]$. Furthermore, since the order of $\check{\Lambda}/\Lambda$ is two, the *first integral* of this minimizer is twice the *first integral* of $\check{\mathcal{F}}_{\min}([\kappa/2])$. Due to Corollary 4.14 all $[\check{\kappa}/2] \in \check{\Lambda}^*/2\check{\Lambda}^*$, whose *complex Fermi curve* $\check{\mathcal{F}}_{\min}(\check{\kappa}/2)$ have genus larger than zero, fulfill this condition. Moreover, if the genus is zero, but $\mathcal{F}_{\min}(\check{\kappa}/2)$ has two *Orbits of type 1* over two different elements of $\check{\Lambda}^*/2\check{\Lambda}^*$, this *complex Fermi curve* also fulfills this condition. But from Corollary 4.14 we conclude that if the genus of $\mathcal{F}_{\min}(\check{\kappa}/2)$ is larger than one, there exists another $[\check{\kappa}'/2] \in \check{\Lambda}^*/2\check{\Lambda}^*$, which fulfills this condition, and whose *first integral* is smaller. In order to determine the absolute minimizer of the *Generalized Willmore functional* 4.5 for all lattices, it suffices therefore to calculate all *complex Fermi curves* $\mathcal{F}_{\min}(\check{\kappa}/2)$ of genus at most one.

In order to do this we parameterize the space of all lattices $\Lambda \subset \mathbb{R}^2$. Any linear conformal transformation

$$A : \mathbb{R}^2 \rightarrow \mathbb{R}^2, x \mapsto Ax,$$

where A is a real 2×2 matrix with $A^t A = \det^2(A) \mathbf{1}$, induces a transformation of the lattice Λ into $A\Lambda = \{A\gamma \mid \gamma \in \Lambda\}$, the dual lattice Λ^* into $\det^{-2}(A)A^t\Lambda^* = \{\det^{-2}(A)A^t\kappa \mid \kappa \in \Lambda^*\}$ and the *complex Fermi curve* $\mathcal{F}$ into $\det^2(A)A^t\mathcal{F} = \{[\det^{-2}(A)A^t k] \in \mathbb{C}^2 / \det^{-2}(A)A^t\Lambda^* \mid [k] \in \mathcal{F}\}$. Since the *Weak Singularity condition* 4.2 and the *first integral* is invariant under

these transformations, we should parameterize the conformal equivalence classes of these lattices. It is convenient to use generators of the lattice Λ in order to parameterize this space. If $\hat{\gamma}$ and $\check{\gamma}$ are some generators of the lattice Λ with positive orientation, there exists a unique conformal transformation, which transforms $\hat{\gamma}$ into $(1, 0)$. The other generator $\check{\gamma}$ is parameterized by $\tau = \check{\gamma}_1 + \sqrt{-1}\check{\gamma}_2$. Since they have positive orientation, we conclude that $\Im(\tau) > 0$. Hence the upper half plane $\mathbb{C}^+ = \{\tau \in \mathbb{C} \mid \Im(\tau) > 0\}$ parameterizes the space of all conformal equivalence classes of lattices up to the choice of two generators with positive orientation. The modular group $SL(2, \mathbb{Z})$ describes the transformation of generators with positive orientation into each other. Hence the space of conformal equivalence classes of lattices is equal to the space of orbits of the action of the modular group on the upper-half plane. There exist many fundamental regions of this action, but the following choice is the most common:

$$\mathcal{M}_1 = \{\tau \in \mathbb{C}^+ \mid -1/2 \leq \Re(\tau) \leq 1/2 \text{ and } 1 \leq |\tau|\}.$$

More precisely, the modular group is generated by the elements $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. The corresponding action on the upper-half plane is given by $S : \tau \mapsto -1/\tau$ and $T : \tau \mapsto \tau + 1$. The first transformation induces an identification of that part of the boundary of $\mathcal{M}_1$, which belongs to the unit circle, with itself, and the second induces an identification of that part of the boundary, which belongs to the line $\Re(\tau) = -1/2$, with the part of the boundary, which belongs to the line $\Re(\tau) = 1/2$. The corresponding dual lattice is generated by the elements $\hat{\kappa}_1 + \sqrt{-1}\hat{\kappa}_2 = -\sqrt{-1}\tau/\Im(\tau)$ and $\check{\kappa}_1 + \sqrt{-1}\check{\kappa}_2 = \sqrt{-1}/\Im(\tau)$. In the sequel we will denote the natural choice of generators corresponding to these moduli τ by $\hat{\gamma}_\tau = (1, 0)$ and $\check{\gamma}_\tau = (\Re(\tau), \Im(\tau))$. Analogously the corresponding generators of the dual lattice are denoted by $\hat{\kappa}_\tau = (1, -\Re(\tau)/\Im(\tau))$ and $\check{\kappa}_\tau = (0, 1/\Im(\tau))$.

All $\mathcal{F}_{\min}(\kappa/2)$ of genus zero: If Y is a *complex Fermi curve* of genus zero, both components of k extend to single-valued meromorphic functions with first-order poles at ∞^- and ∞^+ . If Y is invariant under the involution σ , the function $g(k, k)$, extends to a global holomorphic function and therefore has to be constant. Since Y is invariant under the involution $\eta : k \mapsto -\bar{k}$ without fixed points, this constant has to be a non-negative real number. The *first integral* $\mathcal{W}$ is equal to $4\pi^2 \text{vol}(\mathbb{R}^2/\Lambda)$ times this constant, and the *complex Fermi curve* is given by the equation $g(k, k) = \mathcal{W}/(4\pi^2 \text{vol}(\mathbb{R}^2/\Lambda))$. Strictly speaking, the *complex Fermi curve* is the union of the solutions of this equation shifted by all $\kappa \in \Lambda^*$. We conclude from Corollary 4.14 that we may obtain all $\mathcal{F}_{\min}(\kappa/2)$, by finding a non-zero dual lattice vector $\kappa \in \Lambda^*$ with the smallest length, and the corresponding *complex Fermi curve* is given by the equation $g(k, k) = g(\kappa, \kappa)/4$. Obviously, for all lattices corresponding to $\tau \in \mathcal{M}_1$ the second generator $\check{\kappa}_\tau$ has minimal length among all non-zero dual lattice vectors. However, if $|\tau| = 1$, then the first generator $\hat{\kappa}_\tau$ has the same length. We conclude that if $|\tau| = 1$, then both *complex Fermi curves* $\mathcal{F}_{\min}(\hat{\kappa}_\tau/2)$ and $\mathcal{F}_{\min}(\check{\kappa}_\tau/2)$ coincide and are given by the equation $g(k, k) = 1/(4\Im^2(\tau))$. In all other cases only $\mathcal{F}_{\min}(\check{\kappa}_\tau/2)$ is given by this equation. Furthermore, these are all *complex Fermi curves* of genus zero of the form $\mathcal{F}_{\min}(\kappa/2)$. The corresponding values of the *first integral* are equal to $\pi^2/\Im(\tau)$. This implies that there exists some $\mathcal{F}_{\min}(\kappa/2)$ of genus zero, which contains some other element of $\Lambda^*/2\Lambda^*$, if and only if $|\tau| = 1$ and

κ is either $\hat{\kappa}_\tau$ or $\check{\kappa}_\tau$. In this case the *first integral* is equal to $\pi^2/\Im(\tau)$. In particular, the minimum of the *first integral* of these cases is attained for $\tau = (0, 1)$ and the corresponding value of the *first integral* is equal to π^2 . Therefore, the lowest critical value of the *Generalized Willmore functional* 4.5 is at most $2\pi^2$. Let us now see that this is in fact the absolute minimum.

All $\mathcal{F}_{\min}(\kappa/2)$ of genus one: All these *complex Fermi curves* are *real σ -hyperelliptic* and have exactly one *Orbit of type 2* over the element $[\check{\kappa}_\tau/2] \in \Lambda^*/2\Lambda^*$. Moreover, $\infty^\pm$ are two other fixed points of the involution σ and these four fixed points are all fixed points of the involution σ , since due to the Hurwitz formula (see e.g. [Fo, 17.14]) the involution σ has four fixed points. Moreover, there has to exist an *Orbit of type 1* over some other element of $\Lambda^*/2\Lambda^*$. Therefore, let us first determine the moduli space of all elliptic Riemann surfaces with two commuting involutions, one of which is holomorphic (σ) and the other is anti-holomorphic (ρ), whose composition has no fixed points, and besides the *Orbits of type 2* over $[\check{\kappa}_\tau/2]$ and the points $\infty^\pm$ one *Orbit of type 1*. We describe any such Riemann surface as the quotient of the complex plane $\{z \mid z \in \mathbb{C}\}$ by some two-dimensional lattice. Since all translations induce some biholomorphic map of the Riemann surface into itself, we may chose the origin to correspond to the point ∞^+ . Hence the involution σ is induced by the map $z \mapsto -z$. Since the Riemann surface has *dividing real part*, one generator of $H_1(Y, \mathbb{Z})$ may be chosen to be invariant under ρ and the other to be anti-invariant. These two generators of $H_1(Y, \mathbb{Z})$ correspond to generators of the lattice. By multiplication by some non-zero complex number we may always achieve that the invariant generator is purely imaginary, the anti-invariant generator is equal to 1, and ρ is induced by the map $z \mapsto -\bar{z} + \text{some constant}$. Since ρ and σ commute, twice this constant has to be a period, and the condition that $\rho\sigma$ has no fixed points implies that this constant has to be $1/2$ times the real generator. Hence the lattice is generated by a real positive period usually denoted by 2ω (which may be chosen to be 1) and an imaginary period usually denoted by $2\omega'$, which may be chosen to have positive imaginary part. The two involutions are induced by the maps $\sigma : z \mapsto -z$ and $\rho : z \mapsto -\bar{z} + \omega$. This implies that ∞^- corresponds to the point ω , and the *Orbit of type 2* over $[\check{\kappa}_\tau/2]$ corresponds to the points ω' and $\omega + \omega'$. The real part corresponds to the lines $\Re(z) = \pm\omega/2$. Therefore, the *Orbit of type 1* corresponds to two points $z_1 \in \omega/2 + \sqrt{-1}\mathbb{R}/(2\omega'\mathbb{Z})$ and $z_2 = -z_1$. We conclude that the moduli space of such Riemann surfaces is given by the quotient of the two generators $\omega'/\omega \in \sqrt{-1}\mathbb{R}^+$ and some element z_1 of $\omega/2 + \sqrt{-1}\mathbb{R}/(2\omega'\mathbb{Z})$. This is a real two-dimensional manifold.

The integral of dk along all cycles of $H_1(Y, \mathbb{Z})$, which are anti-invariant under ρ has to vanish. Therefore the lattice $\Lambda_{\text{effective}}^*$ is at most one-dimensional. On the other hand, since Y has an *Orbit of type 2* over $[\kappa'/2]$, there exist two generators $\hat{\kappa}$ and $\check{\kappa}$ of Λ^* , such that $\Lambda_{\text{effective}}^* = \check{N}\check{\kappa}\mathbb{Z}$, with some odd number $\check{N}$. Apriori these two generators are not necessarily the same as the generators $\hat{\kappa}_\tau$ and $\check{\kappa}_\tau$, but we will see that they indeed are. If $\hat{\gamma}$ and $\check{\gamma}$ are the dual generators of Λ , then the function $\hat{p} = g(\hat{\gamma}, k)$ extends to a single-valued meromorphic function with poles only at 0 and ω . The properties $\sigma^*k = -k$ and $\rho^*\bar{k} = k$ implies that $\hat{p}$ may be expressed by the Weierstraß functions

(see e.g. [Bat, 13.12])

$$\hat{p} = \alpha (\zeta(z) - \zeta(z + \omega) + \eta) = \frac{-\alpha}{2} \frac{\wp'(z)}{\wp(z) - e_1},$$

with some non-zero real number α . Since z_1 and $-z_1$ correspond to an *Orbit of type 1* over some non-zero element of $\Lambda^*/2\Lambda^*$ different from $[\tilde{\kappa}/2]$, we conclude that α is given by

$$\alpha = \frac{\hat{N}}{2(\zeta(z_1) - \zeta(z_1 + \omega) + \eta)} = -\hat{N} \frac{\wp(z_1) - e_1}{\wp'(z_1)},$$

with some odd number $\hat{N}$. The conditions on the function $\check{p}$ imply that it may be expressed by the Weierstraß functions:

$$\check{p} = \frac{\check{N}}{2\pi\sqrt{-1}} (2\eta z - \omega(\zeta(z) + \zeta(z + \omega) - \eta)) + \beta p,$$

with some real β . Again the condition that z_1 and $-z_1$ correspond to an *Orbit of type 1* over some non-zero element of $\Lambda^*/2\Lambda^*$ different from $[\tilde{\kappa}/2]$ implies that β is given by

$$\beta = -\frac{\check{N}}{\hat{N}\pi\sqrt{-1}} (2\eta z_1 - \omega(\zeta(z_1) + \zeta(z_1 + \omega) - \eta)) + N,$$

with some integer N . At the point ∞^+ the function $\check{p}/\hat{p}$ takes the value τ . Hence the parameters τ of the conformal equivalence classes of the lattice are given by

$$\begin{aligned} \Re\tau &= \beta = -\frac{\check{N}}{\hat{N}\pi\sqrt{-1}} (2\eta z_1 - \omega(\zeta(z_1) + \zeta(z_1 + \omega) - \eta)) + N \\ \Im\tau &= -\frac{\check{N}\omega}{2\hat{N}\pi} \frac{\wp'(z_1)}{\wp(z_1) - e_1} \end{aligned}$$

Finally, due to Lemma 3.12 the *first integral* is equal to the residue of the form $8\pi^2\sqrt{-1} \text{vol}(\mathbb{R}^2/\Lambda) k_2 dk_1 = 8\pi^2\sqrt{-1} \check{p} d\hat{p}$ at ∞^+ and therefore equal to

$$\mathcal{W} = 8\pi\alpha\check{N}(\eta + e_1\omega) = -8\pi\hat{N}\check{N} \frac{\wp(z_1) - e_1}{\wp'(z_1)} (\eta + e_1\omega).$$

The function $\frac{\wp'(z)}{\wp(z) - e_1}$ is real on the lines $\Im(z) = 0, \Im(\omega')$ and on the lines $\Re(z) = \pm\omega/2$. In particular, $d\hat{p}$ has four zeroes at the points $\pm\omega/2$ and $\pm\omega/2 + \omega'$. Let us calculate the values of this function at these four points. The square of this function is a rational function of $\wp$

$$\left(\frac{\wp'(z)}{\wp(z) - e_1} \right)^2 = 4 \frac{(\wp(z) - e_2)(\wp(z) - e_3)}{\wp(z) - e_1}.$$

Therefore, at the zeroes of $d\hat{p}$ $\wp$ takes the values $(\wp - e_1)^2 = 2e_1^2 + e_2e_3$ and more precisely $\wp(\pm\omega/2) = e_1 + \sqrt{2e_1^2 + e_2e_3}$ and $\wp(\pm\omega/2 + \omega') = e_1 - \sqrt{2e_1^2 + e_2e_3}$. Also the

function $-\frac{\wp'(z)}{\wp(z)-e_1}$ takes at the points $\pm\omega/2$ the values $\pm\sqrt{\frac{3e_1\sqrt{2e_1^2+e_2e_3}+(4e_1^2+e_2e_3)}{\sqrt{2e_1^2+e_2e_3}}}$ and at the points $\pm\omega/2+\omega'$ the values $\pm\sqrt{\frac{3e_1\sqrt{2e_1^2+e_2e_3}-(4e_1^2+e_2e_3)}{\sqrt{2e_1^2+e_2e_3}}}$. If $\omega'/\omega \in \sqrt{-1}\mathbb{R}^+$, then the numbers e_1, e_2 and e_3 satisfy the inequalities $e_3 \leq e_2 \leq e_1$. Hence the restriction of the function $-\frac{\wp'(z)}{\wp(z)-e_1}$ to $\Re(z) = \omega/2$ is positive and takes the maximum at $\omega/2$ and the minimum at $\omega/2+\omega'$. Also the restriction of this function to $\Re(z) = -\omega/2$ is negative and takes the minimum at $-\omega/2$ and the maximum at $-\omega/2+\omega'$. By definition of $\mathcal{F}_{\min}(\kappa/2)$ they have to be minimizers of the *first integral*. This implies that $\hat{N} = 1$ and $\check{N} = 1$. On the other hand the function x is equal to N if $z_1 = \omega/2$ and equal to $1+N$ if $z_1 = \omega/2+\omega'$.

With the help of the limits of the Weierstraß functions (see e.g. [Bat, 13.15]) it is easy to see that the limit $\omega'/\omega \rightarrow \infty\sqrt{-1}$ is exactly the genus zero case on the intersection of the circle $|\tau| = 1$ with the upper half-plane. Hence the Riemann surface is given by the equation $g(k, k) = 1/(4\Im^2(\tau))$ and the *first integral* is equal to $\mathcal{W} = \pi^2/\Im(\tau)$. In the other limit $\omega'/\omega \rightarrow 0\sqrt{-1}$ the Riemann surface tends to the singular curve of two copies of $\mathbb{P}^1$, which are connected by two ordinary double points. This is also the limit $y \rightarrow \infty$ of the unique *complex Fermi curves*, whose normalizations have two connected components, and whose *first integrals* are equal to 4π . In particular, in the limit $y \rightarrow \infty$ all the Riemann surfaces $\mathcal{F}_{\min}(\hat{\kappa}/2)$ tend to this unique singular curve, and the *first integral* converges to 4π . We conclude that with the choice $\hat{N} = 1, \check{N} = 1, N = 0$ the parameter space $(\omega'/\omega, z_1) \in \sqrt{-1}\mathbb{R}^+ \times (\omega/2 + \omega'[-1/2, 1/2])$ is mapped onto the domain $\{\tau \in \mathbb{C}^+ \mid \Re(\tau) \in [-1, 1] \text{ and } |\tau| > 1\}$. Not all of these *complex Fermi curves* are Riemann surfaces of the form $\mathcal{F}_{\min}(\hat{\kappa}_\tau/2)$. But all those elements, which are mapped into the fundamental domain $\mathcal{M}_1$, are of this form. These are all *complex Fermi curves* of the form $\mathcal{F}_{\min}(\kappa/2)$ of genus one.

Let us now calculate the absolute minimum of the *first integral* of this family. For any value of the parameter ω'/ω , the minimum of the *first integral* is attained for $z_1 = \omega/2$ and the value of the *first integral* at this point is equal to

$$\mathcal{W} = 8\pi(\eta + e_1\omega) \sqrt{\frac{\sqrt{2e_1^2 + e_2e_3}}{3e_1\sqrt{2e_1^2 + e_2e_3} + (4e_1^2 + e_2e_3)}}.$$

This expression does not depend on the choice of ω , it depends only on $\omega'/\omega \in \sqrt{-1}\mathbb{R}^+$. In the following lemma we will prove that this function has no critical point for $\omega'/\omega \in \mathbb{R}^+\sqrt{-1}$. As we have shown before, in the limit $\omega'/\omega \rightarrow \infty\sqrt{-1}$ this function converges to π^2 , and in the limit $\omega'/\omega \downarrow 0\sqrt{-1}$ to 4π . Since this function is analytic, this implies that it is strictly monotone decreasing and that the infimum is attained in the first limit, which is the genus zero case described above. This implies that the restriction of the *first integral* to the space of *Generalized Weierstraß curves* 4.4 is not smaller than $2\pi^2$, and that the minimum is attained by some unique *complex Fermi curve*, which corresponds to the *Clifford torus*, as we will see below.

Lemma 4.24. *The first integral of the family of complex Fermi curves $\mathcal{F}_{\min}(\hat{\kappa}_\tau/2)$ of genus one described above, has no critical point as a function depending on $\{\tau \in \mathbb{C}^+ \mid \Re(\tau) \in [-1, 1] \text{ and } \tau \neq 1\}$.*

Proof. We use an extension of the deformations constructed in Proposition 3.7. In fact, if we consider $\check{p}$ as a function depending on $\hat{p}$ and the moduli parameter, the form $\omega = \frac{\partial \check{p}}{\partial t} d\hat{p}$ corresponding to some change of the conformal class (x, y) has poles of third-order at $\infty^\pm$. More precisely, the singular part at $\infty^\pm$ is equal to

$$\left(\frac{\partial \Re(\tau)}{\partial t} \pm \sqrt{-1} \frac{\partial \Im(\tau)}{\partial t} \right) \hat{p} d\hat{p} \pm 8\pi^2 \sqrt{-1} \frac{\partial \mathcal{W}}{\partial t} \hat{p}^{-1} d\hat{p}.$$

We conclude that if we consider all meromorphic differentials on the normalization of the quotient modulo σ , which have poles of second-order at $\infty^\pm$, the analogous construction of Proposition 3.7 describes deformations of the conformal class. Consequently, a relative minimizer of this family cannot have a real (i. e. $\rho^* \omega = \eta^* \omega = \bar{\omega}$) meromorphic one-form on the normalization of the quotient modulo σ , which has at $\infty^\pm$ a second-order pole and a non-vanishing residue and no other poles, and which vanishes at the real point corresponding to the *Orbit of type 1*. Since all *complex Fermi curves* of this family are *real σ -hyperelliptic* the normalizations of the quotient modulo σ is isomorphic to $\mathbb{P}^1$ with one real cycle. Obviously there always exists a meromorphic one-form with the desired properties. Hence this family does not have a relative minimizer of the *first integral*. **q.e.d.**

Remark 4.25. *It is quite easy to describe the corresponding deformations and the analogous deformations of the family of genus two by ordinary differential equations in terms of the coefficients of some polynomials, whose zero set describes the normalization of the complex Fermi curves of this family, and the coefficients of some polynomials, which describe the corresponding meromorphic one-forms. This yields a numerical method to calculate the corresponding complex Fermi curves and their values of the first integrals (compare with [G-S-1, Section 2.2]).*

Remark 4.26. *The analogous calculation for real- σ -hyperelliptic Generalized Weierstraß curves 4.4 with two Orbits of type 1 shows, that for Willmore tori (i. e. critical points with respect to deformations changing the conformal class) the normalization of the quotient modulo σ is isomorphic to $\mathbb{P}^1$, with the two marked points $z = 0$ and $z = \infty$, the anti-holomorphic involution $z \mapsto 1/\bar{z}$, and the two real points $z = \pm 1$ corresponding to the two Orbits of type 1. The corresponding complex Fermi curves are exactly the Weierstraß curves 1.4 of minimal tori in S^3 described in [Hi].*

In conclusion, we want to describe the mappings $\Lambda \mapsto \Lambda_{[\kappa/2]}$ for the three non-zero elements $[\kappa/2]$ of $\Lambda^*/2\Lambda^*$ on $\mathcal{M}_1$ and the corresponding *Generalized Weierstraß curves* 4.4. In the following list we state explicitly the conformal equivalence class of $\Lambda_{[\kappa/2]}$ in dependence of the conformal equivalence class of Λ , considered as a mapping from domains in $\mathcal{M}_1$ into domains in $\mathcal{M}_1$. Moreover, we state the corresponding elements in the preimage of $[\kappa/2]$ under

the canonical map $\Lambda_{[\kappa/2]}^*/2\Lambda_{[\kappa/2]}^* \rightarrow \Lambda^*/2\Lambda^*$. Furthermore, we state the corresponding *complex Fermi curves* with lattice $\Lambda_{[\kappa/2]}$ introduced in Corollary 4.14, to which the *Generalized Weierstraß curve* 4.4. $\mathcal{F}_{\min, [\kappa/2]}(\mathcal{W}_{[\kappa/2]}/2)$ corresponds. Finally, we give the corresponding *geometric genus*. Due to our calculations above the *geometric genus* of $\mathcal{F}_{\min}(\check{\kappa}_\tau/2)$ is zero, the *geometric genus* of $\mathcal{F}_{\min}(\hat{\kappa}_\tau/2)$ is one if $|\tau| > 1$, and finally the *geometric genus* of $\mathcal{F}_{\min}(\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2)$ is two if $\Re(\tau) \neq -1/2$ and $\Re(\tau) \neq 1/2$.

The last two cases describe the neighbourhood of the *Clifford torus*. For all rectangular conformal classes (i. e. $\Re(\tau) = 0$) the minimizers corresponding to $[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]$ have two *Orbits of type 1*, all other minimizers have one *Orbit of type 1* and one *Orbit of type 2*. We shall see in the next section, that only this part of the family corresponds to *Weierstraß curves* 1.4. But due to the generalization of the Weierstraß representation to immersions into $\mathbb{R}^4$, the whole family corresponds to *quaternionic Weierstraß curves*, i. e. to immersions into $\mathbb{R}^4$ (compare with Lemma 2.27 and [P-P]).

We conjecture that for all conformal classes our lower bound for the Willmore functional on tori in $\mathbb{R}^3$ in dependence of the conformal class describes the absolute minimum of the restriction of the Willmore functional to tori in $\mathbb{R}^4$ of the given conformal class.

1. $[\hat{\kappa}_\tau/2]$:

(a) For $|\tau| \geq 2$:

- $\tau \mapsto \tau' = \tau/2$.
- The preimage of $[\hat{\kappa}_\tau/2]$ contains $[\hat{\kappa}_{\tau'}/2]$ and $[\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\hat{\kappa}_\tau/2]}(\mathcal{W}_{[\hat{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2)$.
- They have *geometric genus* two.

(b) For $|\tau| \leq 2$, $|\tau + 2| \geq 2$ and $|\tau - 2| \geq 2$:


- $\tau \mapsto \tau' = -2/\tau$.
- The preimage of $[\hat{\kappa}_\tau/2]$ contains $[\check{\kappa}_{\tau'}/2]$ and $[\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\hat{\kappa}_\tau/2]}(\mathcal{W}_{[\hat{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2)$.
- If $|\tau + 2| \neq 2$ and $|\tau - 2| \neq 2$, then they have *geometric genus* two, otherwise *geometric genus* one.

(c) For $|\tau + 2| \leq 2$:

- $\tau \mapsto \tau' = -2/\tau - 1$.
- The preimage of $[\hat{\kappa}_\tau/2]$ contains $[\hat{\kappa}_{\tau'}/2]$ and $[\check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\hat{\kappa}_\tau/2]}(\mathcal{W}_{[\hat{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2)$.
- They have *geometric genus* one.

(d) For $|\tau - 2| \leq 2$:

- $\tau \mapsto \tau' = -2/\tau + 1$.
- The preimage of $[\hat{\kappa}_\tau/2]$ contains $[\hat{\kappa}_{\tau'}/2]$ and $[\check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\hat{\kappa}_\tau/2]}(\mathcal{W}_{[\hat{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2)$.
- They have *geometric genus* one.



2. $[\check{\kappa}_\tau/2]$:

(a) For $-1/4 \leq \Re(\tau) \leq 1/4$:


- $\tau \mapsto \tau' = 2\tau$.
- The preimage of $[\check{\kappa}_\tau/2]$ contains $[\check{\kappa}_{\tau'}/2]$ and $[\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\check{\kappa}_\tau/2]}(\mathcal{W}_{[\check{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2)$.
- If $\Re(\tau) \neq \pm 1/4$, then they have *geometric genus* two, otherwise *geometric genus* one.

(b) For $\Re(\tau) \leq -1/4$:

- $\tau \mapsto \tau' = 2\tau + 1$.
- The preimage of $[\check{\kappa}_\tau/2]$ contains $[\hat{\kappa}_{\tau'}/2]$ and $[\check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\check{\kappa}_\tau/2]}(\mathcal{W}_{[\check{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2)$.
- They have *geometric genus* one.

(c) For $1/4 \leq \Re(\tau)$:

- $\tau \mapsto \tau' = 2\tau - 1$.
- The preimage of $[\check{\kappa}_\tau/2]$ contains $[\hat{\kappa}_{\tau'}/2]$ and $[\check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\check{\kappa}_\tau/2]}(\mathcal{W}_{[\check{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2)$.
- They have *geometric genus* one.



3. $[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]$:

(a) For $2 \leq |\tau + 1|$, $2 \leq |\tau - 1|$ and $\Re(\tau) \leq 0$:

- $\tau \mapsto \tau' = (\tau + 1)/2$.
- The preimage of $[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]$ contains $[\check{\kappa}_{\tau'}/2]$ and $[\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}(\mathcal{W}_{[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2)$.
- If $\Re(\tau) \neq 0$, then they have *geometric genus* two, otherwise *geometric genus* one.

(b) For $2 \leq |\tau + 1|$, $2 \leq |\tau - 1|$ and $0 \leq \Re(\tau)$:

- $\tau \mapsto \tau' = (\tau - 1)/2$.

- The preimage of $[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]$ contains $[\check{\kappa}_{\tau'}/2]$ and $[\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}(\mathcal{W}_{[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2)$.
- If $\Re(\tau) \neq 0$, then they have *geometric genus* two, otherwise *geometric genus* one.

(c) For $|\tau + 1| \leq 2$ and $2 \leq |\tau - 1|$:

- $\tau \mapsto \tau' = -2/(\tau + 1)$.
- The preimage of $[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]$ contains $[\hat{\kappa}_{\tau'}/2]$ and $[\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}(\mathcal{W}_{[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2)$.
- If $|\tau - 1| \neq 2$, then they have *geometric genus* two, otherwise *geometric genus* one.

(d) For $2 \leq |\tau + 1|$ and $|\tau - 1| \leq 2$:


- $\tau \mapsto \tau' = -2/(\tau - 1)$.
- The preimage of $[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]$ contains $[\hat{\kappa}_{\tau'}/2]$ and $[\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}(\mathcal{W}_{[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2 + \check{\kappa}_{\tau'}/2)$.
- If $|\tau + 1| \neq 2$, then they have *geometric genus* two, otherwise *geometric genus* one.

(e) For $|\tau + 1| \leq 2$, $|\tau - 1| \leq 2$ and $\Re(\tau) \leq 0$:

- $\tau \mapsto \tau' = (\tau - 1)/(\tau + 1)$.
- The preimage of $[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]$ contains $[\hat{\kappa}_{\tau'}/2]$ and $[\check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}(\mathcal{W}_{[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2)$.
- If $\Re(\tau) \neq 0$, then they have *geometric genus* one, otherwise *geometric genus* zero.

(f) For $|\tau + 1| \leq 2$, $|\tau - 1| \leq 2$ and $0 \leq \Re(\tau)$:

- $\tau \mapsto \tau' = -(\tau + 1)/(\tau - 1)$.
- The preimage of $[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]$ contains $[\hat{\kappa}_{\tau'}/2]$ and $[\check{\kappa}_{\tau'}/2]$.
- $\mathcal{F}_{\min, [\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}(\mathcal{W}_{[\hat{\kappa}_\tau/2 + \check{\kappa}_\tau/2]}/2)$ corresponds to $\mathcal{F}_{\min}(\hat{\kappa}_{\tau'}/2)$.



- If $\Re(\tau) \neq 0$, then they have *geometric genus* one, otherwise *geometric genus* zero.

5 Weierstraß potentials

5.1 The *Singularity condition*

In Lemma 4.1 we proved that the *complex Fermi curve* of some potential, which corresponds to some immersion has to fulfill the *Weak Singularity condition* 4.2. But this was only a necessary condition on the *complex Fermi curve* to be a *Weierstraß curve* 1.4. In this section we want to investigate two problems.

First problem: Find a necessary and sufficient condition on the *complex Fermi curve* to be a *Weierstraß curve* 1.4.

Second problem: Given a *Weierstraß curve* 1.4, find a necessary and sufficient condition on a potential, which corresponds to the given *complex Fermi curve*, to be a *Weierstraß potential* 1.3. Section 2.4 suggests a decomposition of all potentials corresponding to a given *complex Fermi curve* into potentials, which corresponds to some one-sheeted coverings of *complex Fermi curve*. In particular, we are interested in the question, which of these subclasses of potentials contains no *Weierstraß potentials* 1.3, which of these contains some *Weierstraß potential* 1.3, and which of these subclasses are contained in the set of *Weierstraß potentials* 1.3.

If U is a potential, whose *complex Fermi curve* fulfills the *Weak Singularity condition* 4.2, then at least one of the following cases takes place:

1. The preimage of $[\kappa/2]$ contains at least two *cusps*. (Due to Corollary 2.17 (iii) the number of such elements has to be even.)
2. The preimage of $[\kappa/2]$ contains two different elements y and y' , such that the values of the eigenfunction at these two points are linearly dependent. (Due to Corollary 2.17 (ii), in this case $\eta(y)$ and $\eta(y')$ are two other points with this property.)
3. The preimage of $[\kappa/2]$ contains only *multiple points* and the values of the eigenfunction at all elements of the preimage are pairwise linearly independent. We may assume that $d\hat{p}$ has no zero in the preimage of $[\kappa/2]$.
 - (a) The preimage contains an *Orbit of type 3*.
 - (b) The preimage contains altogether at least three *Orbits of type 1* or *type 2*.
 - (c) The preimage contains exactly two *Orbits of type 1* and no other orbit.
 - (d) The preimage contains exactly two *Orbits of type 2* and no other orbit.
 - (e) The preimage contains exactly one *Orbit of type 1*, one *Orbit of type 2*, and no *Orbit of type 3*.

In cases 1 and case 2 the preimage contains a singularity of the *Structure sheaf* 2.24 in contrast to case 3. Both former cases can take place simultaneously. Also case 3a and case 3b can take place simultaneously. In case 1 y is a branch point with respect to the coverings over $p \in \mathbb{C}$ and $p' \in \mathbb{C}$ described in the proof of Lemma 4.1. Hence due to Lemma 2.22 the function $\psi(y)$ obeys the *Equivalent form of the Periodicity condition* 1.2. Analogously Lemma 2.22 shows that in case 2 the value of the eigenfunction at y , and in case 3a the values of the eigenfunction at all four points of the *Orbit of type 3* fulfill the *Equivalent form of the Periodicity condition* 1.2.

Lemma 5.1. (i) *In case 3b there always exists some eigenfunction, which fulfills the Equivalent form of the Periodicity condition 1.2.*

(ii) *In case 3c–3e there exists some eigenfunction, which fulfills the Equivalent form of the Periodicity condition 1.2, if and only if the functions $d\check{p}/d\hat{p}$ take the same value at two different elements of the preimage of $[\kappa/2]$, which belong to different orbits.*

Proof. In case 3 there always exists two generators $\hat{\gamma}$ and $\check{\gamma}$ of the period lattice Λ , whose differentials $d\hat{p} = dg(\hat{\gamma}, k)$ and $d\check{p} = dg(\check{\gamma}, k)$ have no zeroes in the preimage of $[\kappa/2]$. Thus, due to Lemma 2.23, the ranges of the projections $\check{P}_{\hat{\gamma}}([\kappa/2])$ and $\check{P}_{\check{\gamma}}([\kappa/2])$ are spanned by the values of the eigenfunction at the elements of the preimage of $[\kappa/2]$. If y_1 and $y_2 = \eta(y_1)$ are two elements of an *Orbit of type 1* or *type 2*, then let ψ_1 and ψ_2 denote the corresponding values of the eigenfunction. Due to Corollary 2.17 we may assume that ψ_2 is equal to $-J\bar{\psi}_1$, and therefore ψ_1 is equal to $J\bar{\psi}_2$. The bilinear forms $\langle\langle J\psi_i, \psi_j \rangle\rangle_{\hat{\gamma}}$ and $\langle\langle J\psi_i, \psi_j \rangle\rangle_{\check{\gamma}}$ form two 2×2 matrices. These bilinear forms are symmetric and transforms under the anti-unitary map $\chi \mapsto J\bar{\chi}$ like

$$\langle\langle J\bar{\chi}, J\bar{\xi} \rangle\rangle_{\hat{\gamma}} = -\overline{\langle\langle \chi, \xi \rangle\rangle_{\check{\gamma}}} \text{ and } \langle\langle J\bar{\chi}, J\bar{\xi} \rangle\rangle_{\check{\gamma}} = -\overline{\langle\langle \chi, \xi \rangle\rangle_{\hat{\gamma}}}.$$

Hence these matrices have the form $\begin{pmatrix} \alpha & \beta \\ \beta & -\bar{\alpha} \end{pmatrix}$ and $\begin{pmatrix} \alpha' & \beta' \\ \beta' & -\bar{\alpha}' \end{pmatrix}$, with complex numbers α, β, α' and β' , respectively. Moreover, the matrices, whose entries are given as $\langle\langle \bar{\psi}_i, \psi_j \rangle\rangle_{\hat{\gamma}}$ and $\langle\langle \bar{\psi}_i, \psi_j \rangle\rangle_{\check{\gamma}}$, are of the form $\begin{pmatrix} -\beta & \bar{\alpha} \\ \alpha & \beta \end{pmatrix}$ and $\begin{pmatrix} -\beta' & \bar{\alpha}' \\ \alpha' & \beta' \end{pmatrix}$, respectively. Since these bilinear forms are hermitian, β and β' have to be real. Due to Lemma 2.16 and Lemma 2.22 the numbers α and α' have to be zero if we consider an *Orbit of type 1*, and the numbers β and β' have to be zero if we consider an *Orbit of type 2*. Due to Lemma 2.23 the other numbers are not equal to zero. Due to the proof of property (iii) of Lemma 2.23 the value of the function $d\check{p}/d\hat{p}$ at the point y_1 has to be equal to β'/β if we consider an *Orbit of type 1*, and equal to α'/α , if we consider an *Orbit of type 2*. If in the first case we change the basis ψ_1, ψ_2 into $\frac{\psi_1 + \psi_2}{\sqrt{2}}, \frac{\psi_2 - \psi_1}{\sqrt{2}}$, then the matrices corresponding to the bilinear forms $\langle\langle J\cdot, \cdot \rangle\rangle_{\hat{\gamma}}$, $\langle\langle J\cdot, \cdot \rangle\rangle_{\check{\gamma}}$, $\langle\langle \bar{\cdot}, \cdot \rangle\rangle_{\hat{\gamma}}$ and $\langle\langle \bar{\cdot}, \cdot \rangle\rangle_{\check{\gamma}}$ transforms to $\begin{pmatrix} \beta & 0 \\ 0 & -\beta \end{pmatrix}$, $\begin{pmatrix} \beta' & 0 \\ 0 & -\beta' \end{pmatrix}$, $\begin{pmatrix} 0 & \beta \\ \beta & 0 \end{pmatrix}$ and $\begin{pmatrix} 0 & \beta' \\ \beta' & 0 \end{pmatrix}$. As well for *Orbits of type 1* as for *Orbits of type 2* after some renormalization these matrices may be brought into the form

$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} \frac{d\check{p}}{d\hat{p}}(y_1) & 0 \\ 0 & -\frac{d\check{p}}{d\hat{p}}(y_1) \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \text{ and } \begin{pmatrix} 0 & \frac{d\check{p}}{d\hat{p}}(y_1) \\ \frac{d\check{p}}{d\hat{p}}(y_1) & 0 \end{pmatrix}.$$

Moreover, $\overline{\frac{d\check{p}}{d\hat{p}}}(y_1)$ is equal to $\frac{d\check{p}}{d\hat{p}}(y_2)$, and both numbers are real and therefore equal if we consider an *Orbit of type 1*. Now let us prove statement (ii). For arbitrary non-zero complex numbers α and β , which are the values of $d\check{p}/d\hat{p}$ at two different elements belonging to two different orbits, we have to determine the solutions of the following four equations with complex numbers z_1, z_2, z_3 and z_4 :

$$\begin{aligned} (z_1\bar{z}_2 + \bar{z}_1z_2) + (z_3\bar{z}_4 + \bar{z}_3z_4) &= 0 & (z_1^2 - z_2^2) + (z_3^2 - z_4^2) &= 0 \\ (\alpha z_1\bar{z}_2 + \bar{\alpha}\bar{z}_1z_2) + (\beta z_3\bar{z}_4 + \bar{\beta}\bar{z}_3z_4) &= 0 & (\alpha z_1^2 - \bar{\alpha}z_2^2) + (\beta z_3^2 - \bar{\beta}z_4^2) &= 0 \end{aligned}$$

These equations imply

$$\begin{aligned} (\alpha - \bar{\alpha})^2(z_1\bar{z}_2)^2 &= ((\bar{\alpha} - \beta)z_3\bar{z}_4 + (\bar{\alpha} - \bar{\beta})\bar{z}_3z_4)^2 \\ &= -((\bar{\alpha} - \beta)z_3^2 - (\bar{\alpha} - \bar{\beta})z_4^2)((\bar{\alpha} - \bar{\beta})\bar{z}_3^2 - (\bar{\alpha} - \beta)\bar{z}_4^2). \end{aligned}$$

The difference of these two equations yields

$$(\bar{\alpha} - \beta)(\bar{\alpha} - \bar{\beta})(z_3\bar{z}_3 + z_4\bar{z}_4)^2 = 0.$$

Thus either α is equal to β or α is equal to $\bar{\beta}$ or the only solution is the trivial solution (i. e. the complex numbers z_1, z_2, z_3 and z_4 are all zero). If α is equal to β , then $z_3 = \sqrt{-1}z_1$ and $z_4 = -\sqrt{-1}z_2$ is a solution for arbitrary z_1 and z_2 . If α is equal to $\bar{\beta}$, then $z_3 = z_2$ and $z_4 = -z_1$ is a solution for arbitrary z_1 and z_2 . This proves (ii). The proof of (i) is similar. **q.e.d.**

Whether case 1 takes place or not depends only on the *complex Fermi curve* and not on the choice of the potential, in contrast, whether case 2 takes place depends on the choice of the potential. In the rest of this section we want to find some property of the *complex Fermi curve*, which is equivalent to the existence of some potential, which leads to the given *complex Fermi curve* and which is the potential of some immersion. To all subcases of case 3 there might exist an analogous subcase of case 2. Thus the main question is, whether to a potential of cases 3c–3e there exists another potential, which leads to the same *complex Fermi curve*, but corresponds to the analogous subcase of case 2. Fortunately there exists some method to transform subcases of case 3 into subcases of case 2, the so-called *Bäcklund transformation* ([E-K],[M-S, Chapter 6.] and [L-McL, Section 4]).

Lemma 5.2. (i) *To all potentials of cases 3c there exists another potential, which leads to the same complex Fermi curve and corresponds to the analogous subcase of case 2.*

(ii) *If the preimage of $[\kappa/2]$ contains exactly four multiple points, at which the function $d\check{p}/d\hat{p}$ takes more than two values, then there exists no immersion, which leads to this complex Fermi curve.*

Proof. If in case (i) the values of the eigenfunction ψ at the four elements of the two *Orbits of type 1* are linearly independent, then one of the Bäcklund transformations described in Lemma 2.29 has the desired properties.

The condition in case (ii) excludes *Multiple points O1/O1* and *Multiple points O3*. It remains to consider the cases *Multiple points O1/O2* and *Multiple points O2/O2*. Due to Lemma 5.1 we have to show that in these cases there do not exist real potentials with the given *complex Fermi curve*, whose eigenfunctions ψ take linearly dependent values at these four *multiple points*. In case of *Multiple points O1/O2* this follows from Corollary 2.17. If in case *Multiple points O2/O2* the values of the eigenfunction ψ at the four points are linearly dependent, then, due to Corollary 2.17 and Lemma 2.23, for any non-vanishing normalization of the eigenfunction ψ the functions $\langle \langle J\sigma^*\psi, \psi \rangle \rangle_{\hat{\gamma}}$ has a zero at the two *Orbits of type 2*. Since this function is invariant under σ , it has to have zeroes of even order at the fixed points of σ . Consequently, the corresponding *Local contribution to the arithmetic genus* 4.3 of the *Structure sheaf* 2.24 has to be larger than one, and the function $d\check{p}/d\hat{p}$ has to take the same value at two points of different *Orbits of type 2*. **q.e.d.**

Thus the desired property of the *complex Fermi curve* is the

Singularity condition 5.3. *There exists some $\kappa \in \Lambda^*$, such that $[\kappa/2]$ belongs to the complex Fermi curve $\mathcal{F}(U, U)/\Lambda^*$. Moreover, the preimage of this point in the normalization of $\mathcal{F}(U, U)/\Lambda^*$ contains either two zeroes of both differentials $d\hat{p}$ and $d\check{p}$, or at least six elements, or exactly four elements. Furthermore, in the last case the function $d\check{p}/d\hat{p}$ takes the same value at two different elements, which are not interchanged by η .*

The discussion above is summarized in

Theorem 5.4. *For a real potential $U \in L^2(\mathbb{R}^2/\Lambda)$ we have the following characterizations:*

- (i) *The complex Fermi curve $\mathcal{F}(U, U)$ is a Weierstraß curve 1.4 if and only if the Singularity condition 5.3 is fulfilled.*
- (ii) *Let $\mathcal{F}(U, U)$ fulfill the Singularity condition 5.3. Then the potential U is a Weierstraß potential 1.3 if one of the cases 1,2,3a or 3b takes place. Furthermore, in the remaining cases 3c–3e the same is true if and only if the functions $d\check{p}/d\hat{p}$ take the same value at two different elements of the preimage of $[\kappa/2]$, which belong to different orbits. **q.e.d.***

5.2 Variations of immersions

Due to the Weierstraß representation [Ta-1, Fr-2] the conformal class of some immersion is given by the conformal class of the flat torus $\mathbb{R}^2/\Lambda$ with the Euclidean metric $g(\cdot, \cdot)$. Thus for all variations of some immersion with potential U , which do not change the conformal class, there exists some variation δU of the potential U , which is a sufficiently smooth function on $\mathbb{R}^2/\Lambda$. Let us now characterize some variations δU of the potential U , which induce a variation of the immersion and do not change the conformal class.

Proposition 5.5. *Let U be the potential of some immersion and $\delta U \in C(\mathbb{R}^2/\Lambda)$ some real variation of this potential. If for all $\hat{\mu} \in \mathbb{C}^2$ and all regular forms ω on the associated complex Fermi curve the pullback of the function $\Omega_{U,U}(\delta U, \delta U)/dg(\hat{\mu}, k) \cdot \omega/dg(\hat{\mu}, k)$ to the normalization of $\mathcal{F}(U, U)/\Lambda^*$ is holomorphic on some neighbourhood of the preimage of $[\kappa/2]$*

and equal to zero at all elements of this preimage, then there exists some variation of the immersion, which corresponds to the variation δU and does not change the conformal class.

In order to prove this proposition we need some preparation.

Lemma 5.6. *Let $\hat{\mu} \in \mathbb{C}^2$ satisfy the condition that $dg(\hat{\mu}, k)$ does not vanish identically on any connected component of the normalization of $\mathcal{F}(V, W)$. Also let χ be an eigenfunction corresponding to the element k' of $\mathcal{F}(V, W)$. Assume that for all regular forms ω of the complex Fermi curve the pullback of the function $\Omega_{V,W}(\delta V, \delta W)/dg(\hat{\mu}, k) \cdot \omega/dg(\hat{\mu}, k)$ to the normalization of $\mathcal{F}(V, W)$ is holomorphic on some neighbourhood of the preimage of k' and equal to zero at all elements of the preimage of k' . Then there exists a variation $\delta_{\hat{\mu}}\chi$ of χ associated with the variations δV and δW (this means*

$$\begin{pmatrix} \delta V & 0 \\ 0 & \delta W \end{pmatrix} \begin{pmatrix} \chi_1 \\ \chi_2 \end{pmatrix} + \begin{pmatrix} V & \partial_{[k']} \\ \bar{\partial}_{[k']} & W \end{pmatrix} \begin{pmatrix} \delta_{\hat{\mu}}\chi_1 \\ \delta_{\hat{\mu}}\chi_2 \end{pmatrix} = 0 \text{ and } \delta_{\hat{\mu}}\chi(x + \hat{\gamma}) = \exp(2\pi\sqrt{-1}g(\hat{\gamma}, k')) \delta_{\hat{\mu}}\chi(x)$$

for all $x \in \mathbb{R}^2$ and all periods $\hat{\gamma} \in \Lambda$), such that $\langle \langle \phi, \delta_{\hat{\mu}}\chi \rangle \rangle_{\hat{\mu}}$ vanishes for all eigenfunctions ϕ of the transposed Dirac operator, which correspond to the element k' .

Proof. Arguments similar to those in the proof of condition (iii) of Lemma 2.23 show that for all $\hat{\mu}, \check{\mu} \in \mathbb{C}^2$ on the complex Fermi curve the identity

$$\langle \langle \phi(k), \psi(k) \rangle \rangle_{\hat{\mu}} dg(\hat{\mu}, k) = \langle \langle \phi(k), \psi(k) \rangle \rangle_{\check{\mu}} dg(\check{\mu}, k)$$

holds. Thus the first assumption of the lemma implies that the denominator of the projection $\tilde{P}_{\hat{\mu}}$ does not vanish identically on the complex Fermi curve, and therefore this projection may be restricted to the complex Fermi curve. Moreover, we choose some $\check{\mu} \in \mathbb{C}^2$, which is linearly independent of $\hat{\mu}$, and parameterize the complex Fermi curve by $\hat{p} = g(\hat{\mu}, k)$ and $\check{p} = g(\check{\mu}, k)$. Now let us consider the normalizations of the complex Fermi curves $\mathcal{F}(V + t\delta V, W + t\delta W)$ locally as covering spaces over $(\hat{p}, t) \in \mathbb{C}^2$, and let us use the convention $\partial\hat{p}/\partial t = 0$. We conclude that the local sum of the function $\tilde{P}_{\hat{\mu}}$ over all sheets, which contain the element $(k', t = 0)$ (compare with Remark 2.19), is locally a holomorphic function depending on $\hat{p}$ and t . Hence the partial derivative of this function with respect to t at the point $\hat{p} = \hat{p}' = g(\hat{\mu}, k'), t = 0$ is some finite rank operator denoted by $\delta\tilde{P}_{\hat{\mu}}(k')$. Now we claim that $\delta_{\hat{\mu}}\chi = \psi_{k'}\delta\tilde{P}_{\hat{\mu}}(k')\psi_{-k'}\chi$ (or equivalently $\delta_{\hat{\mu}}\tilde{\chi} = \delta\tilde{P}_{\hat{\mu}}(k')\tilde{\chi}$) has the required properties. Since the local sum of the function $\tilde{P}_{\hat{\mu}}$ is a projection valued function, we have the identity $\delta\tilde{P}_{\hat{\mu}}(k')\tilde{P}_{\hat{\mu}}(k') + \tilde{P}_{\hat{\mu}}(k')\delta\tilde{P}_{\hat{\mu}}(k') = \delta\tilde{P}_{\hat{\mu}}(k')$, and therefore also $\tilde{P}_{\hat{\mu}}(k')\delta\tilde{P}_{\hat{\mu}}(k')\tilde{P}_{\hat{\mu}}(k') = 0$. This implies the last property. All projections $\tilde{P}_{\hat{\mu}}$ project onto double periodic functions, which implies

$$\delta_{\hat{\mu}}\chi(x + \hat{\gamma}) = \exp(2\pi\sqrt{-1}g(\hat{\gamma}, k')) \delta_{\hat{\mu}}\chi(x) \quad \text{for all } x \in \mathbb{R}^2 \text{ and all } \hat{\gamma} \in \Lambda.$$

Due to our above discussion of Lemma 3.1, the function $\Omega_{V,W}(\delta V, \delta W)/d\hat{p}$ is then proportional to $\partial\check{p}/\partial t$. Therefore, the assumption on the form $\Omega_{V,W}(\delta V, \delta W)$ implies that the pullback of $\partial\check{p}/\partial t \cdot \tilde{P}_{\hat{\mu}}$ to the normalization of $\mathcal{F}(V, W)$ is holomorphic on some neighbourhood of the

preimage of k' and equal to zero at all elements of the preimage of k' . This implies the following operator relation:

$$\left(\sum_{i=1}^l \left(\tilde{D}_{\hat{\mu}, \tilde{\mu}}(\hat{p}') - \check{p}'\pi \mathbf{1} \right)^{i-1} \delta \tilde{D}_{\hat{\mu}, \tilde{\mu}} \left(\tilde{D}_{\hat{\mu}, \tilde{\mu}}(\hat{p}') - \check{p}'\pi \mathbf{1} \right)^{l-i} \right) \check{P}_{\hat{\mu}}(k') + \left(\tilde{D}_{\hat{\mu}, \tilde{\mu}}(\hat{p}) - \check{p}'\pi \mathbf{1} \right)^l \delta \check{P}_{\hat{\mu}}(k') = 0.$$

Here l is the dimension of the range of $\check{P}_{\hat{\mu}}(k')$, $\delta \tilde{D}_{\hat{\mu}, \tilde{\mu}} = \begin{pmatrix} 0 & \frac{\delta W}{\tilde{\nu}_2 - \sqrt{-1}\tilde{\nu}_1} \\ \frac{\delta V}{\tilde{\nu}_2 + \sqrt{-1}\tilde{\nu}_1} & 0 \end{pmatrix}$ denotes the variation of the operator $\tilde{D}_{\hat{\mu}, \tilde{\mu}}(\hat{p})$, and $\hat{p}' = g(\hat{\mu}, k')$ and $\check{p}' = g(\tilde{\mu}, k')$ are the components of k' . Since $\psi_{-k'}\chi$ is a proper eigenfunction of $\tilde{D}_{\hat{\mu}, \tilde{\mu}}(\hat{p})$ with the eigenvalues $\check{p}'\pi$, the variation $\psi_{-k'}\delta_{\hat{\mu}}\chi$ satisfies

$$\left(\tilde{D}_{\hat{\mu}, \tilde{\mu}}(\hat{p}) - \check{p}'\pi \mathbf{1} \right)^{l-1} \delta \tilde{D}_{\hat{\mu}, \tilde{\mu}} \psi_{-k'}\chi + \left(\tilde{D}_{\hat{\mu}, \tilde{\mu}}(\hat{p}) - \check{p}'\pi \mathbf{1} \right)^l \psi_{-k'}\delta_{\hat{\mu}}\chi = 0.$$

Since $\check{P}_{\hat{\mu}}(k')$ is the spectral projection onto the generalized eigenspace of $\tilde{D}_{\hat{\mu}, \tilde{\mu}}(\hat{p})$ associated with eigenvalues $\check{p}'\pi$, the restriction of the operator $\tilde{D}_{\hat{\mu}, \tilde{\mu}}(\hat{p}) - \check{p}'\pi \mathbf{1}$ to the kernel of the operator $\check{P}_{\hat{\mu}}(k')$ is invertible. Thus we have $\delta \tilde{D}_{\hat{\mu}, \tilde{\mu}} \psi_{-k'}\chi + \left(\tilde{D}_{\hat{\mu}, \tilde{\mu}}(\hat{p}) - \check{p}'\pi \mathbf{1} \right) \psi_{-k'}\delta_{\hat{\mu}}\chi = 0$, which is equivalent to $\begin{pmatrix} \delta V & 0 \\ 0 & \delta W \end{pmatrix} \begin{pmatrix} \chi_1 \\ \chi_2 \end{pmatrix} + \begin{pmatrix} V & \partial_{[k']} \\ -\bar{\partial}_{[k']} & W \end{pmatrix} \begin{pmatrix} \delta_{\hat{\mu}}\chi_1 \\ \delta_{\hat{\mu}}\chi_2 \end{pmatrix} = 0.$ **q.e.d.**

Proof of Proposition 5.5. Let $E([\kappa/2])$ be the eigenspace of the Dirac operator $D_{[\kappa/2]}$ with real potential U corresponding to the eigenvalue $\lambda = 0$. Due to Lemma 2.16 this operator is self-adjoint, and $E([\kappa/2])$ is equal to the range of $\check{P}([\kappa/2])$. Moreover, the anti-unitary operator $\chi \mapsto J\bar{\chi}$ leaves this subspace invariant. Due to the *Equivalent form of the Periodicity condition* 1.2 all non-zero elements χ of $E([\kappa/2])$, which satisfy the following two conditions, induce an immersion:

(A) for all $\hat{\mu} \in \mathbb{C}^2$ the bilinear form $\langle \langle J\chi, \chi \rangle \rangle_{\hat{\mu}}$ is zero.

(B) for all $\hat{\mu} \in \mathbb{C}^2$ the bilinear form $\langle \langle \bar{\chi}, \chi \rangle \rangle_{\hat{\mu}}$ is zero.

If δU is some variation of the potential U , which satisfies the assumption of Proposition 5.5, then due to Lemma 5.6 there exists some variation $\delta\chi$ of χ associated with the variation δU of the potential U :

$$\begin{pmatrix} \delta U & 0 \\ 0 & \delta U \end{pmatrix} \begin{pmatrix} \chi_1 \\ \chi_2 \end{pmatrix} + \begin{pmatrix} U & \partial_{[\kappa/2]} \\ -\bar{\partial}_{[\kappa/2]} & U \end{pmatrix} \begin{pmatrix} \delta\chi_1 \\ \delta\chi_2 \end{pmatrix} = 0 \quad \text{and} \quad \delta\chi(x + \hat{\gamma}) = (-1)^{g(\hat{\gamma}, \kappa)} \chi(x)$$

for all $x \in \mathbb{R}^2$ and all periods $\hat{\gamma} \in \Lambda$. Obviously we may add to such a variations $\delta\chi$ some arbitrary element of $E([\kappa/2])$ and again obtain such a variation. In a second step we show that there exists some $\delta\chi$ among these variations, which satisfies

(A') for all $\hat{\mu} \in \mathbb{C}^2$ the bilinear form $\langle\langle J\chi, \delta\chi \rangle\rangle_{\hat{\mu}}$ is zero.

(B') for all $\hat{\mu} \in \mathbb{C}^2$ the bilinear form $\langle\langle \bar{\chi}, \delta\chi \rangle\rangle_{\hat{\mu}}$ is zero.

Again, due to Lemma 5.6, for all $\hat{\mu} \in \mathbb{C}^2$ there exists such a variation $\delta_{\hat{\mu}}\chi$, such that $\langle\langle J\xi, \delta_{\hat{\mu}}\chi \rangle\rangle_{\hat{\mu}}$ vanishes for all $\xi \in E([\kappa/2])$. In particular, $\langle\langle J\chi, \delta_{\hat{\mu}}\chi \rangle\rangle_{\hat{\mu}}$ and $\langle\langle \bar{\chi}, \delta_{\hat{\mu}}\chi \rangle\rangle_{\hat{\mu}}$ vanish. For linearly independent $\hat{\mu}$ and $\tilde{\mu}$ the difference $\delta_{\hat{\mu}}\chi - \delta_{\tilde{\mu}}\chi$ can be non-zero, but has to be an element of $E([\kappa/2])$. Now we again choose two generators $\hat{\gamma}$ and $\tilde{\gamma}$ of the period lattice. Let $\langle\cdot, \cdot\rangle_1$ denote the restriction of $\langle\langle J\cdot, \cdot \rangle\rangle_{\hat{\gamma}}$ and $\langle\cdot, \cdot\rangle_2$ the restriction of $\langle\langle J\cdot, \cdot \rangle\rangle_{\tilde{\gamma}}$ to $E([\kappa/2]) \times E([\kappa/2])$. Both bilinear forms are symmetric and satisfy

$$\langle J\bar{\chi}, J\bar{\xi} \rangle_1 = -\overline{\langle \chi, \xi \rangle_1} \quad \text{and} \quad \langle J\bar{\chi}, J\bar{\xi} \rangle_2 = -\overline{\langle \chi, \xi \rangle_2}.$$

Lemma 5.7. *One of the following two cases hold:*

- (a) *The four linear forms $\langle \chi, \cdot \rangle_1$, $\langle \chi, \cdot \rangle_2$, $\langle J\bar{\chi}, \cdot \rangle_1$ and $\langle J\bar{\chi}, \cdot \rangle_2$ on $E([\kappa/2])$ are linearly independent.*
- (b) *There exists a non-zero linear combination χ' of χ and $J\bar{\chi}$, such that the linear forms $\langle \chi', \cdot \rangle_1$ and $\langle \chi', \cdot \rangle_2$ on $E([\kappa/2])$ are linearly dependent.*

Proof. If the form $\langle \chi, \cdot \rangle_1$ is zero, then case (b) takes place with $\chi' = \chi$. Now let us assume that this linear form does not vanish. The subspace of $E([\kappa/2])$, on which the two linear forms $\langle \chi, \cdot \rangle_1$ and $\langle J\bar{\chi}, \cdot \rangle_1$ vanish, contains the subspace, on which all four linear forms vanish. Furthermore, both subspaces are invariant under the anti-unitary operator $\xi \mapsto J\bar{\xi}$. The same arguments as in the proof of Corollary 2.17 (iii) show that either these two subspaces coincide, or the co-dimension of the second subspace in $E([\kappa/2])$ is equal to four, which implies that case (a) takes place. If the two subspaces of $E([\kappa/2])$ coincide, then there exist two complex numbers α and β , such that the linear forms fulfill the following equations:

$$\langle \chi, \cdot \rangle_2 = \alpha \langle \chi, \cdot \rangle_1 + \beta \langle J\bar{\chi}, \cdot \rangle_1 \quad \text{and} \quad \langle J\bar{\chi}, \cdot \rangle_2 = -\bar{\beta} \langle \chi, \cdot \rangle_1 + \bar{\alpha} \langle J\bar{\chi}, \cdot \rangle_1.$$

If $\begin{pmatrix} \gamma \\ \delta \end{pmatrix}$ is any non-zero eigenvector of the matrix $\begin{pmatrix} \alpha & -\bar{\beta} \\ \beta & \bar{\alpha} \end{pmatrix}$, then $\chi' = \gamma\chi + \delta J\bar{\chi}$ fulfills the condition of case (b). **q.e.d.**

Continuation of the proof of Proposition 5.5. In case (a) we may add to some $\delta_{\hat{\mu}}\chi$ some element of $E([\kappa/2])$, such that the sum fulfills conditions (A') and (B'). In case (b) we may assume that the linear forms $\alpha \langle \chi', \cdot \rangle_1 + \langle \chi', \cdot \rangle_2$ and $\bar{\alpha} \langle J\bar{\chi}', \cdot \rangle_1 + \langle J\bar{\chi}', \cdot \rangle_2$ vanish. Indeed, if the form $\langle \chi', \cdot \rangle_1$ vanishes, then we replace $\hat{\gamma}$ and $\tilde{\gamma}$ (i. e. $\langle \chi', \cdot \rangle_1$ and $\langle \chi', \cdot \rangle_2$). Due to Lemma 5.6 the variations $\delta_{\hat{\gamma}}\chi'$ obey the equations

$$\langle \chi', \delta_{\hat{\gamma}}\chi' \rangle_1 = 0 = \langle J\bar{\chi}', \delta_{\hat{\gamma}}\chi' \rangle_1.$$

If $\hat{\mu}$ is equal to $\alpha\hat{\gamma} + \tilde{\gamma}$ and $\tilde{\mu}$ equal to $\bar{\alpha}\hat{\gamma} + \tilde{\gamma}$, then the variations $\delta_{\hat{\mu}}\chi'$ and $\delta_{\tilde{\mu}}\chi'$ satisfy

$$\langle\langle J\chi', \delta_{\hat{\mu}}\chi' \rangle\rangle_{\hat{\mu}} = 0 = \langle\langle \bar{\chi}', \delta_{\tilde{\mu}}\chi' \rangle\rangle_{\tilde{\mu}}.$$

Since the differences $\delta_{\dot{\gamma}}\chi' - \delta_{\bar{\mu}}\chi'$ and $\delta_{\dot{\gamma}}\chi' - \delta_{\bar{\mu}}\chi'$ belong to $E([\kappa/2])$, the variation $\delta_{\dot{\gamma}}\chi'$ satisfies

$$\begin{aligned} \langle \langle J\chi', \delta_{\dot{\gamma}}\chi' \rangle \rangle_{\bar{\mu}} &= \langle \langle J\chi', \delta_{\dot{\gamma}}\chi' - \delta_{\bar{\mu}}\chi' \rangle \rangle_{\bar{\mu}} = \alpha \langle \chi', \delta_{\dot{\gamma}}\chi' - \delta_{\bar{\mu}}\chi' \rangle_1 + \langle \chi', \delta_{\dot{\gamma}}\chi' - \delta_{\bar{\mu}}\chi' \rangle_2 = 0 \\ -\langle \langle \bar{J}\chi', \delta_{\dot{\gamma}}\chi' \rangle \rangle_{\bar{\mu}} &= -\langle \langle \bar{J}\chi', \delta_{\dot{\gamma}}\chi' - \delta_{\bar{\mu}}\chi' \rangle \rangle_{\bar{\mu}} = \bar{\alpha} \langle J\bar{\chi}', \delta_{\dot{\gamma}}\chi' - \delta_{\bar{\mu}}\chi' \rangle_1 + \langle J\bar{\chi}', \delta_{\dot{\gamma}}\chi' - \delta_{\bar{\mu}}\chi' \rangle_2 = 0, \end{aligned}$$

and therefore also the relations

$$\begin{aligned} \langle \chi', \delta_{\dot{\gamma}}\chi' \rangle_2 &= \alpha \langle \chi', \delta_{\dot{\gamma}}\chi' \rangle_1 + \langle \chi', \delta_{\dot{\gamma}}\chi' \rangle_2 = \langle \langle J\chi', \delta_{\dot{\gamma}}\chi' \rangle \rangle_{\bar{\mu}} = 0 \\ \langle J\bar{\chi}', \delta_{\dot{\gamma}}\chi' \rangle_2 &= \bar{\alpha} \langle J\bar{\chi}', \delta_{\dot{\gamma}}\chi' \rangle_1 + \langle J\bar{\chi}', \delta_{\dot{\gamma}}\chi' \rangle_2 = -\langle \langle \bar{J}\chi', \delta_{\dot{\gamma}}\chi' \rangle \rangle_{\bar{\mu}} = 0. \end{aligned}$$

These arguments carry over to $J\bar{\chi}'$ and show that there exists a variation $\delta_{\dot{\gamma}}J\bar{\chi}'$ such that the relation

$$\begin{aligned} \langle \chi', \delta_{\dot{\gamma}}J\bar{\chi}' \rangle_1 &= 0 & \langle \chi', \delta_{\dot{\gamma}}J\bar{\chi}' \rangle_2 &= 0 \\ \langle J\bar{\chi}', \delta_{\dot{\gamma}}J\bar{\chi}' \rangle_1 &= 0 & \langle J\bar{\chi}', \delta_{\dot{\gamma}}J\bar{\chi}' \rangle_2 &= 0 \end{aligned}$$

hold. Since χ is a linear combination of χ' and $J\bar{\chi}'$ there exists a variation $\delta\chi$ with the desired properties. **q.e.d.**

5.3 Constrained Willmore tori

An immersion, which is a stationary point of the Willmore functional is called a Willmore torus. The following terminology can be found for example in [Wi-2, 7.10]:

Constrained Willmore tori 5.8. *Variations of immersions, which do not change the conformal class, are called conformal variations. An immersion, which is a stationary point of the Willmore functional with respect to all conformal variations of the immersion is called a Constrained Willmore torus.*

Obviously all Willmore tori are *Constrained Willmore tori*. In this section we want to prove the following

Theorem 5.9. *Let U be the potential of a Constrained Willmore torus 5.8. Then the normalization of $\mathcal{F}(U, U)/\Lambda^*$ has a meromorphic function f with finitely many poles at the preimage of the element $[\kappa/2]$ described in the Singularity condition 5.3. Moreover, for arbitrary small $\delta' > 0$ there exists some $\delta, \varepsilon > 0$, such that f maps the open set $\mathcal{V}_{\varepsilon, \delta}^+$ into $\{z \in \mathbb{C} \mid |z+1| < \delta'\}$ and the set $\mathcal{V}_{\varepsilon, \delta}^-$ into $\{z \in \mathbb{C} \mid |z-1| < \delta'\}$. Finally, the function f is invariant under σ and equal to $-\rho^*f$.*

Proof. Let U be the potential of a *Constrained Willmore torus* 5.8. Due to Proposition 5.5 there exist finitely many meromorphic functions $f_1, \dots, f_l$ on some neighbourhood $\mathcal{U}$ of the element $[\kappa/2]$ of the complex Fermi curve $\mathcal{F}(U, U)/\Lambda^*$ such that all real variations δU , whose form $\Omega_{U, U}(\delta U, \delta U)$ multiplied with any of these functions has zero residue on $\mathcal{U}$, induce some variation of the immersion. The pullbacks of all these functions to the normalization of $\mathcal{F}(U, U)/\Lambda^*$ are holomorphic in the complement of the preimage of $[\kappa/2]$. Moreover, the

neighbourhood $\mathcal{U}$ may be chosen to be invariant under σ and ρ , and the functions $f_1, \dots, f_l$ can be chosen to be invariant under σ and to satisfy $-\rho^* \bar{f}_i = f_i$, $i = 1, \dots, l$. In fact, since $\Omega_{U,U}(\delta U, \delta U)$ is invariant under σ and satisfies $\rho^*(\bar{\Omega}_{U,U}(\delta U, \delta U)) = \Omega_{U,U}(\delta U, \delta U)$, for all meromorphic functions f on $\mathcal{U}$ the total residue of $f \cdot \Omega_{U,U}(\delta U, \delta U)$ is equal to the residue of $\sigma^*(f \cdot \Omega_{U,U}(\delta U, \delta U)) = \sigma^*(f) \cdot \Omega_{U,U}(\delta U, \delta U)$ and equal to the complex conjugate of the residue of $\rho^*(\bar{f} \cdot \bar{\Omega}_{U,U}(\delta U, \delta U)) = \rho^*(\bar{f}) \cdot \Omega_{U,U}(\delta U, \delta U)$. Now let $\tilde{A}_1(\cdot), \dots, \tilde{A}_l(\cdot)$ denote the corresponding operator-valued meromorphic functions defined in Lemma 3.2 and $\tilde{A}_0(\cdot)$ the constant function defined in (iv) of Lemma 3.4. Due to Lemma 3.4 the corresponding variations are of the form $(-\delta U_0, \delta U_0), \dots, (-\delta U_l, \delta U_l)$, with real variations $\delta U_0, \dots, \delta U_l \in L^2_{\mathbb{R}}(\mathbb{R}^2/\Lambda)$ and with δU_0 being equal to $2U$. The right hand side of the formula (iv) in Lemma 3.2 for the variations $(\delta U, \delta U)$ and $(\delta U_0, \delta U_0)$ is equal to

$$\frac{1}{\pi \sqrt{-1}(\kappa_1 \kappa'_2 - \kappa_2 \kappa'_1)} \int_{\mathbb{R}^2/\Lambda} \delta U \delta U_0 d^2 x = \frac{2}{\pi \sqrt{-1}(\kappa_1 \kappa'_2 - \kappa_2 \kappa'_1)} \int_{\mathbb{R}^2/\Lambda} \delta U U d^2 x,$$

and therefore proportional to the variation of the Willmore functional. Since U is the potential of a *Constrained Willmore torus* 5.8, this integral must vanish, whenever all the integrals

$$\frac{1}{\pi \sqrt{-1}(\kappa_1 \kappa'_2 - \kappa_2 \kappa'_1)} \int_{\mathbb{R}^2/\Lambda} \delta U \delta U_i d^2 x, \quad i = 1, \dots, l$$

vanish. We conclude that there exist some real constants $c_1, \dots, c_l$, such that $\delta U_0 + c_1 \delta U_1 + \dots + \delta U_l$ is identically zero. Now Lemma 3.2 implies that the meromorphic function

$$\tilde{A}(\cdot) = \tilde{A}_0(\cdot) + c_1 \tilde{A}_1(\cdot) + \dots + \tilde{A}_l(\cdot)$$

commutes with the operator-valued function $\tilde{D}_{\tilde{\gamma}, \tilde{\gamma}}(\cdot)$. Thus these two operators can be diagonalized simultaneously, and the eigenvalue of $\tilde{A}(\cdot)$ defines some global meromorphic function f on the *complex Fermi curve*:

$$f = \text{tr} \left(\tilde{P}_{\tilde{\gamma}} \tilde{A}(p) \right),$$

where p is equal to the corresponding coordinate of the variable of $\tilde{P}_{\tilde{\gamma}}$. Property (i) of Lemma 3.2 ensures that this function is invariant under the action of the dual lattice and therefore defines a meromorphic function on $\mathcal{F}(U, U)/\Lambda^*$. The estimate (iii) of Lemma 3.2 and Theorem 2.12 implies the required estimate on f . Due to the definition of f the pullback of the difference $f - c_1 f_1 - \dots - c_l f_l$ to the normalization of $\mathcal{F}(U, U)/\Lambda^*$ is holomorphic on the complement of the preimage of $[\kappa/2]$. The other required properties of f follows from Lemma 3.4. **q.e.d.**

Corollary 5.10. *All Constrained Willmore tori 5.8 are of finite type.*

Proof. For a given $0 < \delta' < 1/2$ there exist positive $\delta, \varepsilon > 0$, such that the meromorphic function $f^2 - 1$ is bounded on $\mathcal{V}_{\varepsilon, \delta}^+ \cup \mathcal{V}_{\varepsilon, \delta}^-$ by $\delta' < 1/2$. Thus, due to the Maximum principle (see

e.g. [Ah, Chapter 4. Theorem 12.]), this bound holds also on all but finitely many handles. But the set $\{z \in \mathbb{C} \mid |z^2 - 1| < 1/2\}$ decomposes into two connected components. Therefore, the normalizations of all but finitely many handles have two connected components. This proves the corollary. **q.e.d.**

This corollary is similar to the finite type theorem of [P-S]. The main difference is that the authors consider from the very beginning immersions with special properties, and which do not have to be double periodic. These properties result in a global meromorphic function on the corresponding Riemann surface. Afterwards they study double periodic solutions, and obtain some restrictions on the Riemann surface. In particular, the Riemann surface has to be of finite genus. In our approach we consider from the very beginning only double periodic immersions. Afterwards we investigate the consequences of the property, that the immersion is a *Constrained Willmore torus* and again obtain the proof of the existence of some global meromorphic function on the Riemann surface. Now, due to the special form of Riemann surfaces corresponding to double periodic solutions, we can also conclude that the Riemann surface has finite genus. The advantage of our approach in the present case is that the global meromorphic function can be of different degree and therefore will result in different properties of the immersions. In fact, our construction shows that the number of poles is related to the dimension of the eigenspace corresponding to $[\kappa/2]$. The results of [G-S-1] show that even in the case of surfaces of revolutions this dimension can be arbitrary large. But Theorem 5.4 suggests that for relative minimizers the degree of f should be equal to four. Moreover, this global meromorphic function should map the complement of that subset of the Riemann surface, which consists of the two points corresponding to infinite energy, into the complement of some finite subset of $\mathbb{P}^1$. Only the existence of such a global meromorphic function will imply that the potentials satisfy some partial differential equation. Otherwise it would be quite complicated to formulate the property, that the *complex Fermi curve* has a global meromorphic function in terms of the potentials. This property of the global meromorphic function is related to the existence of umbilic points, which are excluded in the description of Willmore tori with the help of the sin-Gordon equation (see [Bo, F-P-P-S, B-B]).

It can happen that not only the *complex Fermi curve*, but the whole complex Bloch variety of some Dirac operator can be compactified to some projective variety. Therefore, the question arises, whether the whole complex Bloch varieties of Willmore tori can be compactified to projective varieties. Here we should mention, that the *complex Fermi curve* alone does not determine the whole spectrum. Moreover, the self-adjoint Dirac operator $\begin{pmatrix} U & \partial \\ -\bar{\partial} & U \end{pmatrix}$ has the same *complex Fermi curve* as the similar operator $\begin{pmatrix} \partial & -U \\ U & \bar{\partial} \end{pmatrix} = \begin{pmatrix} U & \partial \\ -\bar{\partial} & U \end{pmatrix} J$, but the associated complex Bloch varieties are completely different. From the very beginning we could have used the second operator. Indeed, the operators $\tilde{D}_{\tilde{\gamma}, \tilde{\gamma}}(p)$, introduced in Section 5.2 are more closely related to the last operator than with the original Dirac operator. The second operator has a remarkable property: the *complex Fermi curves* corresponding to different energies are all isomorphic. In fact, if we multiply an eigenfunction with the function ψ_k , with $k = (\lambda, 0)$, the energy is shifted by $\lambda\pi\sqrt{-1}$. Thus the whole complex Bloch variety does not contain more information than the *complex Fermi curve* and may be compactified

to a projective variety if and only if the *complex Fermi curve* can be compactified to an algebraic curve. This is not the case for the Dirac operator. Also the property, that the whole complex Bloch variety $\mathcal{B}(U, U)/\Lambda^*$ can be compactified to a projective variety, is much more restrictive.

Obviously, the *Periodicity condition* 1.1 makes sense for spinors in the kernel of a *Finite rank Perturbation* 3.22. The corresponding conformal mappings from $\mathbb{R}^2/\Lambda$ into $\mathbb{R}^3$ can have finitely many poles. Moreover, the whole discussion of Section 5.1 carries over to these conformal mappings with poles. We remark that due to the relation of the *Finite rank Perturbation* 3.22 with the *Singularities of the holomorphic structure* 3.26 the closed forms, which were used to define the immersions, have no monodromy around the singular points. However, suitable inversions transform these conformal mappings with poles into conformal mappings without poles from $\mathbb{R}^2/\Lambda$ into $\mathbb{R}^3$. Due to the conformal invariance of the *complex Fermi curves* [G-S-2] these inversions do not change the *complex Fermi curves*. Therefore we may extend the Willmore functional to those *complex Fermi curves* in $\bar{\mathcal{M}}_{\Lambda, \eta, \sigma}$, which obey the *Singularity condition* 5.3. These *complex Fermi curves*, which obey the *Singularity condition* 5.3 are obviously locally zeroes of holomorphic functions on the compactified moduli space $\bar{\mathcal{M}}_{\Lambda, \eta, \sigma}$. Hence we obtain

Corollary 5.11. *The restrictions of the Willmore functional to an arbitrary conformal class has a minimizer of finite type. In particular, the corresponding immersion is analytic. q.e.d.*

Finally, we remark that this result carries over to immersion into $\mathbb{R}^4$. More precisely, for all $k \in \mathbb{R}^2$ the restrictions of the *first integral* to all *complex Fermi curves* in $\bar{\mathcal{M}}_{\Lambda, \eta}$, which obey the *Quaternionic Singularity condition* 1.6 over an element, whose imaginary part is equal to k , has a minimizer. Due to the results of ‘quaternionic function theory’ developed by F. Pedit and U. Pinkall [P-P, F-L-P-P], these functionals are the restrictions of the Willmore functional in $\mathbb{R}^4$ to fixed conformal classes and fixed holomorphic line bundles (underlying the quaternionic line bundle). We expect that similar arguments may show that these minimizers are of *finite type*. Since the space of holomorphic bundles is compact, the restrictions of the Willmore functional in $\mathbb{R}^4$ to arbitrary conformal classes has also a minimizer. Since the whole family constructed in Section 4.3 obey the *Quaternionic Singularity condition* 1.6, for non-rectangular conformal classes the corresponding minimizers in $\mathbb{R}^4$ take lower values than the corresponding minimizers in $\mathbb{R}^3$.

References

- [Ad] R. A. Adams: Sobolev spaces. Pure and Applied Mathematics **65**. Academic Press, New York (1975).
- [Ah] L. V. Ahlfors: Complex analysis, third edition. McGraw-Hill, Inc., Singapore (1979).
- [A-McK-M] H. Airault, H. McKean, J. Moser: Rational and elliptic solutions of the Korteweg-de Vries Equation and a related many-body problem. Commun. Pure and appl. Math. **30**, 95–148 (1977).

- [A-K] S. Albeverio, P. Kurasov: Singular perturbations of differential operators. London Mathematical Society Lecture Note Series **271**. Cambridge University Press, Cambridge (2000).
- [A-C-G-H] E. Arbarello, M. Cornalba, P. A. Griffiths, J. Harris: Geometry of algebraic curves. Volume I. Grundlehren der mathematischen Wissenschaften **267**. Springer, New York (1985).
- [Ba] M. V. Babich: Willmore surfaces, 4-particle Toda lattice and double coverings of hyperelliptic surfaces. Amer. Math. Soc. Transl. **174**, 143–168 (1996).
- [B-B] M. V. Babich, A. I. Bobenko: Willmore tori with umbilic lines and minimal surfaces in hyperbolic space. Duke Math. J. **72**, 151–185 (1993).
- [Ba-St] C. Bănică, O. Stănăşilă: Algebraic methods in the global theory of complex spaces. Editura Academiei, Bucuresti, John Wiley & Sons, London (1976).
- [Bat] A Erdelyi (ed.), W. Magnus, F. Oberhettinger, F. G. Tricome: Bateman Manuscript Project, Higher Transzendentale Funktionen Volume II. McGraw–Hill Book Company, Inc., New York (1953).
- [B-B-E-I-M] E. D. Belokos, A. I. Bobenko, V. Z. Enol'skii, A. R. Its, V. B. Matveev: Algebro-geometric approach to nonlinear integrable equations. Springer Series in Nonlinear Dynamics, Springer, Berlin (1994).
- [B-S] C. Bennett, R. Sharpley: Interpolation of operators. Pure and Applied Mathematics **129**. Academic Press, Orlando (1988).
- [Bo] A. I. Bobenko: Constant mean curvature surfaces and integrable equations. Uspekhi Mt. Nauk **46:4**, 3–42 (1991); English transl. Russian Math. Surveys **46:4**, 1–45 (1991).
- [B-C-R] J. Bochnak, M. Coste, M.–F. Roy: Real algebraic geometry. Ergebnisse der Mathematik und ihrer Grenzgebiete 3. Folge **36**. Springer, Berlin (1998).
- [B-F-L-P-P] F. Burstall, D. Ferus K. Leschke, F. Pedit, U. Pinkall: Conformal geometry of surfaces in S^4 and quaternions. Lecture Notes in Mathematics **1772**. Springer, Berlin, New York (2002).
- [Ca] T. Carleman: Sur un probleme d'unicite pour les sytemes d'equations aux derivees partielles a deux variables independantes. Ark. Mat. Astron. Fys. **B 26** No. 17, 1–9 (1939).
- [Ch] B. Chen: Geometry of submanifolds. Pure and Applied Mathematics **22**. Marcel Dekker, New York (1973).
- [Co-I] J. B. Conway: Functions of one complex variable. Graduate Texts in Mathematics **11**. Springer, New York (1973).

- [Co-II] J. B. Conway: Functions of one complex variable II. Graduate Texts in Mathematics **159**. Springer, New York (1995).
- [DF] A. Defant, K. Floret: Tensor norms and operator ideals. North-Holland Mathematical Studies **176**. North-Holland, Amsterdam (1993).
- [Do] A. Dold: Lectures on algebraic topology. Grundlehren der mathematischen Wissenschaften **200**. Springer, Berlin (1972).
- [D-K-N] B. A. Dubrovin, I. M. Krichever, S. P. Novikov: Integrable systems I. In: V. I. Arnold, S. P. Novikov (eds.) Dynamical Systems IV. Encyclopedia of Mathematical Sciences vol. 4, pp. 173–280. Springer, Berlin, Heidelberg (1990).
- [E-K] F. Ehlers, H. Knörrer: An algebro-geometric interpretation of the Bäcklund-transformation for the Korteweg–de Vries equation. Comment. Math. Helv. **57**, 1–10 (1982).
- [Ei] L. P. Eisenhart: A treatise on the differential geometry of curves and surfaces. Ginn and Company, Boston, (1909).
- [F-K] H. M. Farkas, I. Kra: Riemann surfaces. Graduate Texts in Mathematics **71**. Springer, New York (1980).
- [F-K-T] J. Feldman, H. Knörrer, E. Trubowitz: Riemann surfaces of infinite genus I–IV. ETH Zürich preprints (1993).
- [F-L-P-P] D. Ferus, K. Leschke, F. Pedit, U. Pinkall: Quaternionic holomorphic geometry: Plücker formula, Dirac eigenvalue estimates and energy estimates of harmonic 2-tori. Invent. Math. **146**, 507–593 (2001).
- [F-P-P-S] D. Ferus, F. Pedit, U. Pinkall, I. Sterlin: Minimal tori in S^4 . J. reine angew. Math. **429**, 1–47 (1992).
- [F-R] A. Figà-Talamanca, D. Rider: A theorem of Littlewood and lacunary series for compact groups. Pacific J. of Math. **16**, 505–514 (1966).
- [Fi] G. Fischer: Complex analytic geometry. Lecture Notes in Mathematics **538**. Springer, Berlin, Heidelberg (1976).
- [Fo] O. Forster: Lectures on Riemann surfaces. Graduate Texts in Mathematics **81**. Springer, New York (1981).
- [Fr-1] T. Friedrich: Dirac-Operatoren in der Riemannschen Geometrie. Vieweg Verlag, Braunschweig (1997). engl. transl.: T. Friedrich: Dirac operators in Riemannian geometry. Graduate Studies in Mathematics **25**, American Mathematical Society, Providence, Rhode Island (2000).

- [Fr-2] T. Friedrich: On the spinor representation of surfaces in Euclidean 3-space. *J. Geom. Phys.* **28**, 143–157 (1998).
- [G-K-T] D. Gieseker, H. Knörrer, E. Trubowitz: The geometry of algebraic Fermi curves. *Perspectives in Math.* **14**. Academic Press, Inc., Boston (1993).
- [G-K] S. G. Gindikin, G. M. Khenkin (eds.): Several complex variables IV. *Encyclopedia of Mathematical Sciences* vol. 10. Springer, Berlin, Heidelberg (1990).
- [G-J] J. Glimm, A. Jaffe: *Quantum physics*. Springer, New York 1987.
- [Gr] H. Grauert: Der Satz von Kuranishi für kompakte komplexe Räume. *Invent. Math.* **25**, 107–142 (1974).
- [G-P-R] H. Grauert, Th. Peternell, R. Remmert (eds.): Several complex variables VII. *Encyclopedia of Mathematical Sciences* vol. 74. Springer, Berlin, Heidelberg (1994).
- [Gr-Re] H. Grauert, R. Remmert: Coherent analytic sheaves. *Grundlehren der mathematischen Wissenschaften* **265**. Springer, Berlin, Heidelberg (1984).
- [Gr-Ha] P. Griffiths, J. Harris: *Principles of algebraic geometry*. Pure and applied mathematics. John Wiley & Sons, New York (1978).
- [G-S-1] P. Grinevich, M. U. Schmidt: Period preserving flows and the moduli space of periodic solutions of soliton equations. *Physica D* **87**, 73–98 (1995).
- [G-S-2] P. Grinevich, M. U. Schmidt: Conformal invariant functionals of immersions of tori into $\mathbb{R}^3$. *Journal of Geometry and Physics* **26**, 51–78 (1998).
- [G-H] B. H. Gross, J. Harris: Real algebraic curves. *Ann. scient. Éc. Norm. Sup.* **14**, 157–182 (1981).
- [Gu-Ro] R. C. Gunning, H. Rossi: *Analytic functions of several complex variables*. Prentice-Hall, Inc., Englewood Cliffs (1965).
- [Har] R. Hartshorne: *Algebraic geometry*. Graduate Texts in Mathematics **52**. Springer, New York (1977).
- [H-K-Z] H. Hedenmalm, B. Korenblum, K. Zhu: *Theory of Bergman spaces*. Graduate Texts in Mathematics **199**. Springer, New York (2000).
- [He] S. Helgason: *Differential geometry, Lie groups and symmetric spaces*. Pure and Applied Mathematics **80**. Academic Press, New York (1978).
- [Hi] N. Hitchin: Harmonic maps from a 2-torus into the 3-sphere. *J. Diff. Geometry* **31**, 627–710 (1990).
- [Ho] N. R. Howes: *Modern analysis and topology*. Universitext. Springer, New York (1995).

- [Je] D. Jerison: Carleman inequalities for the Dirac and Laplace operators and unique continuation. *Advances in Math* **62**, 118–134 (1986).
- [Ka] T. Kato: Perturbation theory of linear operators. *Grundlehren der mathematischen Wissenschaften* **132**. Springer, Berlin (1966).
- [Ke] J. L. Kelley: General topology. *Graduate Texts in Mathematics* **27**. Springer, New York (1955).
- [Ki-1] Y. M. Kim: Unique continuation theorem for the Dirac operator and the Laplace operator. MIT Thesis, (1989).
- [Ki-2] Y. M. Kim: Carleman inequalities for the Dirac operator and strong unique continuation. *Proceedings of the AMS* **123**, 2103–2111 (1995).
- [Kis] S. V. Kislyakov: Classical themes of Fourier analysis. In V. P. Khavin, N. K. Nilol'skij (eds.): *Commutative harmonic analysis I. Encyclopedia of Mathematical Sciences* vol. 15. Springer, Berlin, Heidelberg (1991).
- [Kn] A. W. Knap: Representation theory of semisimple groups. *Princeton Mathematical Series* **36**. Princeton University Press, Princeton (1970).
- [Ko] K. Kodaira: Complex manifolds and deformations of complex structures. *Grundlehren der mathematischen Wissenschaften* **283**. Springer, New York (1986).
- [Kon] B. G. Konopelchenko: Introduction to multidimensional integrable equations. Plenum Press, New York (1992).
- [Kr-1] I. M. Krichever: Methods of algebraic geometry in the theory of non-linear equations. *Russ. Math. Surv.* **32** No.6, 185–213 (1977).
- [Kr-2] I. M. Krichever: Elliptic solutions of the Kadomtsev–Petviashvili equation and integrable systems of particles. *Funct. Anal. Appl.* **14**, 282–289 (1981).
- [Ku] P. Kuchment: Flochet theory for partial differential equations. *Operator Theory Advances and Applications* **60**. Birkhäuser, Basel 1993.
- [Kun] E. Kunz: Holomorphe Differentialformen auf algebraischen Varietäten mit Singularitäten I. *Manuscripta Math.* **15**, 91–108 (1975).
- [La] S. Lang: Differential and Riemannian manifolds. *Graduate Texts in Mathematics* **160**. Springer, New York (1995).
- [L-B] H. Lange, C. Birkenhake: Complex abelian varieties. *Grundlehren der mathematischen Wissenschaften* **302**. Springer, Berlin (1992).
- [Law] H. B. Lawson: Complete minimal surfaces in S^3 . *Ann. of Math.* **92**, 335–374 (1970).

- [L-Y] P. Li, S. T. Yau: A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces. *Invent. Math.* **69**, 269–291 (1982).
- [L-McL] Y. Li, D. W. McLaughlin: Morse and Melnikov functions for NLS Pde's. *Commun. Math. Phys.* **162**, 175–214 (1994).
- [Li] J. E. Littlewood: On the mean value of power series. *Proc. London Math. Soc.* **25**, 528–556 (1924).
- [Ma] N. Mandache: Some remarks concerning unique continuation for the Dirac operator. *Letters Math. Phys.* **31**, 85–92 (1994).
- [M-O] V. A. Marchenko, I. V. Ostrovskii: A characterization of the spectrum of Hill's operator. *Math. USSR Sbornik* **26**, 493–554 (1975).
- [M-S] V. B. Matveev, M. A. Salle: Darboux transformations and solitons. Springer Series in Nonlinear Dynamics, Springer, Berlin (1991).
- [Mi] E. Michael: Topologies on spaces of subsets. *Trans. Am. Math. Soc.* **71**, 152–182 (1951).
- [O] R. O'Neil: Convolution operators and $L(p, q)$ spaces. *Duke Math. J.* **30**, 129–142 (1963).
- [Pi-1] U. Pinkall: Hopf tori in S^3 . *Invent. Math.* **81**, 379–386 (1985).
- [Pi-2] U. Pinkall: Regular homotopy classes of immersed surfaces. *Topology* **24**, 421–434 (1985).
- [P-P] F. Pedit, U. Pinkall: Quaternionic analysis on Riemann surfaces and differential geometry. *Doc. Math. J. DMV, Extra Volume ICM 1998, Vol. II*, 389–400 (1999).
- [P-S] U. Pinkall, I. Sterling: On the classification of constant mean curvature tori. *Annals of Math.* **130**, 407–451 (1989).
- [Pr-Se] A. Pressley, G. Segal: Loop groups. Oxford Mathematical Monographs. Oxford University Press, New York (1986).
- [R-S-I] M. Reed, B. Simon: Methods of modern mathematical physics vol. I: Functional analysis. Academic Press, Inc., London (1980).
- [R-S-II] M. Reed, B. Simon: Methods of modern mathematical physics vol. II: Fourier Analysis, Self-Adjointness. Academic Press, Inc., London (1975).
- [R-S-IV] M. Reed, B. Simon: Methods of modern mathematical physics vol. IV: Analysis of operators. Academic Press, Inc., London (1978).

- [Ro-1] H. L. Royden: The Riemann–Roch Theorem. *Comment. Math. Helvetici* **34**, 37–51 (1960).
- [Ro-2] H. L. Royden: Real analysis. 3 rd edition. Prentice–Hall, Inc., Englewood Cliffs (1988).
- [Sch] M. U. Schmidt: Integrable systems and Riemann surfaces of infinite genus. *Memoirs of the American Math. Soc.* **581** (1996).
- [S-T] H. Seifert, W. Threlfall: A textbook of topology. *Pure and Applied Mathematics* **89**. Academic Press, New York (1980).
- [Se] J.–P. Serre: Algebraic groups and class fields. *Graduate Texts in Mathematics* **117**. Springer, New York (1988).
- [Si-1] L. Simon: Existence of Willmore surfaces. Miniconference, Canberra on geometry and partial differential equations. *Proc. Cent. Math. Anal., Australian National University* **10**, 187–216 (1986).
- [Si-2] L. Simon: Existence of surfaces minimizing the Willmore functional. *Commun. in Anal. and Geom.* **1**, 281–326 (1993).
- [So] C. D. Sogge: Fourier integrals in classical analysis. *Cambridge tracts in mathematics* **105**. Cambridge University Press, Cambridge (1993).
- [St] E. M. Stein: Singular integrals and differentiability properties of functions. *Princeton Mathematical Series* **30**. Princeton University Press, Princeton (1970).
- [S-W] E. M. Stein, G. Weiss: Introduction to Fourier analysis on Euclidean spaces. *Princeton Mathematical Series* **32**. Princeton University Press, Princeton 1971.
- [Str] M. Struwe: Variational Methods, *Ergebnisse der Mathematik und ihrer Grenzgebiete* 3. Folge **34**. Springer, Berlin (1990).
- [Ta-1] I. A. Taimanov: Modified Novikov–Veselov equation and differential geometry of surfaces. *Am. Math. Soc. Transl.* **179**, 133–151 (1997).
- [Ta-2] I. A. Taimanov: The Weierstraß representation of closed surfaces in $\mathbb{R}^3$. *Funct. Anal. Appl.* **32**, 258–267 (1998). and differential geometry of surfaces. *Am. Math. Soc. Transl.* **179**, 133–151 (1997).
- [Tk] V. Tkachenko: Spectra of non–selfadjoint Hill’s operators and a class of Riemann surfaces. *Ann. Math.* **143**, 181–231 (1996).
- [Wh] H. Whitney: Complex analytic varieties. Addison–Wesley Publishing Company, Inc., Menlo Park, London (1972).

- [Wi-1] T. J. Willmore: Total curvature in Riemannian geometry. Ellis Horwood Limited, Chichester (1982).
- [Wi-2] T. J. Willmore: Riemannian Geometry. Clarendon Press, Oxford (1993).
- [Wo] T. H. Wolff: Recent work on sharp estimates in second-order elliptic unique continuation problems. *Journal of Geometric Analysis* **3**, 621–650 (1993).
- [Yo] K. Yosida: Functional analysis. 3 rd edition. Grundlehren der mathematischen Wissenschaften **123**. Springer, Berlin (1971).
- [Zh] K. Zhu: Operator Theory in functions spaces. Pure and Applied Mathematics **139**. Marcel Dekker, Inc., New York (1990).
- [Zi] W. P. Ziemer: Weakly differentiable functions. Graduate Texts in Mathematics **120**. Springer, New York (1989).

Index

- accent tilde, 16
- admissible form, 160
- arithmetic genus, 39
 - local contribution to the $\sim$, 156
 - of the $\rightarrow$ structure sheaf, 39
- asymptotic analysis, 27, 139
- Bäcklund transformation, 47–50, 196
- Baker–Akhiezer function ($\rightarrow$ eigenfunction), 43
 - condition $\sim$ (i)–(ii), 52–53
 - divisor of the $\sim$, 45
- Bloch
 - complex $\sim$ variety $\mathcal{B}$, 25
 - variety, 25
- blowing up
 - of the moduli space, 74
- Carleman inequality, 23
- Clifford torus, 3
- compactification
 - of an isospectral set, 51
 - of the moduli space, 80, 139
- condition
 - Baker–Akhiezer function (i)–(ii), 52–53
 - decomposition (i)–(ii), 130–132
 - divisor (i)–(iii), 45
 - divisor (iv), 57
 - finite rank perturbation (i)–(iii), 94
 - form (i)–(iv), 80–81
 - neighbourhood (i)–(ii), 81–82
 - periodicity $\sim$, 5
 - quasi-momenta (i)–(iv), 43–44
 - quasi-momenta (v), 68
 - quasi-momenta (i') and (vi), 144
 - quasi-momenta (ii'), 80
 - quaternionic singularity $\sim$, 7
 - singularity $\sim$, 7, 197
 - singularity (i)–(ii), 102
 - weak singularity $\sim$, 7, 155
- conformal
- class
 - moduli space $\mathcal{M}_1$ of $\sim$ es, 3, 185
 - of an immersion, 3
 - of the lattice Λ , 185
- immersion, 4
- mapping, 8, 10
- connected components
 - of the normalization, 28
- coordinate
 - $\hat{q}$ and $\check{q}$, 15
 - x, z , 15
- curve
 - complex Fermi $\sim \mathcal{F}$, 6, 25
 - Fermi $\sim$, 6
 - generalized Weierstraß $\sim$, 157
 - quaternionic Weierstraß $\sim$, 190
 - spectral $\sim$, 6
 - Weierstraß $\sim$, 6
- cuspidal, 73, 155, 156, 165
 - of the moduli space, 73
- cut
 - horizontal $\sim$, 83
 - modified pair of parallel $\sim$ s, 84
 - pair of parallel $\sim$ s, 83
- Davey–Stewartson equation, 26
- decomposition
 - condition $\sim$ (i)–(ii), 130–132
- deformation
 - of a complex Fermi curve, 68
 - of a complex space, 68
- Dirac operator D , 15
 - eigenfunction ψ of the $\sim$, 29
 - generalized $\sim$, 56
 - perturbation of the $\sim$, 99
 - resolvent of the $\sim R$, 17
- Dirichlet integral, 84
- disconnected
 - normalization, 158, 173
- dividing

- real part, 157
- divisor
 - condition $\sim$ (i)–(iii), 45
 - condition $\sim$ (iv), 57
 - of the Baker–Akhiezer function, 45
- effective
 - dual lattice $\Lambda_{\text{effective}}^*$, 175
 - lattice $\Lambda_{\text{effective}}$, 175
- eigenfunction ($\rightarrow$ Baker–Akhiezer function), 29
 - ψ of the Dirac operator, 29
 - ϕ of the transposed Dirac operator, 29
 - normalization of the $\sim$, 42
- existence
 - of a minimizer, 3
 - of a potential, 44
- family
 - $\mathcal{F}_{\min, [\kappa/2]}(\cdot)$, 178
 - $\mathcal{F}_{\min}(\cdot)$, 170
- Fermi curve, 6
 - complex $\sim \mathcal{F}$, 6, 25
- finite
 - arithmetic genus, 42
 - geometric genus, 42
 - rank perturbation, 99
 - condition $\sim$ (i)–(iii), 94
 - type potentials, 42
- form
 - admissible $\sim$, 160
 - condition $\sim$ (i)–(iv), 80–81
 - regular $\sim$, 34, 36, 58
- Fourier transform F , 21
- fundamental domain, 15
- generalized
 - Cauchy’s integral formula, 100
 - Dirac operator, 56
 - Hölder’s inequality, 120
 - Weierstraß
 - curve, 9, 157
 - potential, 8, 9
 - Willmore functional, 157
 - Young’s inequality, 121
- genus
 - $\rightarrow$ arithmetic $\sim$, 39
 - $\rightarrow$ geometric $\sim$, 39
- geometric genus, 39
 - finite $\sim$, 42
- gluing rule, 84
- Green’s function
 - $\sim \mathcal{G}_\lambda$, 17
 - $\sim \mathcal{G}_{\mathbb{P}^1, \lambda}$, 109
- handle, 27
- Hausdorff metric, 78
- Hilbert bundle, 16
- index set
 - of handles Λ_δ^* , 27
 - of horizontal cuts Λ_δ^* , 140
- integral
 - Dirichlet $\sim$, 84
 - first $\sim \mathcal{W}$, 7, 79, 84
 - kernel
 - of $R(0, 0, k, 0)$, 94
 - of $\tilde{R}(0, 0, 0, \sqrt{-1}\lambda)$, 18
 - of a finite rank perturbation, 99
- intersection number, 168
 - local contribution to the $\sim$, 168
- involution
 - η , 29
 - ρ , 29
 - σ , 29
 - of the complex Bloch variety, 29
 - of the moduli space, 159
- isospectral
 - set, 12
 - compactification of an $\sim$, 51
 - transformation, 31, 59
 - infinitesimal $\sim$, 59
- lattice
 - $\Lambda, \Lambda_{\mathbb{C}}$, 15
 - conformal class of the $\sim \Lambda$, 185
 - dual $\sim \Lambda^*$, 15
 - dual $\sim \Lambda_{\mathbb{C}}^*$, 169

- dual $\sim \Lambda_{[\kappa/2]}^*$, 176
- dual $\sim$ generator
 - $\hat{\kappa}$ and $\check{\kappa}$, 15
- dual sub- $\sim \check{\Lambda}^*$, 176
- effective $\sim \Lambda_{\text{effective}}$, 175
- effective dual $\sim \Lambda_{\text{effective}}^*$, 175
- generator
 - $\hat{\gamma}$ and $\check{\gamma}$, 15
- Lax operator, 6
- local contribution
 - to the $\rightarrow$ arithmetic genus, 156
 - to the intersection number, 168
 - to the Willmore energy $\mathcal{W}_{\text{sing}}$, 55
- Lorentz spaces, 120
- M-curve, 10
- minimizer
 - $\mathcal{F}_{\min}(\cdot)$, 170
 - existence of a $\sim$, 3
 - of the generalized Willmore functional, 180, 189–190
 - relative $\sim$, 157–170
- moduli space, 8, 67
 - $\mathcal{M}_1$ of conformal classes, 3, 185
 - $\mathcal{M}_{\Lambda, \eta}$, $\mathcal{M}_{\Lambda, \eta, \mathcal{W}}$, $\mathcal{M}_{\Lambda, \eta, \sigma}$, $\mathcal{M}_{\Lambda, \eta, \sigma, \mathcal{W}}$, 79
 - $\mathcal{M}_{\Lambda}$, 145
 - $\mathcal{M}_{g, \Lambda, \eta}$, $\mathcal{M}_{g, \Lambda, \eta, \mathcal{W}}$, $\mathcal{M}_{g, \Lambda, \eta, \sigma}$, $\mathcal{M}_{g, \Lambda, \eta, \sigma, \mathcal{W}}$, 79
 - $\mathcal{M}_{g, \Lambda}$, 68
 - $\bar{\mathcal{M}}_{\Lambda, \eta}$, $\bar{\mathcal{M}}_{\Lambda, \eta, \mathcal{W}}$, $\bar{\mathcal{M}}_{\Lambda, \eta, \sigma}$, $\bar{\mathcal{M}}_{\Lambda, \eta, \sigma, \mathcal{W}}$, 139
 - $\bar{\mathcal{M}}_{g, \eta}$, 80
 - $\bar{\mathcal{M}}_{g, \Lambda, \eta}$, $\bar{\mathcal{M}}_{g, \Lambda, \eta, \mathcal{W}}$, $\bar{\mathcal{M}}_{g, \Lambda, \eta, \sigma}$, $\bar{\mathcal{M}}_{g, \Lambda, \eta, \sigma, \mathcal{W}}$, 80
 - blowing up of the $\sim$, 74
 - compactification of the $\sim$, 80, 139
 - cuspidal of the $\sim$, 73
 - tangent space of the $\sim$, 68
- multiple point, 155, 156, 164, 165
- neighbourhood
 - $\mathcal{U}_{\varepsilon, \delta}^{\pm}$, 27
 - $\mathcal{U}_p^{\pm}$, 82
 - $\mathcal{V}_{\varepsilon, \delta}^{\pm}$, 27
 - $\mathbb{P}_p^{\pm}$, 84
 - condition $\sim$ (i)–(ii), 81–82
- normalization
 - connected components of the $\sim$, 28
 - disconnected, 158, 173
 - of finite genus, 42
 - of the eigenfunction, 42
- orbit, 155
 - of type 1, 155, 156, 164, 165
 - of type 2, 155, 156, 164
 - of type 3, 155, 156
- order of zeroes $\text{ord}_z(\psi)$, 122, 128, 129
- ordinary double point
 - s of the free *complex Fermi curve* $(k_{\kappa}^+, k_{\kappa}^-)$, 27
 - at infinity (∞^-, ∞^+) , 44
- parity transformation $\mathcal{P}$, 31
- periodicity condition, 5
 - equivalent form of the $\sim$, 5
- perturbation
 - $\rightarrow$ finite rank $\sim$, 99
- Picard group, 12, 59
- quasi-momenta
 - $\hat{p}$ and $\check{p}$, 15
 - k , 15
 - condition $\sim$ (i)–(iv), 43–44
 - condition $\sim$ (v), 68
 - condition $\sim$ (i') and (vi), 144
 - condition $\sim$ (ii'), 80
- quaternionic
 - complex line bundle, 45
 - function theory, 7–9
 - line bundle, 4
 - singularity condition, 7
 - Weierstraß curve, 190
- real
 - $-\sigma$ -hyperelliptic, 158
 - part, 30
 - dividing $\sim$, 157
- region of periodic lines, 85
- relative minimizer, 157–170
- residue, 59, 79
- resolvent

- $\sim$ formula, 105
 - of the Dirac operator R , 17
 - weak continuity of the $\sim$, 19
- Sérre duality, 12, 59
- sheaf
 - dualizing $\sim$, 32
 - structure $\sim$, 39
- singularity
 - $\rightarrow$ cusp, 155
 - $\rightarrow$ multiple point, 155
 - condition, 7, 197
 - quaternionic $\sim$, 7
 - weak $\sim$, 7, 155
 - condition $\sim$ (i)–(ii), 102
 - of the holomorphic structure, 55, 102
- Sobolev
 - constant S_p , 17
- spectral
 - curve, 6
 - projection
 - P , 32, 34
 - $P_{\hat{\gamma}}$, 36
 - $\tilde{P}_{\mu}$, 40
- structure
 - $\ell_1^+(\Lambda^*)$ – $\sim$, 145
 - differentiable $\sim$, 9
 - sheaf, 39
 - arithmetic genus of the $\sim$, 39
- symplectic form, 60
- tangent
 - space of the moduli space, 68
 - space of the phase space, 12, 59
- Theta function θ_{Δ} , 93
- topology
 - compact open $\sim$, 78
 - finite $\sim$, 78
 - of the moduli space, 78
- trivialization, 16
- unique continuation property, 23–25
- Weierstraß
 - curve, 6
 - generalized $\sim$ curve, 9, 157
 - generalized $\sim$ potential, 8, 9
 - potential, 5, 9
 - quaternionic $\sim$ curve, 190
- Willmore
 - conjecture, 3
 - functional, 3
 - generalized $\sim$, 157
 - torus, 4
 - constrained $\sim$, 4, 201, 202